00070400

NOT TO BE ISSUED
OUT OF THE LIBRARY.



Digitized with financial assistance from the
Government of Maharashtra
on 18 July, 2018

TREATISE

ON THE

VENOM OF THE VIPER;

ON THE

AMERICAN POISONS,

AND ON THE

CHERRY LAUREL, &c.

AND SOME OTHER

VEGETABLE POISONS.

TO WHICH ARE ANNEXED,

OBSERVATIONS ON THE PRIMITIVE STRUCTURE OF THE ANIMAL BODY; DIFFERENT EXPERIMENTS ON THE REPRODUCTION OF THE NERVES; AND A DESCRIPTION OF A NEW CANAL OF THE EYE.

WITH TEN DESCRIPTIVE PLATES.

TRANSLATED FROM THE ORIGINAL FRENCH OF

FELIX FONTANA,

NATURALIST TO HIS ROYAL HIGHNESS THE GRAND DUKE OF TUSCANY, AND DIRECTOR OF HIS CABINET OF NATURAL HISTORY.

BY JOSEPH SKINNER,

NAVY SURGEON, AND MEMBER OF THE CORPORATION OF SURGEONS OF LONDON.

70400

711 cv 33

IN TWO VOLUMES.

VOL. I.

LONDON:

PRINTED FOR J. MURRAY, NO. 32, FLEET-STREET.

M,DCC,LXXXVII.

00070400

PREFACE

OF THE

FRENCH EDITOR.

THE Firſt Part of this Work was publiſhed in Italian in 1765. Monſieur Darcet, a celebrated Phyſician at Paris, thought it of ſo much importance, that he tranſlated it into French a ſhort time after. Accidental circumſtances delayed the publication of this Tranſlation. The Authour came to Paris in 1776, and gave M. Darcet ſeveral ſheets of corrections and additions, which were likewiſe tranſlated, and added to the reſt. On the following year a pamphlet appeared at Paris on the volatile alkali *(a)*, in which the nature of the venom of the viper, and the uſe of the volatile alkali againſt the bite of this animal, were treated of. Many things were advanced in this pamphlet, entirely contradictory to what our authour had

* Written by M. Sage.

Written

written more than ten years before in Italy. He fancied he had been deceived, and ſat about making freſh experiments on the ſame ſubject, determined to correct the tranſlation I have ſpoken of, before it was given to the publick.

To this new examination, we owe the ſecond third and fourth parts of this work, throughout the whole of which the moſt delicate experiments are conſpicuous. We may ſay in the ſtricteſt ſenſe, that theſe parts are entirely new, as well on account of the ſubjects of which they treat, as of the diſcoveries they contain.

Before François Redi, no one knew in what the venom of the viper conſiſted. This celebrated naturaliſt employed the greateſt part of his reſearches in confuting the errours that prevailed in his time. His work on the venom of the viper is almoſt entirely calculated to demonſtrate, that the opinions held of this venom were in general falſe: a humiliating reflection to man, who is incapable of attaining the truth, in any other way than by paſſing through errour.

We owe to Redi, what has acquired him the greateſt reputation, the firſt diſcovery of the

the humour that renders the bite of the viper venomous. The experiments he has employed to demonſtrate this diſcovery, are in general very well made, although they have not appeared deciſive to M. Charas, a French chemiſt.

The latter, after having made a great many experiments on the bite of the viper, thought it concluſive that the venom of this animal conſiſts in its rage; or rather, that the ſaliva of the viper, exalted by rage, when the creature is furious and bites, becomes venomous and mortal.

Although this opinion is erroneous, it however carries with it marks of probability; ſince it is certain that a viper is more dangerous and deſtructive in proportion as it is more irritated, as will be ſeen in the courſe of this work. This is what engaged our authour to examine the hypotheſis of ſaliva exalted by rage, by deciſive experiments, and the reader may aſſure himſelf, by peruſing the firſt part of this work, that he has been not a little ſucceſsful in this enquiry.

It is certain that François Redi was miſtaken as to the part where the receptacle of the venom is ſeated, and as to the rout the

venom takes, when the viper in biting communicates it to animals. He was of opinion that this humour is ſituated in the membrane that covers the canine teeth, and that it glides exteriourly along the tooth, and at the ſame time inſinuates itſelf into the animals that receive the bite. We ſee this errour repeated half a century afterwards in James's Dictionary, the author of which beſides adopts all the errours of Mead on the ſaline nature of the venom. If it was not therefore abſolutely neceſſary, it was certainly uſeful, to examine this matter afreſh, and to make it as clear as poſſible.

All the other reſearches our authour has made, properly belong to him, and we may ſay with reaſon, that he has begun where others have ended; or, with greater juſtice, that the whole of his work is new, and truly original.

It appears to me, that one of the greateſt merits of this work conſiſts, not ſo much in the rare and numberleſs diſcoveries it contains, as in the luminous method with which the very important enquiries that are introduced in it are treated. If we are aſtoniſhed at the immenſe number of errours that are

every

every where exploded, we cannot at the ſame time help admiring the tracks, hitherto unknown to obſervers, that our authour has beaten, in examining the ſubject of poiſons.

But that which deſerves the greateſt conſideration, is the very delicate analyſis he has made of the moſt obſcure and complicated enquiries; and the ſagacity with which he has conceived the experiments, that could not avoid leading him to truth. It is to be wiſhed that he may ſerve in future as a model to thoſe philoſophers who enquire into facts, without prepoſſeſſion or prejudice. How many diſputes and opinions will then be terminated! How many truths will be diſcovered! How many errours exploded! And how will the number of books be reduced! *The art of interrogating nature by the means of experiment is very delicate. It is in vain that you combine facts, if theſe facts have no affinity with each other; if they preſent themſelves under an equivocal form; if when they are produced by different cauſes, you are incapable of aſſigning and ſeparating with a certain preciſion, the particular effects that have reſulted from each of theſe cauſes**.

* New Experiments on the Reſiſtance of Fluids, in the Preliminary Diſcourſes of Meſſrs. D'Alembert, Condorcet, and Boſſut.

To judge of what our authour has done in this work, and of what in juſtice belongs to him, the reader ought in the firſt place to examine all the publications of Redi and Mead on the ſame ſubject. I exhort him to do ſo, and this is the greateſt eulogium I can pay to the preſent work. Compariſon, a touch-ſtone that can never deceive us, is all that I demand, or rather, what juſtice and impartiality require.

Let this work be compared with thoſe on the ſame ſubject that have immortalized Redi and Mead. There will be no difficulty in judging how much it ſurpaſſes theirs, whether we conſider the number of diſcoveries, or the variety and multiplicity of experiments: indeed it will ſoon be ſeen, that there is no room for a compariſon betwixt them.

I regard the having found the venom of the viper, to be a gummy ſubſtance, as a real diſcovery. The exiſtence of an animal gum is both important and new.

All that we meet with on the blood, and on the nerves, as they relate to the venom of the viper, is entirely new and original. This may be called a giant's ſtride, that clears a new road to freſh diſcoveries.

But

But this is not all the merit of the preſent work. The treatiſe on the American poiſon called Ticunas, and on ſome other vegetable poiſons; the tracts on the Cherry-laurel; and the experiments on ſome other vegetable ſubſtances, are a new field of enquiry, in which the diſcoveries and induſtry of our authour are very conſpicuous. However, when in conſequence of ſo many rich diſcoveries we think that nothing is any longer hid from us, and that we have at length penetrated into the moſt ſecret receſſes of nature, we meet with labyrinths, out of which it does not ſeem permitted us to extricate ourſelves. Such are the new and important conſequences of the obſervations on the poiſon of the Cherry-laurel; a ſubject ſtill obſcure, but intereſting, which will give occaſion to the reſearches of future obſervers.

The excellent experiments our author has made, relatively to the action of poiſons on the nerves, have occaſioned him to enrich this work with ſeveral very important enquiries into the ſtructure of the nerves themſelves; an obſcure matter, in which nothing was known, and in which it ſcarcely ſeemed allowed to man to make any advances. In

the hand of our authour every thing becomes clear, eaſy, and ſimple. I cannot conceive how this double order of bands, how this ſpiral, exteriour, and apparent ſtructure, in the nerves, can have eſcaped the ſight of all our anatomiſts; and I regard the certain knowledge we now have of the firſt elements of the nerves, as one of the fineſt and moſt ſtriking diſcoveries that has ever been made in animal phyſicks; a diſcovery that has eſcaped the ſight of the moſt ſkilful and moſt practiſed obſervers.

After the obſervations of Lewenhoeck, phyſiologiſts and anatomiſts were of opinion that a diſcovery of the laſt diviſions of the nerves could never be attained to; but what did not appear poſſible then, is now unqueſtionably eſtabliſhed, as any one may aſſure himſelf, by following the traces of our authour. We have reaſon to flatter ourſelves that he will ſhortly favour us with his obſervations on the nature and uſes of *the Primitive Nervous Cylinders*. This is all that remains to be known of theſe wonderful organs. He began ſome time ago to buſy himſelf in this enquiry, and what ought we not

not to expect from ſo exact, and ſo penetrating an obſerver!

We are at preſent not only acquainted with the true ſtructure of the nerves, but are much better informed than heretofore of that of the brain.

Our authour has likewiſe examined the internal ſtructure of the retina—it therefore appears that we have ſcarcely any thing more to deſire as to that organ.

But his obſervations do not terminate here. He has developed with the ſame ſucceſs, the ſtructure of the muſcles, and of the tendons. He finds certain characteriſticks to diſtinguiſh theſe two kinds of ſubſtances, as well from each other, as from the nerves.

The firſt organical elements of the nerves, the brain, the muſcles, and tendons, being thus known, our authour paſſes on to the diſcovery of a new and complete ſyſtem of tranſparent, winding, unramified, cylinders, of a much ſmaller ſize than thoſe of the blood, but which are more extended, and in a greater number, than the arterial and venous veſſels. He finds them in the whole cellular ſubſtance, a ſubſtance which penetrates, and compoſes

compoſes, all the organs of the animal machine.

Our authour diſcovers what he calls winding threads, in hairs, in the nails, in the epidermis, and in the bones. He afterwards gives us a detail of obſervations he made on vegetables, in which it appears, that he traces a ſimilar ſtructure. He concludes by a ſeries of excellent obſervations of foſſils, on the ſubject of which he interſperſes a few doubts, that the reader may not be led away by ſimple appearances; and reſerves his ſentiments on this ſubject for another work, which he purpoſes to publiſh under the title of *Microſcopical Obſervations.*——*Obſervations Microſcopiques.*

He terminates his reſearches on the nerves, by relating ſeveral experiments on the reproduction of them,—a very intereſting matter, hitherto unknown to naturaliſts, on which he has thrown all poſſible light.——

To complete this work, I have judged it neceſſary to add a deſcription of a new canal of the eye, diſcovered by our authour more than eighteen years ago, and which he has never publiſhed.—I have taken this deſcription from a letter dated from London, which he

he wrote at the end of the year 1779, to Mr. Adolphus Murray, a celebrated profeſſor of anatomy at Upſal; and have given the part of our authour's letter that relates to this ſubject, in his own words.

We cannot avoid being ſurpriſed at the little value our authour places on his own diſcoveries, whilſt any other anatomiſt, however celebrated, would have haſtened to publiſh them. At the end of eighteen years, he ſcarcely permits this new canal he has diſcovered, to be announced in a few lines in one of his works; at the ſame time that it has been demonſtrated at Vienna for upwards of ten years, in the common courſes of anatomy, to the profeſſors of which it was probably communicated by M. Brambilla, ſurgeon to the Emperor, and director of the military hoſpitals. Our authour ſhowed this canal to M. Brambilla, when he accompanied his Imperial Majeſty, in his travels in Italy.

Although our authour has never publiſhed this ſo long diſcovered canal, he has, however, ſhown it to a great number of his friends, and to ſeveral other perſons. — M. Murray, the profeſſor of anatomy at Upſal, in a letter he addreſſed to our authour ſome time ago, in-

forms him, that the deſcription of his new canal of the eye has been inſerted in the laſt volume of the Acts of the Academy at Upſal. *In ultimo tomo* (he writes) *deſcriptio canalis a te detecti extat.* — This canal was ſhown by our authour to the Swediſh profeſſor, when this laſt, on his coming to Italy, made ſome ſtay at Florence. At his return to Sweden, he addreſſed our authour at Paris, where he then was, to procure the drawing and deſcription of the canal, which he was deſirous of publiſhing in the Acts of the Swediſh Academy, for the advancement of anatomy, and for the good of his countrymen. The drawing and deſcription were ſent from Paris, but were loſt on the road. Our authour ſent him a freſh copy of them from London, but I do not know whether they ever reached him. I have inſerted the figures and deſcription at the end of this work, with the copy of the letter that accompanied them.

Dr. Troja, a diſtinguiſhed profeſſor at Naples, and member of the royal academy of the ſame city, in a diſſertation he publiſhed in the year 1780 on the diſeaſes of the eyes, ſpeaks of this new canal of the eye, obſerving that the diſcovery of it is due to our authour, who

who ſhowed it him at Paris in the eye of an ox.

It only depended on our authour to give this work a more original and novel air, and even to make it appear in certain reſpects more perfect. He had merely to conceal the ſteps by which he arrived at the truths he has diſcovered, and to be ſilent as to the methods and proceſſes that conducted him to them. The enlightened reader will find, that in proportion as he advances in the peruſal of this work, and meets with unforeſeen difficulties, the experiments our authour has imagined to ſurmount them, will preſent themſelves ſo naturally, that he will almoſt imperceptibly be led to believe, that he imagined them himſelf; ſo ſimple do they appear, and ſo appoſitely are they placed. In the ſame way, the new lights that are interſperſed, and the many reſearches that are deſcribed, ſeem to grow out of the matter itſelf, and not to ſpring from the authour. Again, he might not have ſpoken of whatever remains doubtful or undecided; he might have paſſed over in ſilence, the enquiries he has not been able, even after ſo many experiments, to decide upon. His work might have been found more complete,

becauſe

becauſe ignorance only ſuffers in proportion as it is known; but our authour has throughout preferred clearneſs and utility to vain glorioufneſs and a falſe pride.

There is a claſs of readers to which this work will certainly be unpleaſing, and this claſs is neither the leaſt numerous, nor that which has the feweſt ſectators and partizans. It is compoſed of thoſe pretended naturaliſts who explain nature at their ſcrutoire; who meditating on facts ſeen in a wrong point of view, and copied into books, immediately divine the ſprings of them; and who miſtake for real cauſes, the ideal ones they apply to the explanation of effects which have only exiſted in their own imagination: in ſhort, who prefer romances to facts, and to truth.

To folks of this deſcription, accuſtomed either to read, or to make, romances in phyſicks, the preſent work muſt appear barren, tedious, and little philoſophical; and I therefore cannot exhort them to peruſe it;—but to thoſe, on the contrary, who are fond of facts and certain obſervations, it will be infinitely ſatisfactory. I do not for my own part know of any ſubject, either in phyſicks or in medicine, that has been diſcuſſed with a greater

abundance and richneſs of experiments than the one before us.

When a work is founded on certain and new facts, we always gain by reading it, even though it is badly contrived and preſented, and contains falſe reaſonings. The new truths that are found in it are real acquiſitions to the philoſopher, and he may eaſily employ them as a baſis to truer ſyſtems, and to ſurer opinions, and laſtly, to the diſcovery of the true laws of nature.

But what confidence ought not an authour to inſpire us with, who, after having ſaid, *I have made more than* 6000 *experiments*; *I have had more than* 4000 *animals bit*; *I have employed upwards of* 3000 *vipers*; finds no difficulty in adding, *I may have been miſtaken, and it is almoſt impoſſible that I have not been miſtaken!* — What a difference betwixt this authour and many others! betwixt opinion and certainty! betwixt ignorance and knowledge!

This work, ſo enriched by the immenſe number of new facts, and by the length and difficulty of the reſearches it contains, could not have been executed without the protection and conſtant favours of the auguſt Mæcenas the

the authour has the happineſs to ſerve. But whilſt the bounties of a philoſophick ſovereign procure to the enlightened world ſo many experiments and diſcoveries, the uſe our authour has made of the means that preſented themſelves to him on his journies, will, without doubt, entitle him to the gratitude and admiration of men of letters; and it will never ceaſe to be a matter of ſurpriſe, that a work which has coſt ſo much labour, had its birth at Paris and at London, through which places our authour, if I may ſo expreſs myſelf, did nothing more than paſs.

CONTENTS

CONTENTS

OF THE

FIRST VOLUME.

PART THE FIRST.

INTRODUCTION—*In which is ſhown how little authours agree among themſelves on the venon of the viper* - - - 1

CHAP. I.—*Number, ſtructure, and uſe, of the teeth of the viper* - - - 5

CHAP. II.—*Of the yellow liquor that flows from the tooth of the viper* - - - 12

CHAP. III.—*Of the part where the reſervoir of this yellow humour is ſeated* - - 19

CHAP. IV.—*The venom of the viper is no other than the yellow humour that flows from the tooth when the viper bites* - - - 24

CHAP. V.—*The venom of the viper is not a poiſon to the viper itſelf* - - - 27

CHAP. VI.—*The venom of the viper is not a poiſon to every ſpecies of animals* - - 34

CHAP. VII.—*The venom of the viper is not acid* 43
CHAP. VIII.—*The venom of the viper is not alkaline* - - - - 45
CHAP. IX.—*No ſalts are diſcovered in the venom of the viper* - - - - 47
CHAP X.—*The venom of the viper has no determinate taſte, and when put on the tongue cauſes no inflammation* - - - 54
CHAP. XI.—*Properties of the venom of the viper* 61
CHAP. XII.—*Peculiarities of the venom of the viper, and of that of the venomous animals* 63
CHAP. XIII.—*What cauſes the death of animals that have been poiſoned by the viper* - 74

PART THE SECOND.

CHAP. I.—*On the ſource of many errours* 119
CHAP. II.—*Whether the volatile alkali is a certain remedy againſt the bite of the viper* 131
CHAP. III.—*Of the effects of the bite of one, or ſeveral vipers, on the ſame part of an animal, or on two correſponding parts of the ſame animal* 138
CHAP. IV.—*Of the effects of the bite of the viper on different parts of an animal* - 177
CHAP V.—*Experiments on the comb, gills, noſe, and neck, of animals* - - 202
CHAP. VI.—*Experiments on the tendons* 220
CHAP. VII.—*On the nature of the venom of the viper. Deſcription of certain parts of the head of the viper, that relate to the venom* - 233

PART THE THIRD.

CHAP. I.—*Action of the venom of the viper on parts of an animal that have been bit* - 271
CHAP. II.—*Of the time it requires for the effects of the venom of the viper to become ſenſible* 288
CHAP. III.—*On the action of the venom of the viper on the blood of animals* - 312
CHAP. IV.—*Experiments on the nerves* 330
CHAP. V.—*Effects of the venom of the viper on blood expoſed to the open air* - 376
CHAP. VI.—*On the cauſe of the death of animals bit by the viper* - - - 395
INDEX.

PHILO-

PHILOSOPHICAL RESEARCHES

INTO THE

VENOM OF THE VIPER.

PART I.

INTRODUCTION.

In which is ſhown how little Authours agree among themſelves on the Venom of the Viper.

IT is agreed at preſent that there is no other guide in a ſearch into natural truths, than a knowledge of facts; it is only on facts that the philoſopher can hope either to eſtabliſh a reaſonable ſyſtem, or to form a ſound judgement of thoſe already eſtabliſhed. Obſervation is alone capable of diſſipating the miſts that envelop the hidden cauſes of the phenomena

of nature. And laſtly, it is to the labours of obſervers that we owe the rapid progreſs philoſophy has made in our time. But nothing retards this progreſs more, than the little agreement found amongſt authours, even in matters of experiment; that is to ſay, in things that are expoſed to the finger and to the eye. Nothing is more common than to ſee obſervations of this kind, that are nevertheleſs made by men full of candour, frequently belied by others, or in contradiction with themſelves. What then is the cauſe and ſource of theſe errours? Is it the ſpirit of party, or is it the difficulty of nice obſervation? Be it what it will, it is not leſs true, that after having conſulted the moſt celebrated authours on any particular we wiſh to clear up, we often find ourſelves as little informed, and in as great a ſtate of uncertainty, as we were before. In ſuch caſes then, I have apprehended, that without being wanting in the reſpect due to the authority of theſe great men, I might ſolely truſt to my own eyes; and to render my inveſtigations more deciſive, I have endeavoured by a ſingular application, to diſcriminate nicely, to compare the experiments of my predeceſſors with my own, to trace and develope all the circumſtances of them, and in ſhort, to diſcover what may have occaſioned ſo great a variety in the opinions of theſe obſervers, and in their manner of ſeeing.

Such is the true motive that has induced me to give an account of the experiments which follow. Without this I ſhould willingly have paſſed them

over

over in silence, that I might not fatigue the reader, by presenting him with what others may have already published.

The ensuing experiments relate to the viper, and dwell much less on the anatomy and particular structure of some of its parts, that on the nature of the venom of that animal. The facility with which vipers are procured at Pisa, where I made my experiments, has enabled me to multiply my researches, and to vary them exceedingly. It would be losing time to have no other object in view than that of rooting out the popular prejudices on this subject that were so very prevalent in the time of Redi. We are indebted to this authour both for his having made them known, and having disencumbered natural history of them. He himself knew the value of time, since at the conclusion of his letter to Magalotti he says, *che il perder tempo a chi più sa, più spiace. That the more any one is instructed, the more he regrets lost time.*

When I observed that the frequently repeated observations of so celebrated a writer as Mead, clashed entirely with those of Redi, I must confess that a glimpse of the utility of making known the source of the errours of these two great men, and the pleasure of discovering new truths, encouraged me exceedingly to this undertaking, notwithstanding the risk that attends the handling such dangerous animals.

I have deemed it necessary to begin by making a few remarks on the teeth of the viper, and on some

other of its parts, and if in doing this I repeat certain truths that other obſervers have already publiſhed, it is only to give a greater perſpicuity to my work, and the impartial reader, particularly when he ſees that theſe truths are better eſtabliſhed, and that the experiments which ſerve as a baſis to them have been varied in ſo many ways, will readily pardon me.

CHAPTER I.

Number, Structure, and Use, of the Teeth of the Viper (a).

A GREAT deal has already been written on the large or canine teeth of the viper; they had been examined, even with the microscope, before the time of Redi, and were found to be hollow and tubulated to their very points; Redi made himself fully certain of this with the naked eye, and found that if they were bruised when dry, they broke into three or four pieces, from the basis to the point, and plainly showed their internal cavity; but he flatly denies that this cavity is a conduit for the yellow liquor, and that when the viper bites, this venom gushes from the small hole at the extremity of the tooth. He says that he has opened the mouths of vipers, and has always found that this yellow liquor, when they bite, runs along the outer

(a) *Note of the French Editor.* That the parts described in this chapter may be more readily understood, we have borrowed several figures of the head of the viper from Mead's works: see Table I, and the explanation that precedes it. We request the reader to likewise give a glance at Table II, before he goes any further.

part of the tooth from the top to the bottom, and that it never flows from within. *I have aſſured myſelf well of this, continues Redi, by ſeveral experiments, and by the often repeated teſtimony of my own eyes.*

The celebrated Valiſnieri adds moreover, that the canine teeth of the viper are pierced with four very ſmall lateral holes, through which he believes that the moſt ſubtile part of the venom penetrates into the wound from within the tooth, whilſt the thicker and groſſer part of it runs along the outer ſurface. Mead and Nicholls, on the contrary, ſet out from the analogy betwixt the viper and the rattle-ſnake, in the laſt of which this humour is very diſtinctly ſeen to flow from the inner part of the tooth, and maintain that the venom of the viper flows likewiſe from the point of its canine teeth, or at leaſt from an opening they have towards their extremity. I have ſeveral times repeated Redi's experiments, by opening the mouths of theſe animals when living, and acknowledge that I have never been able to aſſure myſelf whether this venomous liquor iſſued preciſely from within the tooth, or whether it ſimply glided along the outer part, from the baſis to the point. If I held the head of the viper with the point of the teeth downwards, I had only to preſs ſtrongly againſt the muſcles of the palate, to make this yellow liquor flow rapidly from the baſis to the point of the tooth; but if I held it with the teeth turned upwards, I ſaw the poiſon collect inſtantly about the baſis of the tooth, and entirely fill the ſheath or bag that ſerves to encloſe it. Redi maintains

tains moreover, that this ſheath is the true reſervoir in which this humour is depoſited and preſerved, and thinks it is ſecreted by a ſmall neighbouring gland ſituated below the orbit. Nicholls ſays, on the contrary, that there is a veſicle, or ſmall bag, ſeparated from the ſheath, and that this gland is deſtined to quite another purpoſe, as the ſecreting ſome lymphatick or ſalivary humour.

In this uncertainty, I conceived that the beſt ſtep would be to examine by my own proper obſervation, the ſtructure of the viper's teeth, and ſo inform myſelf well of their uſes, ſince the deſcriptions theſe authours give of them are obſcure, and the obſervations of Mead and Nicholls contradictory to thoſe of Redi.

At each ſide of the anteriour and upper part of the head, the viper has a moveable bone which forms in part the upper jaw; each of theſe two bones has two ſockets at the ſide of each other, which are only ſeparated by an immoveable, but very brittle lamen, the ſubſtance of which is ſpongy and like that of the bones themſelves: in theſe ſockets are fixed the canine teeth, which are ſometimes found to the number of four, ſeldom of three, and oftener again of two. It is obſerved, that when theſe teeth are four in number, they are not all fixed in their ſockets with the ſame ſtrength and ſtability: there are at that time uſually two, or one at leaſt, moveable, and capable of being eaſily pulled out without breaking: this cannot be done with the others, which are never pulled out entire, notwithſtanding they

have no roots like ours. I have ſometimes found three moveable ones. I have likewiſe ſeen ſome vipers that had only two canine teeth, both of which were however weak and moveable; but this is a very rare caſe.

At the baſis of theſe large teeth, and quite out of the ſockets, ſix or ſeven very ſmall teeth are invariably found; they ſometimes even amount to the number of eight. When they are examined attentively with a magnifying glaſs, they are found to be faſtened at their baſis by a kind of web of a fine and ſoft membranous texture. Theſe ſmall teeth diminiſh in ſize, in proportion as they are more diſtant from the ſockets of the canine teeth; thoſe which are neareſt the ſockets are likewiſe the hardeſt and beſt formed. The other ſmaller ones are more tender, more imperfect, and as it were mucous, particularly at their baſis; they ſeem in reality to owe their formation to a whitiſh and gelatinous matter.

Beſides the two kinds of teeth I have mentioned, the viper has ſtill another order of them, much more minute than the former, and reſembling ſmall hooks; they are ſtrongly fixed to the number of ten, eleven, and ſometimes fifteen, in two ſmall, pretty long, parallel, bones, which on each ſide form the upper jaw; and of eight, nine, and ſometimes twelve, in each of two bones that form the lower jaw.

The canine or large teeth, as well as the other ſmaller ones found at their baſis, are encloſed in a bag or ſheath, which covers them on all ſides, and is compoſed

poſed of very ſtrong fibres, and of a cellular web. It is however open towards the point of the tooth, and terminates there by folding together its two lamina, which at their junction are often dentated. This ſheath ſeems to be a prolongation of the external membrane of the palate.

The canine tooth is ſeldom more than three lines in length, Paris meaſure *(a)*. Its baſis is not more than half a line in diameter; its figure is that of a horn a little flattened, and ſomewhat bent towards its baſis.

This tooth terminates in a very ſharp point, towards which it inſenſibly loſes its curvature, and becomes at length almoſt ſtraight. Below the middle of the tooth, towards its point, and in the convex part, a ſmall opening is viſible to the naked eye, very narrow but exceedingly long, which ending in a ſmall, channelled ſlope, that can ſcarcely be ſeen with a microſcope, terminates in this way at the point. Hairs plucked from the whiſkers of foxes, cats, dogs, &c. may be readily introduced into this opening, and it appears, when viewed with a microſcope, to be a cleft, the length of which is almoſt a fourth part of that of the tooth, and the breadth the ſixteenth part at moſt. It repreſents with its exterior edge a very long or flattened ellipſis, becoming larger towards the baſis of the tooth. This ſlope

(a) The French, or Paris inch, is ſomewhat larger than the Engliſh one, but not ſo much ſo as to make a ſenſible difference in the line, which is its twelfth part---Therefore whenever the latter is introduced in this work, it may be conſidered as the twelfth part of an Engliſh inch.

pene-

penetrates interiourly, and is terminated at its two ſides by two ſhort edges or lips, thick and raiſed up. Another opening is likewiſe found in the convex part of the tooth, towards the baſis, and near the place where it is fixed in the ſocket. This opening likewiſe begins by a ſhallow cavity, immediately at leaving the ſocket. It is much larger than the firſt, but not longer. In proportion as this ſlope or ſmall hollow penetrates into the tooth, it pierces it for its whole length, and forms a channel which terminates in an elliptical hole at the point. A bit of ſilk is eaſily paſſed from one opening to the other, particularly when care is taken to introduce it at the baſis, where the natural entry of this paſſage is found. The ſide of the ſecond opening reſembles a parabole, the baſis of which paſſes over the bony edges of the ſocket, and the other ſides of which end in a ſomewhat obtuſe point towards the ſummit of the tooth. The canine tooth of a viper then is hollow and tubulated for its whole length, from the baſis to the point, and has two holes in its convex part. This hollow is however not ſuch as one would ſuppoſe it to be, on viewing the third figure of Mead, and the deſcriptions of Redi.

The tooth of the viper has a double pipe or tubule almoſt for its whole length, a circumſtance hitherto entirely unknown to obſervers. Theſe two canals or tubes do not communicate with each other, and are ſeparated by a bony partition, very brittle towards the baſis, but which becomes ſomewhat ſtronger in proportion as it advances towards the point. One

of theſe tubes or canals, which I call the external one, becauſe it is at the ſide of the convex part of the tooth, begins, as has been ſeen, at the baſis of the triangular opening, and goes on enlarging by degrees to the middle of the length of the tooth, whence it gradually narrows, and ends at the elliptical opening of the point. The inner canal, on the contrary, which is towards the concave part of the tooth, begins with a large opening at the baſis, from whence it advances, cloſing by degrees, and terminates at length in a blind point above the middle of the tooth. The partition likewiſe that ſeparates theſe two cavities is crooked, and it's convex part is turned towards the hollow of the canal it terminates, ſo that its bony ſubſtance rather preſents an irregular, curvilinear figure, or truncated cone, than a perfect cone itſelf. The blind canal communicates with the ſocket in which the tooth is fixed, and receives veſſels and nerves, which enter by a ſmall oval hole, perceptible to the eye; and opening at the edges of the ſocket itſelf, towards the inner part of the jaw. This bone of the jaw has likewiſe a large round opening, which begins a canal placed a little below and at the ſide, and opening one way into the ſocket, and the other laterally and more below, towards the ſurface of the ſame part of the jaw.

The ſmall teeth placed at the baſis of the large ones, reſemble them very much both in their inner and outer ſtructure. Thoſe particularly that are placed the neareſt the firſt, and which are the firmeſt,

are

are perfectly like them, unleſs it be that their baſis is not ſo well finiſhed. Like the large ones, they have all the elliptical hole towards the point, and a part of the triangular hole at the baſis. The two conduits, internal and external, are alſo ſeen in them.

It is not the ſame with the other very ſmall teeth I have mentioned, which are far the moſt numerous, and in both jaws. Theſe are not channelled, and have no kind of opening, either at the point or baſis.

CHAPTER II.

Of the yellow Liquor that flows from the Tooth of the Viper.

WHEN the viper wiſhes to bite, its canine teeth are raiſed by a mechaniſm, which Nicholls has deſcribed perfectly well in the anatomical appendix he has annexed to Mead's work on poiſons. But thoſe of the large teeth that are not ſo well faſtened in their ſockets, are leſs capable of raiſing themſelves, in proportion as they are more moveable, and badly fixed in the jaw. Nicholls pretends, that when one or two of theſe four large teeth are moveable, the viper can only bite with one tooth on each ſide. Indeed he does not found his opinion on any experiment,

riment, but ſeems to account for this by certain final cauſes which I cannot admit, ſince in phyſicks theſe kind of proofs become of no weight. He remarks that there is ſuch a diſtance betwixt the two canine teeth of the rattleſnake, that the yellow fluid, which is carried by a conduit betwixt them, would entirely enter into the ſheath, inſtead of being conveyed to the animal bit by this ſnake: on this account he does not heſitate to believe that the conduit of this liquor is preciſely fixed oppoſite the hole at the baſis of the ſingle tooth on each ſide, with which the viper ſeizes what it bites. But beſides that no organs are ſeen to execute this function, and that the mechaniſm of it cannot be inveſtigated, I can take upon me to ſay that I have ſometimes ſeen all the four canine teeth of a viper equally firm and ſtrongly fixed in their ſockets, and have ſtill more frequently found three of them, well fixed and very ſtrong, in a ſtate both to ſeize and bite. It is not to be doubted but that the viper, inſtead of ſimply biting with two teeth, one at each ſide, muſt ſeize equally with all thoſe that are firmly fixed in the ſockets, and I have even aſſured myſelf of this by experiment. It is not true then, as Nicholls pretends, that the conduit of this yellow liquor only adapts itſelf to one ſingle tooth on each ſide when the viper bites; beſides, this ſpace which is obſerved betwixt the canine teeth of the rattleſnake, is not alike found in our vipers, the teeth of which, almoſt from the baſis to the point, touch and embrace each other in ſuch a way, that no fluid can paſs betwixt them, much leſs this yel-

low venomous liquor which is ſomewhat glutinous. It is moreover determined that the viper bites and ſeizes not only with the teeth that are fixed in their ſockets, but likewiſe very often with thoſe that are moveable. Of ten vipers I examined, there were three that had two moveable teeth, and two firm in their ſockets; the ſeven others had only one moveable tooth, and two firm and well faſtened. If I except one of the firſt three vipers, and two of the ſeven laſt, all the others to which I held a bit of tendon of beef, boiled and well ſtripped of its coat, ſeized it forcibly, and left the marks of their teeth ſtrongly printed in it: I muſt however obſerve that their leaſt firm teeth were not of the moſt moveable kind, and that when they are very looſe I have found them to riſe ſo little that it is abſolutely impoſſible for their points to touch the body ſeized by the viper.

Since it is certain that this creature never bites without a riſk of loſing ſome of its teeth, Nicholls conjectures with great ſagacity, after Redi, that nature has intended the ſmall ones at the baſis of the others, to replace, when there is a neceſſity for it, thoſe that the viper loſes from time to time. Their crooked ſhape renders it difficult for them to be drawn from the wound, and in the courſe of my reſearches I have ſometimes obſerved, that not only thoſe which are moveable, but even the firmeſt of them, are alike ſubject to accident. The thinneſs of the tooth, and the ſtrength of the animal that has been bitten, contribute equally to this loſs; and this opinion becomes ſtill more probable when we conſider that theſe ſmall moveable teeth are ex-

actly

ſtly formed like the canine ones, that is to ſay, that they have likewiſe two canals, (thoſe however that are the moſt perfect) and the ſame openings at their baſis and at their point. But all theſe reſemblances were in ſhort but one reaſon more why experiment ſhould be conſulted, and the truth eſtabliſhed by nice obſervations.

I have ſometimes obſerved in one of the ſockets, a very moveable tooth, the ill-formed and ſtill gelatinous baſis of which faſtens itſelf to the edges or ſides of the hollow; this tooth may even be drawn a little way out of the ſocket without detaching it entirely, by means of a tender and mucous matter that ſerves as a glue to it. On moving the jaws, I made the one next it raiſe itſelf very well, but the one of which I have ſpoken, abſolutely continued reclined on the baſis of the moveable bone of the jaw. It is clear that this tooth had been of the number of thoſe that are at the baſis of the great or canine ones.

I expreſsly drew from a large viper one of theſe laſt, which was moveable and ill fixed in its ſocket, and obſerved ſome time after that the largeſt of thoſe which are placed beneath the ſheath and beneath the ſocket, had advanced a little towards the empty ſocket. Some days after I thought I perceived it to have approached ſtill nearer. I purſued my obſervations on every ſecond day, and at length ſaw that this tooth had perfectly fixed itſelf in the ſocket, where however it was as yet very moveable, and badly faſtened. At the end of thirty days it was fixed in ſo ſolid a way as to be capable

of

of biting. The neceſſity one is under of frequently handling the viper to be ſatisfied of the ſtate of its teeth, and of opening its ſheath with pincers, or with the blunted point of a bit of wire, makes this experiment very dangerous. The repeated compreſſions the ſmall teeth receive, from the contraction of the muſcles of the jaw, and the action of the ſheath itſelf which conſtantly preſſes on the points of thoſe teeth that are moſt raiſed, are quite ſufficient to puſh the root of the tooth in queſtion, into the ſocket which the falling out of the old tooth has left empty.

The laſt or ſmalleſt teeth are certainly not employed in biting, but are intended to draw nearer to the throat, and to hold firmer, the animal the viper has already ſeized.

The ſingular ſtructure of the canine teeth, ſo different from that of the other teeth of the two jaws, is a powerful perſuaſive of its being from them that the yellow liquor flows; it is not however without ſome apparent reaſon that Redi, otherwiſe ſo exact, has been led into an errour.

To fully aſſure myſelf of this circumſtance, I bound the head of a viper I had juſt killed, to a table. To diſtinguiſh better and to a greater certainty, I took the precaution to remove the lower jaw: the canine tooth, in the way I had fixed the head, was turned upwards, and I obſerved the elliptical cleft with the ſtrongeſt lens of Ellis's microſcope. I gently compreſſed the palate with a ſomewhat blunted iron, when a ſlightly tranſparent yel-

low

low liquor, which formed itſelf into a drop, and at length glided along the outer ſurface of the tooth, inſtantly appeared at the elliptical hole of the point. I repeated this experiment ſeveral times, and always with the ſame ſucceſs. I afterwards cloſed the ſmall opening with wax, and then compreſſed the palate, but not a particle of the venom ſhowed itſelf. I however ſaw it through the tranſparent ſides of the tooth, conveying itſelf from the baſis to the point by the external canal, which it had entirely filled. I put a round bit of wax with riſing edges all about the tooth, in ſome other heads of vipers, immediately below the elliptical hole, and having ſtrongly compreſſed the palate, I inſtantly ſaw the yellow liquor guſhing forcibly, and as it were by ſtarts, from the point, and ſcattering itſelf abundantly on the piece of wax, which it entirely covered all round the tooth.

I have likewiſe been able, although with difficulty, to cloſe the hole at the baſis with wax, and I have then found it to be in vain that I compreſſed all the muſcles of the head ſucceſſively. I could never force a drop of venom from the point, nor could I even diſcover it through the ſides of the tooth. Whenever a viper is held in the hand with the teeth turned upwards, it is eaſy for an attentive eye, accuſtomed to obſervation, to ſee this drop of yellow liquor preſent itſelf at the elliptical opening, in ſuch a way that it may be more or leſs encreaſed at will. I have repeated this experiment a thouſand times, and have invariably ſeen the ſmall drop of venom

exuding from the elliptical hole of the tooth. Nay, what is more, on compreſſing violently, this liquor is ſometimes obſerved to force itſelf out ſuddenly, and to ſpirt to a conſiderable diſtance. It muſt however be remarked, that when the tooth is once wetted with it, particularly when it is entirely covered with the ſheath, this humour, or the drop it forms, glides ſo very precipitately along the tooth, that it is ſuddenly ſeen at the baſis without having been perceived at the point. It in this way imperceptibly fills the ſheath, ſo that it is difficult to perſuade ones-ſelf that it really iſſued from the point of the tooth. This is the way that ſo exact an obſerver as Redi has been led into an errour. It is not proper, after his example, to employ living vipers, nor to open their mouths forcibly, ſince the flowing out of the venom will then be too ſudden, and ſince it will then be dangerous to obſerve as nearly as is neceſſary to prevent being deceived.

I not only ſaw the yellow liquor flow from the point of the tooth that I particularly obſerved, but likewiſe from the neighbouring tooth, when there was one, ſo that it proceeds equally from all the canine teeth at a time, not excepting thoſe which, without being altogether firm in their ſockets, are however ſufficiently ſo to riſe with the others. In a word, in all the heads of vipers I have obſerved, I have ſeen this humour conſtantly flow from all the canine teeth that raiſe themſelves ſufficiently, on the compreſſion of the muſcles of the palate, and on opening the mouth with a force that would be capable

ble of wounding an animal ſeized by the viper. This ſhows that Nicholls is deceived, when he pretends that the venom only flows from one tooth at a time on each ſide.

CHAPTER III.

Of the Part where the Reſervoir of this yellow Humour is ſeated.

THE yellow liquor iſſues then from the point of the vipers tooth, contrary to the ſentiments of Redi, who regarded the bag or ſheath that envelops not only the canine teeth, but likewiſe the others that are found at their baſis, as the true reſervoir of this venom. This opinion is ſtill again belied by the ſtructure itſelf of this ſheath, which has a large aperture next the cheeks, through which this liquor would inceſſantly flow with the greateſt eaſe; ſo that every time the jaws of the viper ſhould be extended, the venom would be conſtantly ſeen oozing through the extremity of the ſheath, even though the viper ſhould not bite. This is what no one has hitherto obſerved. It is beſides certain, that when this ſheath is opened with ſciſſars, neither this yellow humour, nor any kind of fluid that may have collected there, is found in its cavity.

But ſince, as has been ſeen above, this liquor flows from out the elliptical hole at the point of the tooth,

it muſt neceſſarily be carried to the aperture at the baſis by another conduit than this ſheath, in which it is certain that no remains of venom are found. It will not be difficult, according to this hypotheſis, to diſcover the ſmall veſicle that is really deſtined to contain it.

If after having ſtripped the teeth of this bag or ſheath, a preſſure is made on the palate, this humour is obſerved to flow through a ſmall and almoſt imperceptible hole, placed at the anteriour part of the maxillary bone, within the ſheath, and at the ſide of the baſis of the canine teeth; ſo that when theſe teeth are covered by the ſheath, this ſmall orifice comes in contact with the inferiour opening of the tooth. Indeed with the help of a magnifying glaſs, a very ſmall orifice is diſcovered, ſituated in the midſt of a ſmall cleft or furrow, which anſwers to the maxillary bone. I endeavoured to introduce into this orifice a fox's hair, very fine, but nevertheleſs pretty ſtrong, and at length ſucceeded in paſſing it quite acroſs the ſheath, by a long membranous conduit, into a ſmall veſicle placed beneath the muſcles of the upper jaw, on its lateral part. It is a membranous bag of a very ſtrong and cloſe texture, which is again partly covered by tendinous fibres. Its ſhape is nearly that of an equilateral triangle. It differs from other veſicles, which are crooked or ſpheroidal, inſtead of which the baſis of this is in a manner ſtraight. This ſmall veſicle terminates next the eye in a tranſparent canal, which after having proceeded beneath the orbit for the

ſpace

ſpace of two lines, pierces the ſheath, and at length opens at the extremity of the ſockets, into the ſmall cleft of which I have ſpoken. When this conduit reaches the vicinity of the ſheath, it dilates a little, and it is here that the venom finds the greateſt obſtacle to its paſſage, which is owing to the compreſſion it meets with from the bones of the jaw.

The veſicle I have ſpoken of, and which ſerves as a reſervoir for this humour, is three or four lines in length, and two at moſt in breadth at its baſis. It never contains more than four or five drops of the venom, which is forced from it principally by the action of a ſtrong and powerful muſcle, that riſing out of the lower jaw, folds inwards a little, then makes an arch, and preceeding to the upper jaw, runs over a part of it, and faſtens itſelf there. Towards the inner angle of this conſtrictor muſcle, or rather towards the part of its curvature neareſt the upper jaw, the ſmall veſicle begins; it is covered with this muſcle for almoſt the whole of its length. Thus placed, it is as it were encloſed in a preſs; and is bound and fixed to the adjacent bony parts, by means of two tendons, and of the canal, ſo that it can neither force itſelf forward, backward, nor ſideways, and muſt neceſſarily bear the double action of this muſcle, which now compreſſes it, when the viper bites and preſſes forcibly, and now again ſuffers it to dilate, when the muſcle itſelf contracts and enlarges. That which proves this muſcle to be chiefly intended to force the venom from its reſervoir is, that it is faſtened to each jaw in ſuch a way as to be

of very little uſe to the creature in cloſing its mouth, which clearly cannot be its principal purpoſe.

The hairs of a fox's whiſkers penetrate and paſs eaſily from the veſicle through the excretory duct, and go out at the orifice ſituated at the inner part of the ſheath. I have ſometimes ſucceeded in bringing them even to the elliptical hole at the point of the tooth. This is certainly the route the venom takes for the purpoſe of going out at the ſmall orifice of the ſheath, which correſponds preciſely with the height of the parabolical hole of the tooth *(a)*.

As the ſheath is very nicely fitted to the baſis of the canine tooth, the venom that goes out of the conduit at the ſmall orifice, muſt of neceſſity enter entirely into the hole of the tooth; and although it guſhes in abundance through this canal, it does not at all ſcatter itſelf in the ſheath, ſince the orifice out of which it flows is infinitely ſmaller than the parabolical hole to which the nice fitting of the ſheath makes it immediately correſpond. In a word, it paſſes entirely into it, particularly when there is only one of theſe teeth. Still more, I have obſerved on folding the ſheath back above the baſis of the teeth,

(a) It muſt appear very ſtrange that Doctor James, who has written after Mead, has aſſerted in his Diſpenſatory, that the true reſervoir of this liquor is the bag (ſheath) which covers the roots of the large teeth of the viper, and that a ſmall veſicle is found at the top of this bag, which opens at its extremity, to give a paſſage to the teeth that ſhed the venom. It appears however that this writer made many experiments on the viper, and with the intention of making them well.

and

and preſſing ſlightly and gradually upon the conduit, that the venom is carried by a natural declivity towards the hole of the tooth, which it entirely fills before a drop of it is ſpilt in the ſheath. This natural declivity is ſimply cauſed by a ſmall hollow in the jaw, which extends as far as the parabolical hole, and is ſcarcely ſeen with a microſcope. I do not however deny but that in ſome particular caſes this liquor may flow directly into the ſheath, and even glide from thence to the points of the teeth, particularly when two of them are ſo cloſe to each other as to touch and leave nothing but a hollow betwixt them, and when the viper bites in ſuch a way as to force its teeth deep into the fleſh, and to ſtop up the parabolical hole; it muſt here ſqueeze with ſufficient force and long enough to compreſs the veſicle, and give time to the liquor to glide betwixt the two teeth. In theſe caſes, which cannot however but be very rare, there is no doubt but that the animal may even kill without the venom having made its paſſage through the uſual conduit of the tooth. I repeatedly ſtopped up with pitch, ſometimes the parabolical hole, at others the elliptical hole, and ſometimes again both of them, and found that the yellow liquor did not then reach the bottom of the ſheath but with great difficulty, and after a ſtrong compreſſion had been made for a long time on the conſtrictor muſcle. I lay it down from hence, as an infallible concluſion, that the venom flows from the point of the tooth, and not form the

bag or ſheath, whether the viper of itſelf cauſes it to flow in biting, or whether a compreſſion is purpoſely made on the veſicle I have ſpoken of.

CHAPTER IV.

The Venom of the Viper is no other than the yellow Humour that flows from the Tooth when the Viper bites.

It happens very often, in vipers particularly that have been lately killed, that this yellow humour dries, ſtops up both holes, and totally obſtructs the canal of the tooth. As it cannot then enter into the tooth ſo as to find a paſſage through it, it muſt conſequently flow from the excretory conduit into the ſheath. This obſervation is ſo much the more neceſſary, as without it 'twould be eaſy to fall into an errour, and to preſume that the venom, inſtead of being conveyed by the tooth, flows from the ſheath, and is carried from thence into the wound.

I was deſirous of aſſuring myſelf how far one may rely on the opinion of thoſe who believe that the bite of the viper is only mortal on account of the rage the creature is thrown into before it bites. I omit the mention of an infinite number of experiments I have made to be certain, with Redi, that the yellow humour which flows from the tooth of the viper is mortal when introduced immediately into the blood h e medium of a wound. I ſhall only obſerve, that the various experiments of Redi and Mead agree

agree perfectly as to the truth of this circumſtance, and I cannot conceive how certain celebrated writers have continued to perſuade themſelves to the contrary, and to attribute the mortal effect of the bite of the viper to the rage of the animal, and to the power of the exalted ſtate of the ſaliva, rather than to the ſpecifick character of this humour.

I have frequently enraged vipers, and afterwards opened the mouth in ſuch a way that they could neither compreſs nor bite with it. I have then ſoaked bits of cotton well in the foam or ſaliva with which it was covered, and applied them to the wounds of animals, the bleeding of which had ceaſed. I have never ſeen any accident cauſed by this, nor have the animals appeared to be the leaſt diſordered. Neither the foam then, nor the other humours of the viper's mouth, are capable of cauſing death when introduced into the blood of an animal.

I have ſevered at one ſtroke the heads of ſeveral vipers from their bodies, at a time when far from being enraged, they were in a calm and tranquil ſtate. I have then taken the venom from the tooth itſelf, to be ſure of having it pure and unmixed. I have taken it from ſome of them immediately on ſeparating the head, and from others ſome hours after, when the head had dried in a great meaſure, and had ceaſed to move. On applying this venom carefully to the wounds of different animals, they have all without any exception been killed by it. We muſt conclude then that the humour alone which flows from the tooth, has a deadly quality,

to

to which the rage of the animal does not at all contribute. But to obviate all objection, and to prevent the being reproached with having neglected to make a viper bite after having enraged it, and having contented myſelf with introducing its foam into wounds, I took one, and made it bite ſeveral animals. When I conceived that there could be no longer any remains of the venom, I began to prick and torment the creature, and in a word, employed all the means I could think of to enrage it. When I ſaw by its hiſſings, and the rapid vibrations of its tongue, that it was become furious, I preſented animals to it, which it bit with all the force it was capable of. Neither of them however died, or ſeemed to feel any inconvenience. This was a very natural reſult, ſince the liquor that flows from the tooth, which alone has the faculty of killing, had already been entirely waſted, and ſince nothing more now remained than the foam and other humours that are in no way venomous, even during the moſt exceſſive rage of the animal. I repeated this experiment on two other vipers, with the very ſame ſucceſs.

I was deſirous of making another experiment, which, to prevent its being dangerous, requires a great deal of precaution and addreſs on the part of the obſerver, although after all it cannot be more deciſive than the preceding one. This was to entirely remove the two veſicles that contain the venom. After ſeveral fruitleſs attempts I at length ſucceeded, without doing much injury to the viper, and without tearing its mouth. I made an inciſion into

into the ſkin that covers the two veſicles, and having ſeized them with pincers, cut them entirely out with a biſtoury. Thoſe who are accuſtomed to diſſect this ſpecies of animals, muſt ſee clearly that this experiment is attended with more danger than difficulty. To ſucceed in it, the neck of the viper muſt be ſeized by ſome one, or it muſt be well tied to a table, in ſuch a way that the creature cannot raiſe its head to bite. Having removed the veſicles, I had two frogs bits, ſo as to diſcharge whatever venom might remain in the teeth or in the extremity of the conduit: neither of them died. I kept this viper a long time, and at different times made it bite both large and ſmall animals, as well with warm blood as with cold. Neither of them died, nor had any other complaint than what muſt have neceſſarily been cauſed by the ſimple mechanical wound of the tooth.

CHAPTER V.

The Venom of the Viper is not a Poiſon to the Viper itſelf.

Very grave authours have likewiſe imagined, that this humour, which deſtroys other animals, is not leſs hurtful to the viper itſelf; and this is the opinion of thoſe who have written lately on the venom of animals. The examples of ſcorpions and ſpiders

which kill each other with the bite or ſting, ſeemed to favour this opinion very ſtrongly. We read in the philoſophical tranſactions that rattle-ſnakes die in a very few minutes, when they bite each other. It is at preſent known that this ſnake is a ſpecies of viper, larger than ours, and therefore they have by analogy drawn the ſame concluſion as to the viper and other venomous animals.

Certain Spaniards had brought from the Eaſt Indies three ſnakes called *Cobras de capello*, and one only had ſurvived the frequent combats they had amongſt themſelves. Doctor Mead concluded that the other two died of the venom, and conſequently that the viper's venom ought to be likewiſe mortal to the viper itſelf. It ſeems to me that he ought rather to have drawn a contrary concluſion, for it is not likely but that the victorious ſnake which ſurvived, would have been ſometimes bit by the two others; and yet we ſee that it contrived to live.

'Twould have been undoubtedly much better to have made experiments, than to have founded an opinion ſo ſlightly on a mere matter of fact as Mead has done, and on a ſimple analogy drawn from caſes that are very rare; particularly as the fury with which ſcorpions and ſpiders combat and mangle each other does not prove that they die of the venom they have received. Beſides, it has been obſerved that the ſpider which leaves the combat victorious, only dies when it has loſt ſome one of its organs neceſſary to life. As to the rattle-ſnakes, the examples we have of their combats are too rare and not ſufficiently authenticated

ticated to furniſh a good analogy. This could beſides be never any thing more than a ſimple analogy, rendered ſo much the weaker by there being certainly a very great difference betwixt this ſnake and our viper, whether we regard their ſtructure, or the activity of their venom.

It is not eaſy to provoke vipers to bite each other, whatever care may be previouſly taken to kindle them to fury. I employed the following method to get the better of this repugnance. Having ſeized the neck of a viper with pincers, I kept its tail in my other hand to manage it with greater ſafety. I employed an aſſiſtant to ſeize a ſecond one in the ſame way. I held the body of one of them to the head of the other, which perceiving itſelf to be cloſe graſped by the neck, hiſſed, twiſted itſelf, and fell with fury on every thing that came near it. The former one, which it bit ſeveral times, was much ſmaller, and expreſſed each time, by the livelineſs of its motions, the violence of what it ſuffered. I found a ſuperficial wound at the part where it had been bit, moiſtened with blood and with the venom from the tooth. I encloſed it in a glaſs vaſe, where it continued tranquil for ſome minutes. Two hours after I found the part where it had been bit a little ſwelled; this ſwelling however continued but a little time, and the viper, recovering its natural vivacity, crept along the ſides of the vaſe, and raiſed its head with the ſame ſtrength as if it had not been bitten. After twelve hours I ſet it free, when it appeared as ſtrong

ſtrong as another one I placed with it by way of compariſon. I put it again into the vaſe, and on the following day found it ſtill as lively and healthy as before. At length, thirty-ſix hours after, ſeeing no appearance of its having been envenomed, I killed it. I found ſeveral holes in the ſkin at the part where it had been bit; the muſcles themſelves of the back were very deeply pierced; and the blows of the tooth had in more than one place forced it through the body, and through the abdominal viſcera. And laſtly, the wounds were a little inflamed, but there was no appearance of ſwelling or tumour.

Two days after I tried two very large vipers, which threw themſelves furiouſly on the animals that were preſented to them. I made them bite another middle ſized viper, which received, from one of them two very deep wounds made with the teeth, from the other four. One of them even left a tooth in the wound. At every blow the viper received on its belly, all of which were directed to the ſame ſpot, it gave violent ſymptoms of pain, hiſſed, and nearly eſcaped from the hands of the perſon that held it. I put it into a glaſs vaſe, in which it remained for a few minutes in a ſtate of inſenſibility: however, on afterwards placing it on the ground, it crawled about with great agility. I could never diſcover any ſwelling at the part where it had been bit; the ſkin had notwithſtanding been lacerated, and the muſcles laid bare: there was no hemorrhage. I kept it four days in the vaſe, during

ring which time it did not appear in the leaſt diſordered. The ſecond day I held an animal to it, which it inſtantly bit, and which died two hours after. I at length killed it and found that the blows of the teeth had pierced it through and through: the wounds were ſomewhat inflamed. The ſame thing happened to five other vipers which I had bit repeatedly. I even forced a ſixth to bite itſelf at the tail. Neither of them died, nor appeared to be in the leaſt diſordered.

But to prevent any one from thinking that the hardneſs of the ſkin had kept the venom from penetrating, and to introduce it with greater certainty into the blood, I removed a conſiderable portion of the ſkin from the backs of four vipers, and had them bit by ſeven others, from which they actually received ſeveral blows of the teeth; neither of them died, or became ill, and only one of them appeared to be heavy and languid, and had a ſwelling at its back.

Again, I irritated another viper, by pricking it on the body with a pointed iron, and afterwards made it bite a piece of jagged glaſs. The venom ſpread from the tooth over the whole mouth, which the glaſs had cut and made bleed. I kept it ſtill, and waited the event. For the three firſt days it crawled about a little. On the fourth it was more lively, but did not yet make any attempt to bite, even when provoked. On the ſeventh day I opened its mouth, and found it entirely healed, without any ſcar. On the ſame day I made it bite a ſmall animal, which died an hour after.

I re-

I repeated this experiment on three other vipers, and employed the following method. From one of them I removed a portion of the ſkin of the neck, from another a portion of that of the back, and laid bare the muſcles of the third juſt above the tail. I wounded each of them at the part laid bare, bending the point of the lancet a little, to open the wounds the better. Into each of theſe wounds I introduced a ſmall drop of the venom, that is to ſay, as much as was neceſſary to entirely fill it. I afterwards returned theſe vipers, each into its vaſe, where they remained very tranquil, and ſeemed to have ſuffered but little. Their wounds were however inflamed, but there was no ſwelling. I kept them alive for ſeveral days.

We now ſee what opinion ſhould be entertained of the analogy betwixt the venom of the viper and that of other animals, and may judge into how great an errour thoſe have fallen who have believed that the yellow humour which flows from the tooth of the viper, and which is mortal to other creatures, is likewiſe ſo to itſelf, and that the bites of theſe dangerous animals are capable of poiſoning each other. If analogy could have any weight in this inſtance, I ſhould be tempted to belive that, contrary to the opinion of Mead, the venom of the ſcorpion would have no ill effect on the ſcorpion itſelf, and that there is probably no venomous animal on earth, the venom of which can be hurtful to thoſe of its own ſpecies. If it be ſo, it can only be in a very few animals, and only in the ſmalleſt of them, the venom

of

of which is acrid and caustick, such as bees, wasps, and hornets. It may likewise be true that the scorpions of Asia and Africa carry a venom that is mortal to their own species, since the venom of the Italian scorpion, when put on the tongue, is acrid and pungent. It appears to me, that the general errour which many observers, who are otherwise very exact, have embraced, has its source in a deceitful experiment. It had been remarked, that when a scorpion was surrounded by live coals, it was first agitated, and then turned its sting towards its back, as if to wound itself. As it at length died, and was even burned up, from its violent agitation amongst the live coals, it was simply believed that it died from its venom, and from its own wounds. The experiment is equivocal; it is even false. I have repeated it a thousand times, and have never observed that the scorpion struck itself with its sting; it died roasted and burned up, and not envenomed.

It has likewise been observed, that the fresh water polypus, in swallowing its prey, sometimes swallows the arms or claws with which it holds it; and likewise, that when two of these polypusses dispute together, the stronger frequently prevails, and swallows the claws of the weaker. The polypus, however, in neither of these cases dies, although its venom, as we shall see hereafter, is very active. The parts thus swallowed, leave the stomach soon after, entire and alive, without having suffered any apparent change, and continue as before to serve as claws to the animal.

CHAPTER VI.

The Venom of the Viper is not a Poiſon to every Species of Animals.

Thus far we have ſeen that the venom of a viper is neither a poiſon to the viper itſelf, nor to thoſe of its own ſpecies; and this ſingularity led me to ſuſpect that it might alſo be innocent to ſome other kinds of animals. Indeed why ſhould it not be ſo, as well as to the viper? If, in a word, it is not capable of decompoſing the ſolids, and altering the fluids, of any particular living machine; if it can neither diſturb the harmony of it, nor occaſion death; why ſhould there not be other living organized creatures, on which it may have as little action? We know but little of the manner in which poiſons in general act, but we know that there are many very active ſubſtances which produce the moſt terrible effects on certain parts, and which notwithſtanding have no effect on others. Stibiated tartar, for example, a ſubſtance that is introduced without danger into the eyes, is a very violent emetick when received into the ſtomach. There are perſons who are thrown into convulſions by the fragrant ſmell of the roſe. Theſe various accidents are owing without doubt to the ſtructure and organization of the animal machine.

machine. Certain ſubſtances are known to be poiſons to certain animals, whilſt far from being hurtful to ſome others, they even ſerve as an aliment to them. Such is hemlock for inſtance, which deſtroys the human ſpecies, and nouriſhes goats. It is thus that the bitter almonds we eat on account of their flavour, kill certain birds, and do no injury to others. It may likewiſe be the caſe then, that the venom of the viper may not be a poiſon to all kinds of animals, particularly if it act like narcoticks, that do not cauſe death by corroding the ſolid parts. Corroſive ſublimate is a poiſon deſtructive to all living animals, becauſe its mechanical action is in fact capable of exerciſing itſelf on all the organs of the animal machine. Narcoticks, on the contrary, ſo dangerous to men, produce no ill effect on dogs. The different ſtructure of the organs of animals may then be the occaſion, that the ſame ſubſtance may at the ſame time be a very active poiſon to certain ſpecies of them; and altogether of an indifferent nature, or an aliment, or even an excellent remedy, to others.

It is on theſe conjectures that I engaged in the long courſe of experiments I am going to relate. I had already obſerved, that of all animals, leeches are inconteſtibly killed with the greateſt difficulty. When they are cut in pieces, each portion preſerves for ſeveral months the motions it had before it was ſeparated from the others. I conceived that an animal ſo tenacious of living, might well reſiſt the venom of the viper, without dying or even being incommoded

by it. I fixed then upon leeches, but before I had them bit, I took care on removing them from the water, to wipe them very dry with a piece of linen, fearing that the mucoſity or kind of glue that covers them, and which they emit when touched, might prevent the ſucceſs of my experiment. I had one of the largeſt kind, that are called horſe leeches, bit by a very ſtrong viper which I had previouſly thrown into a violent rage. It pierced its body, from which a ſmall quantity of blood flowed, ſeveral times through the whole ſubſtance of it. I afterwards put the leech into water, and it continued to move as uſual. On the day following I changed its water, and found it very lively and ſwimming perfectly well in the glaſs. It lived in this way for ſeveral days, and would certainly have lived much longer, had I not applied it to another purpoſe.

I took a ſmaller one, of that ſpecies that have different coloured ſtripes on the back, and that are employed in medicine. I had it bit by two vipers, which pierced its body in ſeveral places. It was bit the next day by a third, and the day following again by two others. Its ſkin was full of holes, from which a viſcous and black matter flowed, on preſſing with the fingers. Notwithſtanding this it continued to live, and move about in the water. Laſtly, I had many other leeches of both kinds bit in the ſame way, at ſome times in the head, at others in the body, and neither of them died of the venom.

I did

I did not ſtop here, but fearing that the venom might have been enveloped and deadened by the glutinous humour of the leeches, which oozes out in greater abundance the moment the tooth of the viper has pierced the ſkin, I made deep wounds in ſeveral with a biſtoury and with ſciſſars, and poured into them large drops of the venom. I introduced into the bodies of others, bits of tow moiſtened with the venom, and paſſed quite through, and this method, which I had always found mortal when tried on other animals, was quite without effect on this occaſion, ſince neither of the leeches died. I preſerved for ſeveral months, in glaſſes filled with water, parts of leeches quite alive. Each of theſe pieces preſerved the motions there, that it poſſeſſed when united with the other part of the body. I had ſeveral of theſe bit by vipers, and made notches in others, paſſing into them bits of tow ſteeped in the venom: neither of them died. They preſerved all their motions, and did not ſeem at all incommoded. The leech then has the property of reſiſting the venom of the viper, which is quite innocent to it.

I afterwards wiſhed to try what would be the effect of the venom of the viper, on ſnails and ſlugs. I procured the largeſt of them, and of different kinds. I had ſome of them bit in ſeveral parts of the body, and by ſeveral vipers. I likewiſe made inciſions, into which I introduced the venom, taking good care before hand to wipe off the glutinous matter that covers them, that the poiſon

might penetrate the readier. Out of twenty-ſeven ſlugs and ſnails on which I made theſe experiments, one ſlug only died, twenty hours after it had been bit. I could not even ſucceed in killing them with the envenomed bit of tow introduced into their bodies. The greater part of them covered themſelves with their viſcous ſlaver on being bit.

In the country about Piſa a ſnake is found, which the peaſants call aſpick, and which they repreſent as more venomous than the viper. This creature has ſome exteriour reſemblance to the viper, but has neither the canine teeth, the ſheath, nor the veſicle or reſervoir of the venom; and the experiments I have made, have convinced me that it is an animal in no way dangerous. The ſnake with two heads that was brought to Redi, and of which he gives a deſcription at the beginning of his obſervations *on living animals that are found in living animals*, was of this ſpecies. That of Redi was ſingular however in having two heads. I wiſhed in the firſt place to be certain whether the venom of the viper is mortal to this kind of ſnakes, and had one of them bit twice in the tail by a large viper. Two days after, I had it bit by two others in the back, from which a little blood flowed, and after two days more held three other vipers to it, which gave it ſeven or eight blows on the neck with their teeth. It was rendered a little torpid by them, and was ſlower in its motions, but on examining it two days after I found it alive, and on putting it to the ground

ground it crawled along and ſeemed in perfect health.

The venom of the viper has no greater action on another larger ſnake, which in Tuſcany is particularly diſtinguiſhed by that name; this is the *adder*. I had ſeveral of them bit on the back, tail, neck, and belly; to ſome of them I have even held three vipers at a time: neither of them died. They did not even ſeem ſurprized at it, neither were they benumbed. I at length employed the envenomed tow, forcing it into wounds I purpoſely made. In ſome of them I even removed the ſkin at particular parts, to convey the venom the better to the blood. All theſe attempts were without effect. It ſeems certain then that the venom of the viper is in no way either mortal or dangerous to this ſpecies of ſnakes. It is not alone then on animals of the *worm* claſs that the venom of the viper is deſtitute of action; there are others again, the organization of which is more compoſed, and which have a heart and many viſcera, and are notwithſtanding out of the reach of its fatal influence.

I have found another ſnake called cecilia, the *orvai* of the French, which likewiſe reſiſts the bite of the viper. I have frequently made experiments on theſe ſnakes, and have had them bit by ſeveral vipers at a time, and in different parts of the body. This creature, naturally ſluggiſh, did not ſeem incommoded by the venom, even when I introduced it into its body by the means of inciſions.

Theſe three ſnakes, the aſpick, the cecilia, and adder, are not venomous, ſo that one incurs no riſk, even when they bite ſo as to draw blood; they have no canaliculated teeth, no ſheath ſuch as that which covers the teeth of the viper, nor a reſervoir for venom; in a word, they are creatures perfectly innocent, as I have aſſured myſelf by many experiments.

I had two turtles bit by a very large viper, rendered furious, in the hind feet where the ſkin is the leaſt hard. I kept them alive for more than ten days, during which time they did not appear to ſuffer, and walked as uſual. I had another bit ſeveral times in the neck, and as a clear proof that the teeth of the viper penetrated through the rough ſkin, one of them was left in that had forced its way into the vertebræ. On the day following, I had it bit in the neck by another viper, and by a third in the fore feet. Laſtly, on the third day it was again bit by two vipers in the neck and hind feet. It not only ſurvived, but did not ſeem to have ſuffered the ſmalleſt inconvenience. One would have ſaid, on the contrary, that it was become more ſenſible and active.

I had five others bit in the breaſt and belly by eight vipers, after removing the inferiour ſhell, and laying the fleſh bare. Neither of them died. They were even living four days after, which is uſually the caſe with thoſe that have been deprived of the lower ſhell only. In others I made deep wounds in the feet, and even removed the ſkin in

ſome

ſome of them to introduce the poiſon the better. At length I forced into the wounds, large bits of tow ſoaked in the venom. Neither of them died nor had the ſmalleſt ailment.

I do not believe however that turtles are abſolutely beyond the reach of the effects of the venom. One of them died, after it had been bit by eighteen vipers. The blood oozed from the wounds theſe animals had made, in every part of its body. A third died twelve hours after it had been bit in the neck by three vipers only; and a third again in the ſpace of twenty-four hours, notwithſtanding it had been ſimply bit in the feet by two large vipers. It appears then that this venom does but rarely penetrate and diffuſe itſelf in the bodies of turtles, and that its action there is much ſlower and weaker than in the other animals with cold blood. Theſe, laſt in general, die from the effects of this poiſon, at leaſt all thoſe I have had bit, not even excepting eels, which however die later, and not till the end of eighteen or twenty hours. The other kinds of fiſh are likewiſe killed by the venom of the viper. And laſtly, the ſmaller lizards ſcarcely ſurvive its bite for a few minutes.

Animals with warm blood are univerſally killed by this poiſon; I have at leaſt never met with any one that ſurvived its action. A ſmall goſs hawk died in leſs than three minutes. In a few ſeconds it began to open its beak, as if it felt a difficulty of reſpiration, and had an inclination to vomit. A few inſtants after it fell on its breaſt, and could not again

again recover its feet. It at length died with all the ſymptoms of an extreme debility. I have generally obſerved that animals with warm blood, and the action of the heart of which is very lively, die much ſooner than the others.

There are ſeveral kinds of animals, very diſtinct from each other, to which the venom of the viper is not a poiſon; or if it be ſo, it is but very rarely, and that with the leaſt poſſible energy. There are perhaps many others we are ignorant of, that reſiſt its action. I have myſelf found ſeveral of the ſpecies of inſects and worms to which this venom is not hurtful. I ſhall perhaps ſpeak more fully of them in another work, in which I ſhall treat of the remedies againſt the bite of the viper.

All theſe particulars ought to render the philoſopher who ſtudies nature very circumſpect, unleſs he wiſhes to bewilder himſelf at every ſtep; they likewiſe ſhow us how little truſt is to be repoſed in the ſimple analogy that may be found betwixt different animals, either as it regards life, or the economy of their motions. Nature does not ſuffer herſelf to be devined. Experiment alone, in the hands of an attentive and diſcerning obſerver, can ſnatch from her her ſecrets.

CHAP-

CHAPTER VII.

The Venom of the Viper is not Acid.

IN a ſmall publication of Mead on poiſons, printed in 1739, with the falſe indication of Amſterdam and Naples, the venom of the viper is ſaid to be acid, and to change the blue colour of the turneſol to red; of the truth of which he ſays he is convinced by his own experience. To be certain of this, I received the venom of a viper I had juſt killed, on a bit of glaſs, forcing it immediately from the point of the tooth, by a gentle compreſſure of the palate. I afterwards poured this venom on a bit of blue paper, which ſoaked it up, but inſtead of becoming red, turned a little yellow, and preſerved this appearance even after it was dry. It appeared extraordinary to me that ſo learned a man as Mead ſhould have been deceived in ſo eaſy an experiment. I therefore took a greater quantity of venom, with which I rubbed ſeveral pieces of blue paper, and that nothing might be neglected, varied the experiment a hundred different ways. At times, to have the venom the purer, I took it immediately from the tooth, before it had touched the other parts of the mouth; and at others again, either forced a bit of cotton into the mouth of a

living

living viper at the moment of its biting, or introduced it into that of a dead one filled with the venom. I diluted a quantity of venom in water, and wet blue paper with it. I tried to find whether the mixture of the venom with the other humours of the animal, had not deceived Mead as to the colour, and varied my experiments for that purpoſe infinitely, but in vain. I could never turn the paper red. It ſimply took the yellowiſh tinge found in the venom itſelf. Mead likewiſe maintains, that he has ſeen the mixture of this liquor with violets become ſomewhat red: I have tried this, but the event has not been the ſame. When the venom is in a greater proportion than the ſirop, the mixture does indeed become a little yellow, but never becomes red. I increaſed, I diminiſhed the quantity of the venom; I have taken it pure, and again have employed it mixed with the foam of the viper: I could never perceive any thing beſides a ſlight yellow tinge, and all my experiments have only ſerved to confirm me in the opinion that the venom of the viper neither changes red the ſirop of violets, nor the dye of the turneſol (*a*).

In the ſame work on poiſons, Mead maintains that the venom of the viper is a true acid, and that it effervesces with alkaline ſubſtances. In conſequence of this I took ſeveral fluid alkalies, ſuch

(*a*) Doctor James is likewiſe of opinion that the venom of the viper is acid, becauſe, according to him, it changes the dye of the turneſol and ſirop of violets red, as other acids do.

as

as the ſpirit of hartſhorn, and oil of tartar *per deliquium*, with which I mixed different quantities of the venom, always pure and unmixed with the other liquors of the mouth. I never could obſerve the ſmalleſt motion nor the leaſt effervescence, at the moment of their union. It was in vain that I had recourſe to a microſcope, I could never obſerve the ſmalleſt air-bubble diſengage itſelf; the colour remained the ſame, and I met with nothing that gave me the ſmalleſt ſuſpicion of the exiſtence of an acid in the venom. It muſt not be thought that the rapidity of the effervescence prevented my ſeeing it, ſince the drop of venom was ſo ſlow in uniting itſelf to the alkalies, that it was eaſy to follow it with the microſcope, and to ſeize the preciſe moment of their perfect union.

CHAPTER VIII.

The Venom of the Viper is not Alkaline.

AS authours are to be found who pretend that the venom of the viper is alkaline, and not acid, and as it is principally on the activity and ſuddenneſs of its effects that they have founded this their hypotheſis, I thought it adviſable to conſult experiment thereupon. I took then different acid liquors,

liquors, ſuch as vinegar, ſpirit of ſalt, ſpirit of nitre, ſpirit of vitriol, and laſtly, ſeveral acid ſalts extracted from plants. I united with all theſe acids a larger or ſmaller proportion of the venom, but could perceive no other than a yellow colour, which appeared whenever the quantity of venom exceeded that of the acid. I armed myſelf with a good microſcope, and never found either efferveſcence, motion, or air bubble, to reſult from this mixture. I tried it afreſh with ſirop of violets, but it did not turn it green, as alkaline ſubſtances uſually do.

It is equally without foundation then that naturaliſts pretend that the venom of the viper is acid or alkaline; and it is ſtill with leſs reaſon that they have contrived to explain by theſe hypotheſes, the pernicious effects of this poiſon. Their irrational theories are completely belied by experiment, the only guide to thoſe who enter into the ſearch of phyſical truths. It muſt however be acknowledged, that Dr. Mead has corrected many errours as to facts, in a new edition of his work on poiſons, printed in Paris in 1751, which has reached me too late. He there retracts what he had advanced on the acid quality of the venom of the viper. He confeſſes that the experiments made with the turneſol and ſirop of violets are falſe, and that the venom neither efferveſces with acids nor alkalies. This avowal prevents me from endeavouring to account for the contradiction betwixt his experiments

ments and mine, and from pointing out what may have occaſioned his errour.

CHAPTER IX.

No Salts are diſcovered in the Venom of the Viper.

THUS have I the ſatisfaction of being the firſt to confirm, by experiments more numerous and more diverſified than his, the truths which Mead has diſcovered, and which no one that I know of has buſied himſelf about ſince him. This conformity fixes in an unvariable way the certainty of my obſervations.

In the courſe of my reſearches I have examined with the moſt ſcrupulous nicety into the exiſtence of that pungent and cauſtick ſalt, which Mead himſelf in his laſt work, and all the obſervers after him, ſay they have ſeen in the venom of the viper *(a)*.

Mead regards it as a neutral ſalt. He pretends that he has ſeen it floating in the ſtill liquid venom, and deſcribes it as formed of very ſharp

(a) James maintains with Mead that he has ſeen theſe ſalts, although in a ſmall quantity, in the diluted venom. They both ſay, that the net-work it forms in drying, is entirely compoſed of ſmall cryſtals.

points. But what was my ſurprize when on examining the venom with a microſcope, I could never diſcover this collection of ſaline cryſtals which the learned Engliſhman believed he invariably ſaw! I even employed, but ineffectually, the very ſtrong lens' made in England. I could find nothing throughout beſides a yellowiſh and viſcous humour, without any determinate ſhape, without diſtinct floating corpuſcles or particles, and alike in all its maſs, as an oil of any kind appears when viewed with a microſcope. The venom I employed was pure and taken from the tooth alone. I varied this experiment an hundred different ways, and even had recourſe to the ſolar microſcope; I at length ſatisfied myſelf that there are in reality no ſalts in the venom, and that Mead muſt have been impoſed upon by ſome particular circumſtance.

I then recollected that I had formerly ſeen with a microſcope, certain tranſparent bodies which floated on the human ſaliva, and which might eaſily have been taken for ſalts. Indeed any one who is not very converſant in the uſe of the microſcope, and who is not well acquainted from habit with the ſhape of the different ſalts that are found in liquors, particularly whilſt they are drying, would eaſily perſuade himſelf that the ſmall diaphanous particles which float on the ſaliva, are abſolutely of a ſaline nature. They are however too light, too large, and not ſufficiently tranſparent, to be really ſalts. They vary both in ſize and ſhape. The direction of theſe ſmall bodies is rather

ther crooked than ſtraight; they have hollows and folds on their ſurface; and laſtly, they become ſhrivelled and obtuſe in proportion as the ſaliva dries. Thus are they to the eyes of a practiſed obſerver, nothing more than ſmall pellicles or light, plaited, membranes, and as it were the relicks of almoſt digeſted aliments. In reality, they diſappear on waſhing the mouth well, and I have obſerved on touching them with a fine and ſharp needle, that they lengthen or ſhrivel up like ſmall bits of ſkin. I have met with ſmall floating bodies ſimilar to thoſe that are found in the human ſaliva and in that of animals, with the aſſiſtance of a microſcope, in the ſalivary humour of the viper. I have likewiſe ſeen ſome of them floating in a drop of venom I had caught on a ſmall ſilver ſpatula, put into the mouth of a viper, the palate of which was ſtrongly compreſſed. I then conceived how Mead was led into this errour. He certainly took the venom from the mouth of the animal, and not in the way I did, immediately from the tooth; and regarded the ſmall bodies which proceeded alone from the ſaliva, as belonging to the venom.

It is likewiſe true that ſmall bodies or globules, ſomewhat yellow and tranſparent, are often found in the venom of the viper whilſt yet fluid. This never happens but when a ſtrong compreſſion is made on the palate or veſicle, at which time, far from being pure, the venom flows mixed with other corpuſcles ſupplied by the reſervoir.

In Mead's works we likewiſe find an obſervation, which is repeated in the Paris Edition, and which appears to eſtabliſh the exiſtence of theſe ſalts in a clear and evident way. He ſays, that in examining with a microſcope the venom of the viper put on a bit of glaſs, the ſaline particles, in proportion as it dries, are ſeen to form themſelves into very fine and ſharp chryſtals, reſembling a very fine ſpider's web; and that theſe tranſparent chryſtals or needles continue perfect for ſeveral months, ſo ſtrong and firm are they, notwithſtanding their ſmallneſs.

I took then a drop of the viper's venom, perfectly pure and free from any mixture with the other liquors of the mouth. I dried it on a piece of glaſs, and viewed it with a microſcope. What was my ſurpriſe on obſerving, inſtead of the drop, a heap of different tranſparent bodies, of an equal ſurface, and diſpoſed with great ſymmetry and regularity! their ſhape was in general quadrilateral or triangular, and their points very ſharp, ſo that they ſtrongly reſembled the net-work Mead has deſcribed. Their regularity and tranſparency might at firſt ſight very eaſily cauſe them to be taken for ſalts, but they were too large, and arranged with too much order, not to make one miſtruſt this appearance. What at length fixed me in the perſuaſion that they were not chryſtals was, that I did not ſee any of them in cluſters, as they are found in other ſalts; they were all diſtinct, and placed at equal diſtances from each other. Perſons who are accuſtomed to view the ſalts of other fluids, muſt perceive the weight of theſe laſt proofs.

proofs. I now ſuſpected that the venom had ſplit and cracked in different places in drying, and that this had occaſioned its being thus divided on the glaſs, as happens to ſeveral ſubſtances, which when they dry, ſplit in this way into thouſands of fragments, either pretty regularly ſquared, or in a triangular form, and all at equal diſtances. If theſe cracks are throughout of the ſame ſize, it is owing to the ſame cauſe, that is to ſay, evaporation, acting at the ſame time and with ſame the force on the whole ſurface; it is from this that it repreſents a kind of net-work with different meſhes, exactly like the web of a ſpider.

Laſtly, to make myſelf ſtill more certain that theſe were not ſalts, but rather ſo many ſcales and broken pieces of the dried venom, I fell upon a new experiment which I thought a deciſive one. I dried a few drops of the venom, in a very pure ſtate, in a ſmall concave glaſs; I then examined them with a microſcope, and found them as uſual, full of ſmall crevices, repreſenting a ſpider's web. It was however very clear that theſe chinks, towards the bottom of the glaſs, were larger in proportion as the dried humour had a greater thickneſs. Theſe pretended ſalts were no other than the fragments of venom ſeparated and dried on the glaſs. Thoſe that were the thickeſt had little or no tranſparency. They were of a yellow colour, like the venom itſelf in a fluid ſtate. They are ſimply cauſed then by the parts of the venom retreating from each other dur-

ing the evaporation; and this is even viſible to the eye, without the aſſiſtance of a microſcope.

But to remove all doubt and ſuſpicion on a matter ſo important and ſo generally adopted, and on which Mead has in ſhort founded his hypotheſis of the action of the venom carried into the blood of animals, I made another experiment, which abſolutely proves the nonexiſtence of this pretended ſaline net-work. I put a drop of venom on a flat and ſmooth glaſs, and followed it very attentively with the microſcope during the whole time of its drying: nothing occurred however ſimilar to what happens to ſalts diſſolved in water. The ſaline particles during the progreſs of the evaporation collect together and approach towards the centre from the circumference, forming at firſt very ſmall chryſtals, which encreaſe in bulk, from the addition of ſaline particles of the ſame nature which unite with them. Here, on the contrary, I found nothing beſides a humour which, in proportion as it dries, cracks and preſents furrows, that form the quadrilateral and triangular fragments I have mentioned. Theſe crevices, which are like the ſpaces betwixt the threads of a net, appear at firſt at the circumference, and proceed gradually towards the centre in proportion as the deſiccation advances. But the quadrilateral and triangular fragments that fill the ſpaces betwixt the crevices, and repreſent meſhes, do not encreaſe here as the ſaline particles do in a diſſolution of ſalt during the progreſs of the evaporation. I repeated this ſeveral times with a ſingular pleaſure. I mixed

the

the venom with a few drops of very pure ſpring water, which I left to evaporate, and obſerved it patiently with a microſcope, hoping in this way to diſcover any ſalts it might contain, but I was not ſo fortunate: no better method can however be fallen upon for this purpoſe.

Two celebrated Profeſſors of the Univerſity of Piſa, Meſſieurs Perelli and Lampredi, were witneſſes to my experiments. They very gladly honoured me with their preſence, and conſtantly aſſiſted me, particularly when I made my reſearches on the ſalts of the venom of the viper. They both agree that whatever reaſon they might have previouſly had to ſuſpect their exiſtence, my experiments, joined to a little reflection, have been more than ſufficient to deſtroy the very ſhadow of a ſuſpicion.

It muſt likewiſe be remarked, that the clefts which form when a large drop of the venom is evaporated, are much larger than when the drop is ſmall, or when it is diſſolved in water, or very much ſpread on the glaſs; theſe large clefts are diſpoſed like rays that proceed to an union with each other, towards the centre of the dried venom. The ſpace betwixt theſe rays is likewiſe diſſected by other tranſverſe rays, which become cloſer in proportion as they approach the centre, and form the above-mentioned figures, beſides many other very irregular ones. Theſe tranſverſe clefts are ſmaller at the circumference, are at greater or leſs diſtances from each other, and are bent into ſegments of a circle.

When the venom of the viper is viewed with a microſcope, very ſmall and tranſparent particles or ſpots are likewiſe ſometimes obſerved in it, which are the laſt to dry.

Thus have I fully ſatisfied myſelf of the nonexiſtence of thoſe ſalts which phyſicians and naturaliſts have hitherto admitted with ſo much confidence. I have ſeen the theories founded on this principle, to explain the action of the viper's venom, fall and vaniſh before experiment, which proves that no ſalt, either acid, alkaline, or neuter, exiſts in this humour.

CHAPTER X.

The Venom of the Viper has no determinate Taſte, and when put on the Tongue cauſes no Inflammation.

FROM the teſtimony of Redi, the venom of the viper was at firſt thought to be inſipid, and ſomewhat ſimilar in taſte to the oil of ſweet almonds. We however find in no part of his works that he experienced this himſelf. He ſeems on the contrary to have truſted in this reſpect to a certain Jacques, a viper-catcher, who was venturous enough to taſte this dangerous liquor. He boaſted that he could ſwallow a whole ſpoonful of it, and Redi tells us that he has been ſeen to take it ſeveral times.

Mead

Mead, on the other hand, aſſures us that he has taſted it himſelf, that he has made others taſte it, and that it is acrid and pungent; he ſays that it leaves a ſenſation of burning on the tongue for ſeveral hours, notwithſtanding it is diluted with warm water. He adds, that a pain and ſwelling of the tongue ſoon rewarded the temerity of him who taſted it pure. Theſe contradictions reduced me to the philoſophical neceſſity of taſting the venom myſelf. I did ſo, but not without repugnance; and as the celebrated Morgagni obſerves in his excellent letter on poiſons *(a)*, I ſhall adviſe no one to try it in the gaiety of his heart, leſt he ſhould happen at the time to have ſome excoriation on the tongue, which is a circumſtance not always eaſy to determine. Here however a point was to be ſettled which has divided the opinions of the moſt modern and moſt reputed authours.

I put a drop of the venom then on a bit of glaſs, and diluted it with ten or twelve drops of water; I touched the tip of my tongue very ſlightly with it, and immediately felt a ſenſation as it were of cold and inſipidity. I waited a little, in expectation of that burning ſenſation, which acid and cauſtick liquors occaſion, and at length withdrawing my tongue, paſſed it acroſs my lips, gums, and palate, that I might better come at the ſavour of the venom: notwithſtanding this I could find no taſte in it, except that of a very inſipid liquor. I then took all the venom I could expreſs from a viper, and

(a) De ſedib. et cauſ. morb. Epiſt. 49.

ventured to put it in a pure ſtate on my tongue, the point of which I rubbed well with it, as the moſt ſenſible part; I likewiſe rubbed my lips with it. I found a degree of conſiſtence and viſcoſity in it, but nothing acrid, pungent, or burning; in a word, it had no determinate taſte. It is however not ſo inſipid as pure ſpring water. There is ſomething in it that reſembles the almoſt inſenſible ſavour of the freſh fat of animals, with a very ſlight ſlavour which one can ſcarcely diſtinguiſh, but which would be pretty like that of the viper's fat, if this laſt was not ſtronger and more nauſeous.

I found no greater taſte nor ſmell in it on drying it, and reducing it to powder. As I could meet with no naturaliſt bold enough to make the ſame trial, and as a ſupport to my opinion, I gave it to my ſervant, a native of Tirol, named Jacques Benvenuti, to taſte. This man, as intrepid as the one Redi ſpeaks of in ſuch terms of admiration, ſwallowed it repeatedly at different times, at ſome times pure, and at others diluted in water, varying the quantity, and never perceived it to ſwell or burn either the tongue or the mouth. He ſaid however that when he took it pure and in a large quantity, the ſenſation he felt was very different from that excited by oil of ſweet almonds, pure water, or either acid or ſharp ſubſtances: but he could not tell in what this difference conſiſted. A ſenſation ſometimes continued on the tongue for ſeveral hours, not of pain, but as he deſcribed it, ſuch as is felt on taking

ſome-

ſomething aſtringent. His obſervation was juſt, for I myſelf have experienced this diſagreeable kind of ſenſation, which frequently continued for five or ſix hours, in the parts of my mouth where the poiſon had remained a long time. If it is taken in a ſmall quantity and mixed with water, it leaves no ſenſation on the tongue; and this ſpecies of diſorder in the mouth is not felt the inſtant the venom is taſted, nor immediately after, but only at the end of a certain time, and it is likewiſe neceſſary that the venom be kept a long time in the mouth. I have repeated theſe trials more than an hundred times, and have never had my tongue either ſwelled, inflamed, or painful. What is ſtill more, the venom when even applied to the eyes, cauſes neither pain nor inflammation. I have laid ſome of it ſeveral times on the tunica conjunctiva of different animals, ſuch as dormice, cats, and dogs, and neither tumour nor inflammation has ever ſupervened in this part, which is otherwiſe ſo ſenſible to the impreſſion of ſubſtances, frequently thoſe that are the moſt innocent. I have even introduced it into the noſes of theſe animals, without their ever betraying any ſign of ſuffering the ſmalleſt inconvenience from it.

It is certain then that the venom of the viper is in no way ſimilar to cauſticks, and that it is not acrid and hot like that of the bee or ſcorpion. Scarcely had I put an atom of the venom of the bee on my tongue, either pure or mixed with a little water, than it ſtung and burned with as much violence

lence as if I had applied the ſtrongeſt cauſticks that chemiſtry affords. The venom of the waſp and that of the hornet are not leſs acrid and pungent than the bee's, and the pain that each of them excites laſts a long time. I took it ſometimes from the ſting, and ſometimes from the ſmall veſicle that ſerves as a reſervoir to it, and found it in both caſes alike, and invariably productive of the ſame pain. It ſtill preſerves its ſtrength and cauſticity after having been dried, and kept for ſeveral days.

It is the ſame with the venom of the ſcorpion; the white and viſcous humour it throws out by its ſting when it darts it, cauſes a ſenſation on the tongue ſimilar to that occaſioned by the venom of the viper, but much weaker. It is on this account that the ſting of the bee is more painful than that of our ſcorpions. Probably the venom of thoſe of Africa is exceedingly cauſtick, ſince it kills animals in a very ſhort ſpace of time.

I afterwards made a trial of the viper's venom on other animals, which, although they are not, like man, gifted with ſpeech, are not backward in manifeſting by ſigns, the pleaſure or diſguſt they feel on eating any thing. I put then a drop of the viper's venom into the mouth of a dog; the creature ſwallowed it with avidity, licked its lips for a long time, as if it had met with ſomething agreeable to its taſte. I then ſteeped a bit of crum of bread in the venom, to ſuch a degree that it became quite yellow, and gave it to the ſame dog, at a time when it had already fed ſo plentifully as to refuſe

food,

food. It ſmelt to it, and inſtantly devoured it, manifeſting the ſtrongeſt deſire for more: in a word, every time a drop of venom approached its lips, it licked it up with the greateſt ſatisfaction.

Every body knows that dogs, like children, are ſworn enemies to whatever is bitter and acrid, and that they are paſſionately fond of whatever is ſweet and unctuous. Hence we muſt conclude, that if the dog found the venom agreeable, it was undoubtedly owing to its ſweetneſs. Thus is it abſolutely falſe and imaginary, that it is acrid and fiery; as it alſo is that the tongue, on taking it, ſwells, inflames, and becomes painful.

Mead had an idea that the venom of the viper, when applied to the wounds of a living animal, cauſed a very painful ſenſation: a natural conception to thoſe who believe like him, that it abounds in ſalts, which render it hot and cauſtick. He endeavours to eſtabliſh his opinion on an experiment he made on a dog. This animal did not ſeem very ſenſible of the pain occaſioned by piercing the noſtrils with a crooked grooved needle; but when the venom entered the wound, it howled and became furious. I made the very ſame experiment on a young dog, and it appeared inſenſible to the entrance of the drop of venom into the wound. I muſt acknowledge however that I have ſeen a cat ſhake itſelf and become more agitated, at the moment the venom was forced into the lips of a wound that had been made in its noſe. But this experiment is always liable to error, ſince the needle not only remains in the wound,

wound, but the motion of the animal is ſtill another cauſe of its being more agitated there, and of its being forced ſtill deeper, and cauſing a greater laceration of the parts. This is doubtleſs ſufficient to renew the pain, and even to wound the nerves that eſcaped the firſt introduction of the needle.

I have often poured the venom into inciſions made with a lancet, and could at no one time aſſure myſelf to a certainty, that the introduction of it was productive of pain; although it ſometimes happened that I was pretty much convinced to the contrary.

But were it well proved that the venom of the viper cauſes pain, does it follow that we can draw an indubitable concluſion that it abounds with ſalts, or that it is acrid and cauſtick? As if we had not examples of a juice which, though inſipid to the taſte, brings on violent pain when applied to a wound. I have myſelf known people, who having been bit by a viper, have notwithſtanding felt but a very ſlight pain, ſuch as might very well have been cauſed by the ſimple blow of the tooth. We have a dexterous viper-catcher at Piſa, named Bongi, who having one day been bit in the finger, did not perceive it till he ſaw the blood flow, a proof that he did not feel any pain. His father relates a ſimilar circumſtance; he likewiſe had been bit in the finger, and compares the pain it occaſioned to the bite of a fly. However they both in the concluſion became very ill of their wounds, a clear demonſtration of the venom having found its

way

way into the blood. I am well perſuaded then from experience, that the venom of the viper is neither acrid nor burning *(a)*, and that it does not contain thoſe ſalts which ſo many writers have imagined, either for the purpoſe of explaining its mode of action on the blood, or becauſe they have been imperfect in their obſervations.

CHAPTER XI.

Properties of the Venom of the Viper.

THE deadly humour furniſhed by the viper, which I have neither found to be acid, alkaline, nor cauſtick, ſubſides inſtantly on being thrown into water, like certain heavy oils drawn from vegetables. In this ſituation the parts of it preſerve their viſcoſity and natural union, and remain in that ſtate for ſome time, without changing either their primitive colour, or their tranſparency. This poiſon then is heavier than water, and differs in that reſpect from common oils, from the fat of animals, and from that of the viper itſelf, all of which float on water. Oils and other liquids that are heavier than water, ſhould be at leaſt ſuſpected, and are indeed often found, to be very violent poiſons. Without mentioning the oils of the common and cherry lau-

(a) The modification this expreſſion admits of, will be ſhown in the ſequel.

rel,

rel, the red oil of bitter almonds by diſtillation is a poiſon.

My next enquiry has been to know whether the venom of the viper is inflammable, that is to ſay, whether the phlogiſtick principle it contains is capable of taking fire. I have thrown it on burning coals. I have ſteeped a piece of paper and a bit of wood in it, and I have collected it in ſmall drops on the point of a needle: it has never taken fire, and I have not found it to be more inflammable than the other fluids of animals.

This obſervation holds good as to the venom of the bee, and thoſe of the waſp, hornet, and ſcorpion, which ſo far reſemble that of the viper. They all conſume and dry in the fire, without kindling into a flame.

If a pure and freſh drop of the viper's venom is applied to the mouth, it is found to poſſeſs a certain viſcoſity; but when it is dried in large drops on a bit of glaſs, it has the appearance of a tranſparent and yellow jelly: it then, like pitch, adheres ſo ſtrongly to the teeth, that it is with difficulty detached from them.

CHAPTER XII.

Peculiarities of the Venom of the Viper, and that of other Venomous Animals.

IT has been ſeen that, contraty to the opinion of Redi, the venom of the viper flows from the hole at the point of the tooth, and that it enters by the hole ſituated at its baſis. According to this diſpoſition one would be tempted to believe, that theſe teeth have been formed for the expreſs purpoſe of killing, ſo much does the ſmall hole at the point ſeem calculated to convey the venom into the blood of the animal bitten. But I do not here pretend to recur to final cauſes, and am very far from thinking that all this ſingular mechaniſm in the viper, has been expreſsly made for the deſtruction of other living creatures. The venomous liquor with which it is provided is perhaps neceſſary to its digeſtion, and I ſhall ſhow that it ſingularly diſpoſes the fleſh on which the viper feeds, to a ſpeedy putrefaction; a degree of change 'tis neceſſary for it to undergo for the purpoſe of being well digeſted. However, by an unlucky but neceſſary mechaniſm, the ſame tooth at once conveys the poiſon into the animal the viper bites, and into the aliments it feeds on. Who knows but that the depriving it of this venomous humour would expoſe it to acci-

dents

dents ſimilar to thoſe that happen to other animals, from a defect or depravity of ſome one of their digeſtive juices?

If it were true, for example, that the human ſaliva, as has been believed, is a poiſon to certain kinds of animals, and a philoſopher in this number, reflecting and reaſoning on its nature, ſhould obſerve that this ſaliva is one of the principal juices that concur to our digeſtion, would this reaſoning animal be miſtaken? Would it not, on the other hand, have divined nature? But if, on the contrary, one of theſe ſpecies of animals ſhould pretend that our ſaliva has been ſupplied us for the purpoſe of poiſoning them, ſince it actually does deſtroy them, would it not be fallen into a very abſurd errour? See however where thoſe incautiouſly hurry themſelves, who inceſſantly recur to final cauſes, in the examination and explanation of natural facts and events.

Finally, it is a general law of venomous animals that wound either with the tooth or ſting, to convey the venom into the wound by holes or orifices they have in thoſe parts. As to the ſcorpion, writers do not agree either on the number or ſituation of theſe orifices. Redi, by an inconceivable fatality, could never diſcover them; and as he had ſeen only a ſingle drop of venom on a plate of iron againſt which he had made a ſcorpion ſtrike its ſting repeatedly, he inferred from thence that there was a ſingle hole only at its extremity. Valiſnieri reckons as many as three. It is however very cer-

tain that thoſe of Tuſcany I have examined, never had more than two lateral openings, through which the venom flowed, and that neither a ſingle one nor three are to be found in them, as theſe two obſervers have maintained. When the ſmall veſicle which terminates the tail of the ſcorpion, and at which the ſting begins, is gently compreſſed, the two lateral apertures, and likewiſe the venom at the criſis of its flowing out of them, are ſeen with the help of a good magnifying glaſs.

But to return to the viper; its venom preſerves itſelf in the cavity of the tooth for ſeveral years, without looſing its colour or tranſparence. If this tooth is then put into warm water, the venom diſſolves very quickly, and is ſtill capable of killing animals. It beſides preſerves its activity for ſeveral months after being dried and reduced to powder, as I have many times experienced in common with Redi. It is here ſufficient that it is conveyed into the blood as uſual, by the medium of ſome wound: it muſt not however be kept too long, ſince I have frequently known it to loſe its effect at the end of ten months.

I do not heſitate to believe, that the animals the death of which is occaſioned by their touching the heads of vipers, even a long time after they are dead, are in reality ſimply poiſoned by the venom lodged in the cavity of the tooth, which being diſſolved by the blood of the wound, may have flowed out at the elliptical hole at the point of the tooth. A bit of dried venom that may happen to adhere to

the outer ſurface of the tooth is likewiſe capable of producing this effect. I am well aſſured by all the obſervations I have made, that the head of the viper dies in leſs than twenty-four hours, and that its muſcles dry in a few days provided they are in a very dry place, or ſoon become putrid if the place is wet. The teeth of the viper are beſides very ſharp, ſo that they pierce the ſkin, however ſlightly they touch it. I have twice ſucceeded in killing animals by ſimply wounding them with a tooth which had been plucked from a viper ſeveral hours before, and which was filled with coagulated venom. If the nephew of the famous Jaques the viper catcher, as Redi informs us, wounded himſelf ſeveral times in the hand ſo as to draw blood, with viper's teeth he had juſt plucked out, without his ever experiencing any other ill effects than what would have reſulted from the prick of a pin or of a thorn, he did not however at any time make the experiment without the greateſt riſk of there being ſome remains of this mortal poiſon in the tooth. The chickens likewiſe that Redi wounded in ſeveral parts of the body, with the teeth plucked from a living viper, all incurred the ſame riſk.

I do not deny but that the venom contained in the veſicle may be likewiſe capable of killing, even on the day following that on which the head has been ſeparated from the body of the viper. To effect this it will be ſufficient that the animal has not bit before it was killed, and that the head is

neither too dry nor too rotten, ſince in theſe caſes, the veſicle would either be deſtroyed, or could no longer convey the venomous humour to the tooth, by the excretory conduit already obſtructed and dried up.

From what has hitherto been ſaid, it may be conceived how certain mountebanks, according to the relation of the authour of a work on Theriaca, dedicated to Piſo, ſuffered themſelves to be bit by vipers with impunity. "There are men," ſays this authour, "who under the pretence of poſſeſſing "an antidote, have themſelves bit by vipers; they "previouſly give them a certain paſte which ſtops "up the holes in their teeth, and thus renders "their bites ineffectual, to the great aſtoniſhment "of the ſpectators, who are ignorant of the method "employed by theſe people to conceal their im-"poſture." This paſſage clearly ſhows us, that even in theſe times they had ſome knowledge of the ſtructure of the viper's teeth, and that they were of opinion, that the venom was carried by this hole into the wound. We likewiſe find in the work of Chryſogonus, entitled *De Artificioſo modo Curandi Febrium*, that this authour, who lived a long time after, was of the ſame opinion. Speaking of the viper, he ſays, "it has two teeth, the right and the "left, fixed in the lower jaw, and each of them "perforated; they are longer than the others, and "are ſhed every year when theſe animals quit their "ſkin: theſe two teeth are enveloped in two ve-

" ficles filled with venom, whence it flows into the " tooth by the hollow canal, the inſtant the viper " bites."

This authour ſeems only to have added miſtakes to what was known of the natural hiſtory of the viper before his time. It is falſe, for example, that it ſheds its teeth every year, when it changes its ſkin; it is falſe that the two veſicles ſurround the teeth; it is falſer ſtill that theſe two teeth are placed in the lower jaw. This alone proves fully that he never examined the mouth of the viper.

I endeavoured myſelf to have animals bit with impunity, and for this purpoſe prepared a paſte with pitch, turpentine, and yellow wax. I made two vipers bite ſeveral times at this compoſition, and they remained for ſome days without being able to do any miſchief. I found that their teeth towards the point, were indeed filled with this glutinous paſte, which ſtopped up the orifice out of which the venom ſhould have flowed.

I do not believe however that this method is a certain preparative againſt the bite of theſe animals. We have ſeen that there are circumſtances in which the venom may likewiſe paſs immediately from the excretory conduit into the ſheath. The ſureſt way then would be to entirely remove the reſervoir; and thus the mountebank would impoſe on the ſenſes of the vulgar with greater certainty, ſince he would no longer have any thing to dread from theſe dangerous animals.

There

There are excellent naturaliſts who believe that the fly, which they call in Tuſcany, *Aſſillo*, (the ox-fly) throws out a venomous and cauſtick juice from the end of the ſting it has at the extremity of its belly. Valiſnieri, who has written ſo well on this inſect, thinks that when it pierces the hide of the larger animals with this very ſharp ſting, it inſinuates into it a ſpecies of poiſon of a very corroſive nature, which irritates and as it were burns the tender filaments of the nerves of the part, ſo as to produce ſpaſms, throws their blood into an effervescence, and drives them to madneſs *(a)*.

Reaumur, that great and exact obſerver of the minuteſt animals believes, in oppoſition to the opinion of Valiſnieri, that this pain is rather the effect of a ſimply mechanical wound, than of a venom or any other cauſtick matter that the ox-fly may throw out of its ſting *(b)*.

The celebrated Morgagni, after having nicely weighed theſe two opinions, does not preciſely embrace either of them, but ſeems to combine them ſo as to form one opinion out of two. He maintains that the pain which the ſting of this fly cauſes to animals, frequently depends on two cauſes at the ſame time; that of a conſiderable nerve wounded by the ſting, and of an acrid and cauſtick venom which irritates the nerves *(c)*.

(a) Tom. II. Page 229. Venezia.
(b) Hiſtoire des Inſect. Tom. IV.
(c) De Cauſis et Sedibus Morborum. Lib. II.

The opportunity I had of procuring theſe flies, inſpired me with the wiſh of examining them. The ancients were acquainted with a fly that, with its ſting, threw whole herds into fury. The Greeks called this fly *Oeſtros*. The Latins have likewiſe mentioned a fly, the ſting of which produced the ſame effect on large animals. This they named *Aſſillus*. I do not doubt but that the *Oeſtros* of the Greeks, and the *Aſſillus* of the Latins, is the ſame with the *Tabanus* of Varro and Pliny. And although the ancients have diſcovered their uſual negligence in the deſcription they have given of this fly, it is however impoſſible not to ſee that it is no other than the *Aſſillo* of the Tuſcans, and the *Taon* (ox-fly) of the French. We muſt otherwiſe determine within ourſelves, that a fly which was ſo common amongſt the Greeks and Latins, has not deſcended to us, and that its ſpecies has been long deſtroyed and extinct. I flattered myſelf that I could find with eaſe the ſmall veſicle that contains the venom of this fly, and the hollow ſting that conveys it, as they are readily found in the bee, the waſp, and the hornet: I was however deceived. The ſting, much larger than that of the bee, is notwithſtanding neither hollow nor channelled, and I could never diſcover any cavity in it, either in its outer or inner part. I did not ſucceed better in finding the reſervoir of this pretended humour; in ſearching for which the ſtrongeſt lens were ineffectual; it was in vain that I compreſſed the extreme part of the belly and the root of the ſting, I could never perceive a fluxion

of

of this liquor, as it is ſeen in the bee, waſp, and hornet; and in a word, in all the animals that convey venom into the wounds made with their ſtings.

But to leave nothing undecided on this ſubject, I endeavoured ſeveral times myſelf, and engaged others in the ſame trial, to diſcover the venom by its taſte, by applying the ſting and the parts of the belly moſt adjacent to it to the mouth. I bruiſed it betwixt my teeth, and rolled it in my mouth, but could not find any thing acrid or burning in it, and did not feel the ſmalleſt pain or inconvenience. If it were however true, that this humour is ſo very acrid and cauſtick as to burn, as it were, the nervous filaments of the oxen, I certainly ought to have felt it on my tongue, ſince the venom the bee carries in its ſting, cauſes an intolerable ſmart and pain in that part.

It is falſe then that the ox-fly, at the ſame time that it pierces the hide of oxen, ſheds a poiſon. The pain it cauſes is ſimply mechanical, and ariſes from the particular ſhape of its ſting. This is compoſed of three ſmall, ſharp, and pointed hooks, of a horny ſubſtance, which when united together form a kind of pincers. The ox-fly does not uſually cauſe any great pain by its ſting, but if it accidentally wounds a large nerve or other ſenſible part of the animal, or if, which is more probable, it withdraws its ſting with fear and precipitation, and in a direction oppoſite to that in which it entered, it then happens that by tearing the ſkin and dragging the nerves forcibly with its hooks, it muſt neceſſarily cauſe that

very violent and inſupportable pain, which throws the herds into fury. We know how great a difference there is betwixt the ſlight pain cauſed by a ſharp inſtrument, and that excited by a weapon that tears and lacerates the nervous parts.

I have likewiſe had an opportunity of examining into the nature of leeches. There are naturaliſts who believe them to be venomous, becauſe the wounds they make are very painful, remain a long time open, and ſometimes cauſe a ſwelling of the adjacent parts. But it is clearly proved that theſe ſmall animals, ſo uſeful in medicine, are deſtitute of venom, and ſimply make a mechanical wound with the very ſingular weapon they have at the bottom of the mouth. This inſtrument is formed by three ſemilunar ſubſtances placed at the entrance of the oeſophagus, towards the centre of which their edges would meet each other, did not this cavity ſeparate them: they are placed perpendicularly in a direction with the length of the animal. The curved edges of theſe half-moons terminate in a horny ſubſtance diſpoſed in ridges, the diſtance betwixt which gradually widening, they at length form a kind of very fine teeth, like thoſe of a ſaw.

Theſe worms employ the following method in ſucking blood. They make a forcible application to the ſkin with the outer edges of their mouth. They then make a vacuum by enlarging that cavity in ſuch a way that the ſemilunar inſtrument approaches the ſkin, at which time they move the three ſaws circularly, and by ſucceſſively drawing them

73

them to and from each other, they make three notches in the ſkin, which unite in a ſingle point. In proportion as theſe ſaws recede from each other, the oeſophagus dilates and draws into its cavity the blood that has been pumped up.

I have tried what I advance here on myſelf. I applied a large leech to my arm, after I had cut away half its mouth, and was enabled in this way to view the whole of the mechaniſm at my leiſure.

The teeth and channellings of theſe ſaws are eaſily ſeen with a good microſcope. They may even be felt by paſſing the end of the finger over them; and by drawing the edge of a lancet acroſs them, particularly when they have been left to dry a little, may be heard to grate. In this ſtate they may be employed in ſawing the ſkin, provided they are held firm with pincers, or turned round with their edges conſtantly oppoſed to the part. I have even been able to effect this, notwithſtanding the ſoft parts of theſe ſemilunar bodies, ſuch as the muſcles, were not yet become dry. It is eaſy then to comprehend how the leech, after having contracted and ſtiffened the muſcles that form the greater part of theſe ſaws, contrives to pierce the moſt obdurate hide; and why it is that the wounds it makes are ſo very painful, and bleed for ſo long a time, ſince it only obtains this blood in conſequence of having torn with its ſaws, and made an opening in ſo ſenſible a part as the ſkin, and one ſo abundantly provided with nerves and veſſels.

I here

I here conclude the experiments, which, as I have obſerved in the beginning of this treatiſe, are the moſt certain guide to conduct us to the diſcovery and knowledge of natural truths. Facts alone are however not ſufficient to diſſipate the obſcurity that envelops them. A train of obſervations, without the help of a ſkilful hand to apply them, would be at beſt but the uſeleſs proof of a painful application. In the ſame way the moſt brilliant ſyſtems the rich and fertile imagination of a philoſoper can ſupply, do not deſerve the attention of naturaliſts, unleſs they are founded on good experiments. To come at the cauſes of the laws which regulate the courſe of the celeſtial bodies, nothing leſs was needed than the long ſeries of obſervations of the Chaldean ſhepherds, and the powerful aid of the creative genius of Newton.

CHAPTER XIII.

What cauſes the Death of Animals that have been Poiſoned by the Viper.

THE firſt object of my obſervations on the venom of the viper, was to diſcover the origin of the contradictions which, notwithſtanding they are attacked on all ſides by learned men of the firſt rank, are found in the various experiments that have been

been made on that ſubject. I muſt confeſs, however, that in verifying and analyzing all theſe particulars, my aim has likewiſe been to find in their combination, if poſſible, a ſatisfactory explanation of the ſpeedy and deadly manner in which this poiſon acts.

I ſhall aſk then, with Redi, "in what way the "venom of the viper extinguiſhes life, and brings "on death? Whether its action depends on a la- "tent cauſe beyond the reach of human intelli- "gence? Whether, on its penetrating to the heart, "it chills and freezes up the principle of heat; or "whether, on the contrary, by multiplying this "very principle and giving it more activity, it "kindles it afreſh and conſumes it, and in this way "diſſipates and reſolves the animal ſpirits? Whe- "ther it acts by deſtroying the ſenſation of this or- "gan? Whether, by the means of a painful irri- "tation it excites, the blood does not flow back "too precipitately to the heart, ſo as to bring on "ſuffocation? Whether it ſtops its motion, by "congealing the blood in its two ventricles, ſo that "they can no longer dilate or contract? and laſtly, "whether it coagulates, not only the blood in the "heart, but likewiſe in the whole venous ſyſtem?" "To reſolve theſe queſtions with truth," continues Redi, "is a taſk I am unequal to, and I "place them amongſt the infinite number of things "I now am, and ſhall probably always be, ignorant "of." Other authours, bolder than he, are not afraid of expoſing their ſentiments, whether badly or well founded. Before I propoſe mine, I think it ne-

neceſſary to relate the moſt reaſonable opinions that have been held by naturaliſts, as well ancient as modern, on this ſubject.

The learned Brogiani, profeſſor of anatomy at Piſa, has written a treatiſe full of erudition, on the venoms of animals. He there examines, as a ſkilful critick, the different ſyſtems and various opinions that have been eſtabliſhed on the mode of action of theſe poiſons.

It was at firſt believed that the venom, on entering into the blood, cauſed a univerſal coagulation of it, preciſely as acids do when they are introduced at an aperture made in a vein. The animals on whom this experiment is made, die in a very ſhort time, with tremblings, convulſions, and vomitings. On opening them afterwards, their blood is entirely coagulated in the veins, and as it has likewiſe been found coagulated in certain animals which, after having been attacked with the ſame ſymptoms, died of the bite of the viper, a trifling and hazardous inference has been drawn, that the venom brings on death by coagulation. But if, according to the teſtimony of Redi and the Memoirs of the Academy of Sciences of Paris, this appears not to be equally true of all the ſubjects that die of this poiſon; if it is likewiſe falſe, that they all have theſe tremblings, vomitings, and convulſions; if the blood is frequently found coagulated in this way, in every kind of dead bodies; it follows, that the queſtion yet remains undecided, and the difficulty as great as before: beſides, may there not

not be other circumſtances capable of coagulating the blood, and exciting the tremblings, convulſions, and other accidents, without recurring to the acid of the venom of the viper? My experiments themſelves have ſhown me that this acid does not exiſt, and that no ſtreſs ſhould therefore be laid on it.

Others have believed on the contrary, that this venom kills by exciting an univerſal inflammation. But how can it be thought capable of exciting it ſo as to occaſion death in ſo ſhort a time? I will go farther, and aſſert that the fever which conſtantly attends inflammation is not always found in thoſe that die of the bite of the viper. There are even no traces of inflammation in their dead bodies, and when any ſuch are found, it is rather the effect of ſome particular circumſtance in the temperament, than of a proper and peculiar quality reſiding eſſentially in the venom of this dangerous animal.

The diſciples of Hoffman, who, at the example of their Maſter, explain every thing by the atony and ſpaſm of parts, endeavour on this occaſion to avail themſelves of a truth to ſupport their opinion. They pretend that this poiſon excites, they know not how, an univerſal ſpaſm in the machine. But again, if this ſpaſm does not exiſt in all the animals that die of this poiſon, how can it be regarded as an univerſal cauſe? It is on the contrary certain, that they invariably die, rather from an atony and univerſal reſolution, than from the rigidity and contraction of their members.

I paſs over ſeveral other hypotheſes, which are nothing more than ſimple conjectures, and, far from being ſupported by any deciſive obſervation, are on the contrary belied by experience.

I however think it incumbent on me to relate the opinion of Mead. He ſets out on the exiſtence of cauſtick ſalts in the venom of the viper, and on this foundation grounds the whole of his theory of its effects. In the edition of his book on poiſons, printed in 1739, we find an ample detail of the different opinions of philoſophers, followed by a chain of ſyſtematick reaſonings, which, as any one may ſatisfy himſelf, are very tedious and filled with ſuppoſitions. His object is to ſhow that theſe ſalts decompoſe the globules of the blood, and deſtroy the temperament of it; and as it is difficult to comprehend how they can in this way deſtroy the whole maſs in ſo ſhort a time, he ſays that when once the venom has inſinuated itſelf into a wound, a very ſubtile and very elaſtick fluid riſes out of it, which in an inſtant extends its action to, and brings on a decompoſition of, all the parts, even the moſt diſtant ones, of this fluid. It is thus that a ſingle ſpark which touches a long train of gunpowder makes a rapid progreſs along it, and cauſes an univerſal exploſion, by the ſimultaneous diſengagement of the air encloſed by each particle.

It is without doubt unneceſſary to endeavour to combat this ſyſtem, ſince theſe pretended ſalts do not exiſt in the venom of the viper, and ſince nothing is falſer than the idea of theſe ſmall globules

bules of blood filled with elaſtick air. It is beſides certain that the venom does not alter the ſhape of theſe globules, which when they are obſerved with a microſcope, are found to be exactly the ſame as before, that is to ſay, obſcure and dark coloured at their circumference, and more tranſparent, at the centre, as ſmall round bodies generally are when viewed with a microſcope. I cannot conceive how Baker, otherwiſe very exact in his obſervations, could ſay in his *Treatiſe on Microſcopes*, that the bite of venomous animals, or even an atom of their venom, corrupts the whole maſs of blood, and alters the ſolidity and ſhape of the red globules that compoſe it.

It is not on this occaſion alone that a belief has been held without any foundation, of the change of ſhape of the globules of blood. The ſmall rings that have been endeavoured to be ſubſtituted to theſe globules, are a proof that the light, the microſcope, and the obſerver who relies upon appearances, are frequently the ſource of the pretended changes that do not in reality exiſt. I ſhall ſhow in a ſmall diſtinct work (*a*), that all ſmall globular corpuſcles, viewed with a microſcope, ſeem to be ſhaped like rings, becauſe the rays of light meet the eye of the obſerver in a greater number, from the centre than from the edges.

(*a*) The work announced here was printed ſome years ago at Lucca. It is entitled, *Oſſervazioni ſopra i Globetti del Sangue.*

The

The decompoſition of the globules of blood, ſo frequently advanced by phyſicians, is one of the rareſt phenomena in the animal economy. The phyſicians who are mechanicians, ſuppoſe that the globules of blood are ſo many ſmall round veſicles filled with a very elaſtick air encloſed in a fine membrane; they likewiſe believe that theſe globules (*a*) may eaſily crack and alter their ſhape, even from much ſlighter cauſes than that of the action of a cauſtick ſalt: but the fact is, that they are not veſicles, as they have perſuaded themſelves, and that they very rarely alter their ſhape.

Convulſions themſelves, that are ſcarcely ever felt by animals with cold blood, do not prove that the venom of the viper contains cauſtick ſalts, the inviſible points of which prick the nerves and irritate the muſcular fibres. Narcoticks and opium bring on convulſions, but muſt we therefore believe that they act by like mechanical agents? Still more, convulſions are not always the effect of an irritating *ſtimulus*; they rather ariſe from the deſtruction of equilibrium betwixt the antagoniſt muſcles. Weak languiſhing animals, that die from a loſs of blood, periſh in dreadful convulſions; and yet there are in this caſe neither points nor irritating ſalts. The convulſions are here likewiſe un-

(*a*) Let it not be underſtood that they are really globules; their true ſhape will be ſeen in a work of microſcopical obſervations which I propoſe to publiſh ſoon, and in which I ſhall ſpeak of whatever relates to their properties.

juſtly

juſtly attributed to the ſuperabundance of animal ſpirits; it ſeems more reaſonable to believe on the contrary, that it is to a defect of them, or to their irregular diſtribution in the muſcles, or rather to an irregularity in the circulation of the blood, that they owe their origin.

That opium cauſes convulſions is owing, in my opinion, to its deſtroying at different times and in an irregular way, the irritability of the muſcular fibres. It is beſides certain that men and women of a delicate and weak frame are always the moſt ſubject to convulſions; and it is not poſſible to ſuppoſe in theſe people a ſuperabundance of animal ſpirits. We know that all the muſcles, even in a relaxed ſtate, preſerve notwithſtanding a certain tenſion of their fibres, which, when they are cut, never fail to contract themſelves, and to enlarge the wound. When a muſcle becomes paralytick, it lengthens, and its antagoniſt then contracts the more; which ſhows that the repoſe of the muſcles depends on the equilibrium of ſtrength betwixt the different muſcles, and betwixt their different fibres. The powers thus balanced, deſtroy and renew themſelves at every inſtant, without producing any motion or ſenſible change. This natural tenſion of the muſcular fibres certainly depends on an equal and exact diſtribution of the fluids in the whole ſubſtance of the muſcles. This truth is demonſtrated in a diſſertation which I publiſhed in the third volume of the Acts of Sienna, which was in part reprinted ſome time after at Lucca, with ſeve-

ral conſiderable additions, and which was afterwards inſerted in the firſt volume of my animal phyſicks. [*Phyſique Animale.*]

But if theſe muſcles do not receive the ſame proportion of fluids, or if theſe fluids reach them, or are diſtributed amongſt them, with an unequal quickneſs and energy, the equilibrium of the mutual effort of the muſcles is immediately deſtroyed; the ſtrongeſt of them contract; and hence ariſe the convulſions and violent agitations of the whole frame. This is the reaſon why thoſe who die of an hemorrhage, as well as thoſe who periſh by poiſon, are ſeized with convulſions: for it certainly is not probable that the loſs of blood and of ſtrength ſhould bear an equal proportion in every part, in every muſcle, and in every fibre, whilſt the circulation itſelf is unequal, and the muſcular irritability is deſtroyed gradually, and in a very irregular way according to time and circumſtances.

But even though it might be concluded from the preſence of convulſions, that the matter which occaſions them is acrid and cauſtick, this would not determine it to be a ſalt; and becauſe ſalts prick, irritate, and corrode the nerves, can we ſay that they alone poſſeſs theſe properties? too few experiments have been made to warrant the maintaining this.

The convulſions ſome of thoſe are ſeized with who have been bit by the viper, do not furniſh a certain argument to explain the nature of the kind of jaundice that ſometimes attacks thoſe who die of

of this bite, or who ſicken with the diſeaſe of the venom. Some authours have aſcribed this jaundice to the contraction of the biliary pores at their origin in the liver, by which all ſecretion of bile being interrupted, the blood becomes charged with this humour, and depoſits the greater part of it in the organs of the ſkin.

Others have conceived, with greater appearance of truth, that theſe convulſions, and this violent irritation of the nerves, cauſe a conſtriction of the biliary ducts, ſo that the already ſeparated bile is carried into the blood, and ſpreads itſelf over the whole ſuperficies of the ſkin. Both theſe hypotheſes however are founded on a falſe principle, ſince anatomy teaches us that the nerves are not irritable, and that the biliary ducts are not compoſed of muſcular fibres. The firſt of them is abſurd on another account, for if the bile is not primarily ſeparated in the liver, and afterwards returned into the blood, how can it ſhow its quality and colour? I cannot conceive how very great naturaliſts have brought themſelves to think that it is not neceſſary for it to be ſeparated in the liver, to enable the blood to take a yellow tinge, and to give this colour to the ſkin.

It is not ſufficient that the blood contains all the ingredients of the bile, the fixed and volatile ſalts, the oil, and the water, to enable it to form bile. It is likewiſe neceſſary that the organs which concur to its generation, appropriate the matter of it, and regulate the proportions; ſo that the ſame

ſubſtances which in the proper viſcus might have formed bile, can never acquire, when mixed in the blood with the other principles of that fluid, either its nature or properties. But when once it is ſeparated, and thrown again into the maſs of blood, it preſerves its ſeveral qualities in ſuch a way, that all the principles of the blood can no longer decompoſe them, or break their combination. It may be compared to a drop of oil, which conſtantly preſerves its nature in the midſt of another fluid, although agitated and divided *ad infinitum*; each ſeparate particle continues to be oil as before. Thus for example, the principles of *muſt* (new wine) and of oil certainly exiſt in the vine and in the olive tree, but they only ſhow themſelves in the grape and olive.

A more appoſite circumſtance ſtill, and one which ruins this hypotheſis, is the example of eunuchs. The partizans of it will agree, that it is in vain for theſe unfortunate people to preſerve in their blood, during their whole lives, the principles that conſtitute the ſemen, ſince it does not manifeſt itſelf by any of its effects; they reſemble women, and never have the ſmell that characterizes the male. I will go further, and allowing that not only the principles of the bile are contained in the blood, but likewiſe the bile itſelf, it will not yet follow that it has the property of giving a yellow tinge to the ſkin. Animals have been known to have a ſcirrhous liver, or a very large abſceſs in that viſcus, for a long time, without being jaundiced.

diced. Let us agree then, that if thoſe who are attacked with the diſeaſe of the venom become ſo, the cauſe which produces this effect muſt have intercepted the courſe of the bile after its ſeparation in the liver, without its having done any previous injury to that ſecretion. I am firmly perſuaded that it does not thus pour itſelf into the maſs of humours, but becauſe its courſe is intercepted in the ductus communis choledochus, before it diſcharges itſelf into the duodenum. The convulſions of the ſtomach and inteſtines that attack thoſe who have been bit by the viper, may very readily irritate and contract the duodenum, and ſo ſtop up this orifice. Neither muſt we be aſtoniſhed at ſeeing the ſame jaundice make its appearance in thoſe who have taken other poiſons, ſince they alſo have the ſame convulſions, with a painful drawing together at the pit of the ſtomach, bilious and convulſive vomitings, a contraction about the navel, and other complaints in the abdomen. It may likewiſe happen in certain caſes, that the bile of thoſe that have been bit, may be ſo attenuated and exalted, that it may even penetrate through the ſubſtance of the liver, and immediately make its re-entrance into the torrent of the circulation, conveying the jaundice to the whole ſurface of the body. It is thus that, in conſequence of its being exalted in certain diſeaſes, it paſſes through the thickeſt membranes, and depoſits itſelf abundantly on the colon, duodenum, meſentery, epiploon, and peritoneum, on which, as may be found by opening

dead bodies, it beſtows its colour. It is well known that there are very few humours in the animal body that corrupt ſo readily as the bile; and we ſhall ſoon ſee that it is this principle of putrefaction particularly, which the venom of the viper conveys into animals.

But to return to the opinions of authours, as to the immediate cauſe of the death of thoſe that are attacked by the diſeaſe of the venom. The celebrated De Buffon maintains, in his great work of Natural Hiſtory, that the activity of the venom of the viper, as well as of other active poiſons, depends on thoſe microſcopical animalcules, which are diſcovered in the infuſions of vegetable and animal ſubſtances, and which he believes to be ſimple *organical particles*. I can certify that nothing like them exiſts, either in the venom of the viper, or in the other poiſons, whether of the animal, vegetable, or mineral kingdoms, particularly thoſe of the laſt. I have rendered myſelf very certain of this, by experiments made with the greateſt care, and in which I employed the ſtrongeſt lens.

The authour of a book, entitled, *On The Reproduction of Individuals*, [*De la Reproduction des Individus*] or rather Monſieur de Buffon himſelf, aſſerts, that the venom of the viper, as well as all other active and penetrating poiſons in animals and vegetables, can be nothing elſe than theſe organical particles; and he ſays, that the ſalts Mead obſerved are preciſely the ſame particles carried to their higheſt degree of activity. He likewiſe believes

believes that the pus of wounds is filled with theſe moving corpuſcles: but all this is without foundation. I have ſhown that theſe pretended ſalts are not found in the venom of the viper, any more than the particles ſuppoſed to be in motion. I have likewiſe examined all kinds of wounds, whether of a good quality, gangrenous, or cancerous, and have never been able to find the leaſt veſtige of theſe particles: I could only diſcover a quantity of ſmall unequal corpuſcles, more or leſs round, and ſwimming in a tranſparent liquor. But what appeared ſtill ſtranger to me, and which is however inconteſtible, theſe microſcopical animalcules are not found even in wounds of living animals, that come of themſelves, whilſt they are always to be traced in animal and vegetable ſubſtances, put to putrify in water, and expoſed to the air.

This illuſtrious French naturaliſt has been miſtaken then in all he has written on the nature and action of the venom of the viper, and of other poiſons. The acids ſalts of Mead which have never exiſted in nature, and the neutral ſalts of the ſame authour, which are not real, have been metamorpoſed by the fertile imagination of the elegant French writer, into what are ſtill more abſurd, organical particles, endued with motion.

It is falſe, that the corpuſcles which are ſeen with a microſcope in continual motion, in the infuſions of animal and vegetable ſubſtances, are ſimple organical particles, ſince they are real animals. It is falſer ſtill that theſe organical particles are ſeen

in the venom of the viper, and in other poiſons. No motion is obſerved in any poiſon whatever, and there is nothing that can give one the ſlighteſt ſuſpicion of the exiſtence of theſe particles there. It is beſides impoſſible that the ſalts of Mead can be the particles of Buffon, ſince theſe ſalts are merely imaginary. There is no greater truth in the exiſtence of theſe particles in the pus of wounds, ſince there is no motion in this ſubſtance. It is with regret that I ſee myſelf obliged to dwell on the errours of this writer, but his authority may eaſily miſlead thoſe who can only judge after others. How many are there who judge in this way! we may include in this number all thoſe who are not capable of immediately conſulting nature; who prefer hypotheſis to fact, and eloquence to truth: a ſevere and candid poſterity will, without doubt, be aſtoniſhed to find, that there have been philoſophers and naturaliſts in the eighteenth century, who, even in the moſt important particulars, have ventured to ſubſtitute conjecture to experiment, notwithſtanding that the latter would have been made with as much eaſe, as it would have been deciſive.

"Let indolent men," obſerves Senac, (a) "ſeek an amuſement in deviſing the ſecret ſprings "of nature, as obſcure politicians divine and re- "gulate what paſſes in the cabinets of princes: it "is a philoſophical delirium that only hurts the

(a) Traité du Coeur, page 29. Preface.

mind.

" mind. But where life is intereſted, if we are per-
" mitted to form conjectures, it is for the purpoſe
" of ſubmitting them to the proof of experiment,
" which ought to decide."

In this uncertainty, ſeeing that the opinions of the greateſt philoſophers are ſubject to the greateſt difficulties, I thought it expedient to have recourſe to my own obſervations. Neither of their ſyſtems is ſatisfactory, when we conſider the quickneſs with which the venom of the viper kills animals. I could not comprehend how creatures with cold blood, ſuch as the frog, are ſo ſoon deſtroyed by this poiſon, whilſt they ſurvive for ſo long a time the loſs of the heart, inteſtines, and other viſcera, and even that of the brain and head.

Doctor Mead, as we have already ſeen, aſſerted with the generality of philoſophers, in the firſt edition of his works, that poiſons, particularly thoſe of the animal kingdom, act on the blood, and are carried by this fluid into the innermoſt parts. But having paid attention to the quickneſs with which the venom of the rattle-ſnake cauſes death, this illuſtrious naturaliſt has changed his opinion in his laſt work on the ſame ſubject, and has ſubſtituted the animal ſpirits to the blood. He maintains then, that the primary action of the venom of the viper and that of other animals, is on the nervous fluid, which, being depraved by it, produces inflammation in the organs, and deſtroys the animal machine; ſo that the diſeaſe cauſed by theſe venoms, communicates itſelf to the whole body in

no

no other way than by the medium of the animal ſpirits, which finally vitiate the blood, with which they unite themſelves. The falſity of this hypotheſis of Mead will be demonſtrated by and by.

Nothing is leſs known than the manner in which this poiſon acts and brings on death; if we reflect however on the effects of opium, its mode of action may inſtruct and enlighten us a little on that of the venom of the viper. That vegetable juice begins by rendering an animal weak and torpid, and ſoon kills it by deſtroying the irritability of the muſcular fibres, as I have ſeveral times obſerved in animals with cold blood, and as the famous Haller demonſtrated a long time ago, even in thoſe that have the blood warm. The ſymptoms and accidents that follow the bite of the viper, do not differ eſſentially from thoſe I have juſt ſpoken of, and may at leaſt induce one to ſuſpect that the venom of that animal likewiſe kills by totally deſtroying the irritability of the fibres.

I recollect that being ſome years ago at Bologna, and reflecting attentively on the action of mephitick vapours, whether natural or artificial, I could not bring myſelf to be ſatisfied with all that the different authours have written on their nature, and on the proximate cauſe of the ſudden death they cauſe to animals. Some will have it to be owing to the exceſſive elaſticity of the air, and others aſcribe it to the total loſs of this ſame elaſticity: now theſe two hypotheſes are equally belied by facts, which prove on one hand, that the

changes

changes the elaſticity of the air may undergo in mephitick vapours, is never ſufficient to kill animals ſo ſuddenly; and on the other, that there are mephitick vapours in which the air abſolutely loſes part of its elaſticity. Others have conceived that this peſtilential vapour kills by irritating the nerves of the bronchiæ, and cauſes a criſpation and an univerſal conſtriction in the lungs, to ſuch a degree as to cloſe the paſſage of air, and to prevent their dilation. Laſtly, there are thoſe again who have conceived, that the vitriolick particles of the mephitick vapours act with a repulſive force againſt the particles of the animal fluids, ſo that the pulmonary veſicles, deprived of their animal ſpirits, fall into an abſolute ſtate of relaxation. It is however very certain, that even thoſe animals which live a long time without reſpiring, and without there being any circulation in the lungs, ſuch as frogs, and other animals with cold blood, and the generality of inſects, in which the circulation frequently remains for a long time intercepted, without endangering life in the leaſt; that all theſe animals and inſects, I ſay, are very ſoon killed in mephitick vapours. Beſides, the nerves are neither ſuſceptible of contraction nor of irritability, and the pulmonary veſicles are not formed of muſcular fibres. It is likewiſe certain, that there are mephitick vapours without ſulphur, ſmell, or taſte, and which do not contain either an acid or alkaline ſalt; and even though they ſhould contain any ſuch, we ſhould not comprehend the reader how they are capa-

capable of killing ſo ſuddenly, animals in which life is ſo tenacious, and which the knife, the fire, the very extraction of the heart, lungs, and all the other viſcera, and laſtly that of the brain, do not deſtroy without great difficulty. In conſequence of theſe conſiderations, I formed a determined reſolution to make artificial mephitick vapours, and to examine the effects of them on living animals. I collected the vapours of ſulphur in a recipient, in which I placed a frog; it died almoſt inſtantly, after making a few leaps, and being violently agitated. I opened it, and found all its parts flaccid and relaxed. The heart ſtill moved, but feebly and with great difficulty, and in a ſhort time entirely loſt theſe little remains of action. I endeavoured ineffectually to irritate, not only that, but likewiſe the other muſcles; neither of them would contract. I forced a needle into the ſpinal marrow, and ſaw with ſurprize, that it no longer awakened the motion of the limbs. The colour of the blood was changed to brown, but its globules ſtill preſerved their round and ſpherical ſhape.

I placed two other frogs beneath a glaſs recipient, into which I had introduced the vapour of a ſolution of iron filings in the nitrous acid. They died inſtantly. I opened them, and found the blood of a browniſh hue, and collected in the auricles. The heart was no longer in motion, and was inſenſible to ſtimulations. The fleſh was throughout flaccid, and had likewiſe loſt all irritability. On pricking

pricking the crural nerves, the legs remained motionleſs.

During this period, the celebrated Doctor Veratti likewiſe made experiments on artificial mephitick vapours. I aſſiſted at them myſelf, in company with other profeſſors, and they proved very conformable to mine. It reſults clearly from all theſe circumſtances, that mephitick vapours kill animals, by deſtroying the irritability of the whole muſcular ſyſtem. This is the immediate cauſe of their action, and the reaſon why theſe pernicious exhalations kill animals as it were inſtantaneouſly.

About the time when the firſt part of the preſent work appeared in Italian, (at Lucca in 1767) I found, as has been ſeen above, that artificial airs kill frogs by deſtroying the irritability of the heart; and the examination of the effects that mephitick vapours produce on living animals, made me conclude, that they occaſion death by deſtroying the irritability of the whole muſcular ſyſtem. But a celebrated Phyſician (Tiſſot) ſeems not to be of this opinion in his excellent work on the nerves. He there expreſſes himſelf in this manner *(a)*, "One of the greateſt modern naturaliſts "thinks that factitious airs abſolutely deſtroy the "irritability of the heart, and that their effects are "to be explained accordingly: but there is no

(a) Traité des nerfs, &c. T. i. Seconde Partie, Article des Effets des Poiſons, § 218. en note

"convey-

" conveyance by which their action can be carried
" to the heart. Fixed air, which kills when reſpi-
" red, being applied in the way of injection to the
" muſcular fibres of the inteſtines, revives their
" action, awakens the principle of life, and reco-
" vers ſick perſons at the point of death. Applied
" then to the muſcles themſelves, it excites their
" irritability, inſtead of deſtroying it."

This is not the place to ſpeak in an expreſs way of the effects of artificial airs on the living body. I purpoſe to do this in another work on reſpiration, which has been finiſhed for ſome time, and in which I ſhall relate the detail of the experiments I have made on this ſubject, and give my ſentiments on the cauſe of the death brought on in mephitick airs. However in the mean time I think myſelf under the neceſſity of obſerving, that the arguments of the learned Tiſſot have not hitherto been deciſive; that the queſtion remains in its original ſtate; and that it ſhould only be decided by having recourſe to experiment, to which an authority of ſo great a weight as this philoſopher's, is but too capable of preventing an application.

The firſt difficulty Tiſſot oppoſes is, that we do not know the channel by which mephitick airs deprive the heart of its irritability.

But it muſt be acknowledged, that the ignorance of one truth does not exclude the knowledge of another; and that we may know the effects without underſtanding the cauſes, and ſtill know the manner of acting. All human ſcience is of this nature.

nature. We know effects, of which we are entirely ignorant of the cauſes; and we know cauſes, of which the mode of action is abſolutely concealed.

The queſtion then is reduced to this; to determine by experiment, whether mephitick vapours deſtroy, or do not deſtroy, the irritability of the heart; and the difficulty propoſed above is of no weight, whether we know or not, the way in which this is brought about, provided the experiment be certain, and the illuſtrious writer oppoſe nothing which diſproves it.

I do not beſides ſee how we can be certain that there are abſolutely no channels by which the action of theſe vapours may reach the heart.

They deſtroy animals that are made to reſpire them. In theſe circumſtances there is an immediate communication betwixt the lungs and theſe vapours. Fluid ſubſtances are continually ſeparated from the lungs, and this viſcus may receive others, if they chance to act on it. There may be a real communication then betwixt theſe airs and the lungs, betwixt them and the ſubſtances that are ſeparated from that viſcus. But the lungs are known to receive the blood from the heart, and to convey it thither again. I do not therefore ſee why the communication, or rather the action, of theſe airs on the heart ſhould be impoſſible.

The other difficulty Tiſſot oppoſes is, that fixed air, which kills when reſpired, when immediately applied to the muſcular fibres of the inteſtines, revives their action, and cures diſeaſes; whence he deduces,

deduces, that when applied to the muſcles themſelves, it muſt neceſſarily excite irritability inſtead of deſtroying it, and conſequently cannot deprive the heart of its irritability.

But, in the firſt place, nothing is more common in medicine, than to find ſubſtances which, when applied to one part of the animal machine, act as a remedy; inſtead of which they occaſion diſeaſes, and even death, when applied to others. Several medicines, particularly in the claſs of poiſons, operate preciſely in this way; freſh examples of which will be given in the continuation of this work.

Electricity occaſions death by depriving the heart and fleſhy fibres of their irritability, as I have proved in my work on *Animal Phyſicks (a)*; and this ſame electricity is notwithſtanding one of the ſtrongeſt ſtimulants to the muſcular fibres that are known. It reſtores life by exciting irritability, in the very animals in which it had an inſtant before deſtroyed it. Amongſt all the ſtimulants that can be employed to call the animals back to life, that the electrical ſhock has thrown into a ſtate of inſenſibility, a proper application of gentle ſparks appears to me the moſt efficacious remedy.

In the ſecond place, the application of fixed air has a very different effect when introduced into the inteſtines, than when it is reſpired. In the firſt

(a) The firſt volume of this work, which I have already had occaſion to quote ſo often, was printed at Florence in 1775, and is entitled *Ricerche Sopra la Fiſica Animale.*

caſe its action is immediate; in the ſecond, it ſeems to need the aſſiſtance of the blood, to convey its action to the heart. Whence it follows, that its effects may be very different in theſe two circumſtances.

Theſe particulars naturally led me to think, that the venom of the viper likewiſe kills animals by deſtroying their irritability. I procured fifty of the ſtrongeſt and largeſt frogs I could meet with. I preferred theſe animals, becauſe they are livelier than others; becauſe they die with greater difficulty; becauſe they are more irritable; and laſtly, becauſe their muſcles contract even ſeveral days after they are dead.

I had each of them bit by a viper, ſome in the thigh, others in the legs, back, head, &c. Some of them died in leſs than half an hour, others in an hour, and others again in two, three, hours, or ſomewhat more. There were ſome again that were not affected, whilſt others that did not die, became nevertheleſs ſwelled. There were likewiſe others amongſt them that fell into a languiſhing ſtate, their hind legs that had been bit continuing very weak, and even paralytick. In ſome of them I contented myſelf with introducing cautiouſly into a wound, made with a lancet at the very inſtant, a drop of venom. Theſe laſt lived longer than thoſe I had had bit; neither of them however eſcaped. I conſtantly took the precaution to prevent the venom I introduced into the wound, being carried out by the blood that flowed from it. Some of

theſe frogs ſwelled very much, others but little, and others not at all. The wounds of almoſt all of them were inflamed more or leſs. There were ſome however that died very ſuddenly, without the ſmalleſt mark of inflammation. A ſhort time after theſe animals had either been bit, or wounded and *venomed* (*a*), the loſs of their muſcular force, as well as that of the motion of their extremities, was very evident. When they were ſet at liberty, they no longer leaped, but dragged their legs and bodies along with great difficulty, and could ſcarcely withdraw their thighs, when they were pricked with a needle, of the pain of which they ſeemed almoſt inſenſible: by degrees they became motionleſs and paralytick in every part of the body, and after continuing a very ſhort time in this ſtate, died.

I now opened the abdomen, and ſtimulated the nerves that paſs through it in their way from the vertebræ to the thighs. I employed the ſtrongeſt corroſives, but could excite no motion nor tremulus in the lower extremities. I pricked the muſcles with as little effect, and thruſt a long pin into the ſpinal marrow, without producing any motion or trembling either of the muſcles or limbs. In none of theſe parts, all of which had died at the

(*a*) I thought I might be allowed this term, to expreſs in one word, that an animal, or any part of it, had received the venom, or at leaſt that it had been applied to it. *Envenomed* would be the proper term, but cuſtom has given it a figurative and moral ſignification, which makes me afraid to apply it in its proper ſenſe. In a work of ſcience, the uſe of a new word ſhould be permitted, to avoid tediouſneſs or ambiguity.

ſame

ſame time, was there the ſmalleſt veſtige of life. The nerves were no longer the inſtrument of motion. The muſcles no longer contracted, and were no longer ſenſible to ſtimuli. The heart alone in ſome few of them continued to move languidly, and its auricles were ſwelled and blackened by the blood with which they were ſurcharged. This organ did not however ſeem to have ſuffered much from the activity of the venom. It continued its motion, notwithſtanding the entire death of the other parts, and renewed its vibrations on being ſtrongly ſtimulated with needles. This motion and theſe oſcillations were however but of ſhort duration after the death of the animal.

Perſons have been ſometimes met with, who having been bit by a viper, have remained paralytick in ſome particular part of the body during life. A ſhort time ago a woman in Tuſcany, who had been bit in the little finger by a viper, became after various other complaints, paralytick throughout the whole right ſide of her body, and could never be cured. In a word, it is certain that all thoſe who have met with this accident, complain ſoon after of an univerſal weakneſs. Their muſcles refuſe their office. They become dull and heavy, have no longer the free exerciſe either of body or mind, and fall inſenſibly into a kind of lethargy: ſo true it is, that this venom induces a palſy of the muſcles, and robs them of that active property, called by the moderns, animal irritability. In the continuation of this work, I ſhall ſhow what opi-

nion ought to be held of that ſyſtem, and the changes I have made in it.

Thus then it appears, that animals die of the bite of the viper, from their fibres loſing that irritability, which is the grand principle, both of voluntary and involuntary motions in the animal economy (*a*).

From theſe experiments on frogs, it ſeems that the venom of the polypus is very analogous to that of the viper. Scarcely has a polypus ſeized an earth-worm, when it periſhes, and has no longer any motion: theſe worms are known however to be very tenacious of life, and to move a long time after they are cut in pieces. Let us ſay then, that the venom of the polypus (for it is one, ſince it kills ſuddenly, and in a very ſmall doſe) attacks the animal irritability, and extinguiſhes life, preciſely as does that of the viper.

After having found that the venom of the viper occaſions death by deſtroying the irritability of the fibres, let us examine what are the changes that happen to the muſcles after they are deprived of this property. It has been conſtantly obſerved, that the fleſh of animals loſes its motion and irritability in proportion as it has been penetrated by a putrefactive principle. We have many examples to prove that the loſs of the former invariably accompanies the firſt progreſs of the latter. Mephi-

(*a*) The propoſition I advance here is a very general one; the different modifications it is capable of will be ſhown hereafter.

tick

tick airs, which deſtroy irritability, likewiſe haſten putrefaction, and the muſcles of the animals that are killed by them, become flaccid and livid. We likewiſe find that thoſe of the animals bit by the viper become putrid in twenty-four hours. In both caſes the very principles of the elementary fibres are attacked, and the diſunion of theſe occaſions the loſs of their moſt innate natural properties. This disjunction of parts, which the putrefaction of the muſcles invariably cauſes, muſt neceſſarily deprive the latter of their irritability and fitneſs for motion.

I am led to think that the venom of the viper produces a ſomewhat ſimilar effect, and I found my opinion principally on the analogy of the other poiſons. Indeed we find that the fleſh of animals which has been cut with a knife dipped in the juice of napel, inſtantly becomes more tender, and fitter for culinary purpoſes. Travellers inform us that in both Indies, as well as in Africa, the inhabitants uſually hunt with poiſoned arrows, and that in the ſpace of ſix minutes, or more or leſs according to the degree of the poiſon's activity, theſe arrows kill the largeſt animals, ſuch as lions, tigers, and even elephants. They likewiſe obſerve that the fleſh of theſe animals immediately ſoftens and becomes tender; an unequivocal proof that all theſe poiſons equally diſpoſe the fleſh to a ſpeedy putrefaction. I have myſelf obſerved the ſame thing, in frogs and other animals bit by the viper. Their fleſh ſoftens much ſooner than uſual, to ſuch

70463

a degree as to crumble at the leaſt touch; it ſeparates of itſelf from the bones, and corrupts and ſmells in a very ſhort time.

If after theſe obſervations it is almoſt impoſſible to deny that the venom of the viper deſtroys irritability by conveying a putrefactive principle into the fleſh of animals that have been bit, and into their fluids, we muſt agree as to the inutility of having recourſe, at the example of mechanicians, to cauſtick, ſtimulating, and inviſible ſalts, in explaining the action of this venom. Very far from favouring this action, we know that ſalts are in general better calculated to ſuſpend and ſtop it; and I cannot conceive how naturaliſts, otherwiſe very enlightened men, have imagined and wrought themſelves into a belief, that the poiſons drawn from animals, and even thoſe from vegetables, owe all their activity to certain ſalts of this nature. Beſides, we ſcarcely find the ſmalleſt trace of ſalts in the juices of ſome of theſe plants, even the moſt venomous of them. I have examined ſeveral of them with the microſcope, and have not the ſmalleſt idea of having found any ſalts in them, except in the *toxicodendron*, in which tree, as in other plants, we only find a few ſhining globules, ſwimming in a more or leſs tranſparent fluid. What I am very certain of is, that there does not exiſt in the venom of the viper the ſmalleſt veſtige of thoſe formidable ſalts, that have been ſuppoſed capable of killing animals the moment they are introduced into the blood.

It

It is the facility then with which, by the help of theſe pretended ſalts, the action of poiſons is explained, that has ſeduced theſe mechanick phyſicians. They have conceived to themſelves *ſpiculæ* throughout, calculated to diſunite the animal fibres, and decompoſe the humours. But what will they reply to the example of opium? It kills by weakening, by even deſtroying, the irritability of the fibres. If the virulence of this vegetable juice reſides eſſentially in its gummy, and reſinous part, will they likewiſe ſuppoſe the exiſtence of ſalts there? Theſe hypotheſes have had their birth in a chymical laboratory, and are not the reſult of conſtant obſervations of the phenomena of nature. We muſt agree that theſe imaginary ſalts have been but too much abuſed. There are thoſe who have not heſitated to place them every where, and who have even gone ſo far as to believe that they alone are capable of awakening the ſenſes of taſte and ſmelling, whilſt nothing is leſs demonſtrated than the preſence of theſe ſalts in ſapid and odoriferous ſubſtances. Beſides, they do not conſider that ſalts are capable of altering their ſhape without loſing their natural taſte; how then can they likewiſe change their taſte whilſt they preſerve the ſame ſhape? It is not therefore on a certain determinate ſhape that their action muſt be made to depend, as certain naturaliſts will have it, who, when they ſet about explaining the ſenſations, ſee nothing throughout beſides edges and points; in an infinity of caſes this is not only ſuppoſed, but is likewiſe belied by experience. If there is only need

of awakening the ſenſations in ſome of our organs, why is there ſo great a neceſſity for theſe ſalts? Cannot this be brought about without their aſſiſtance? Have not other particles of bodies likewiſe the properties of contact and mechanical ſtimulus? Are light and air ſalts, becauſe they ſtrike the eye and the ear? A ſubſtance of any kind that acts on a nerve, may drag and relax the medullary ſubſtance, and may either compreſs or irritate it, independent of the cauſe that afterwards conveys the impreſſion to the mind or brain. If all the external ſenſations are reduced to a change in an organ, other bodies may then operate as well as ſalts. A fluid may likewiſe relax the tender parts of a nerve laid bare, and may equally ſhrivel and dry them. There are ſpirits and oils that dry and harden the fleſh of animals, and irritate the nervous and muſcular ſyſtem, and notwithſtanding contain no ſalts. In the ſame way, poiſons may kill without ſuppoſing ſalts throughout, in the three kingdoms. May not an action of one body upon another exiſt without the aſſiſtance of wedges and points? Can any one ſay, that ſalts are found every where where theſe figures are met with? or that they pre-exiſted in all the ſubſtances whence chemiſtry at length ſucceeds in extracting them? There is no more need of this, than to ſuppoſe that there are ſalts and points in camp and jail fevers, in the ſcurvy, and in a word, in all putrid diſeaſes, where the corruption of the ſolids and fluids is alike general. We muſt have recourſe to ſomething very different from ſalts to explain the deſtruc-

deſtructive force of theſe hazardous diſeaſes which overturn and deſtroy the whole animal economy in ſo ſhort a ſpace of time. Their effects, and thoſe of many other diſeaſes analogous to them, as well as the ſymptoms that accompany them, are well calculated to lead one to believe that they convey a latent *virus* into the machine, which, like the venom of the viper, brings about the deſtruction and univerſal decompoſition of the ſolids and fluids. Indeed theſe diſeaſes are invariably obſerved to be attended with convulſions, great faintneſs, a proſtration of ſtrength, drowſineſs, an exceſſive ſtench exhaling from the yet living body; and laſtly, a ſpeedy putrefaction, which almoſt immediately follows death. The very ſudden failure of vital ſtrength in the whole muſcular ſyſtem, is a certain indication that the diſeaſe attacks the animal irritability, and the principle of motion in the fibres. It is only in this way that, without having recourſe to ſyſtems, and to free and arbitrary hypotheſes, we can comprehend and explain how it is that the ſeeds of death are capable of ſpreading themſelves in an inſtant over the whole animal economy.

I preſume that it will not be poſſible for the future to entertain any doubt as to the true proximate cauſe of the death brought on ſo ſpeedily by the venoms of the viper and aſpick; and amongſt the three ſpecies of the latter, principally of that called *nintipolenga zeilanica*. This aſpick kills by occaſioning a ſudden drowſineſs and univerſal weakneſs, followed by death, in the animal ſtruck by it. In a word,

a word, it ſeems that all the poiſons ſupplied by the animal kingdom, occaſion death by deſtroying the irritability of the muſcular fibres, and diſpoſing both ſolids and fluids to a ſudden corruption. The ſame may be ſaid of thoſe vegetable poiſons, that are no ſooner introduced into the blood, than they are ſucceeded by death.

But of all the poiſonous animals hitherto known, the polypus ſeems to poſſeſs the moſt powerful and active venom. However irritable theſe creatures may be in other caſes, and difficult to kill, it ſucceeds inſtantly in extinguiſhing the principles of motion and life in water worms. What is very ſingular, its mouth or lips have no ſooner touched this worm than it expires, ſo great are the force and energy of the poiſon it conveys into it. No wound is however found in the dead animal. The polypus is neither provided with teeth, nor any other inſtrument calculated to pierce the ſkin, as I have aſſured myſelf by obſerving it with excellent microſcopes.

Let us likewiſe be very cautious how we believe, at the example of many naturaliſts, that life conſiſts in general in the circulation of the blood and motion of the heart, and that it abſolutely ceaſes when this circulation is interrupted. The circulation is not general in animals; polypuſſes have not even a heart or other analogous viſcus, to bring about its operations. It is proved too, that ſeveral animals with cold blood live a long time without heart, and without viſcera, as is ſeen in frogs, turtles, and ſeveral

veral kinds of fiſh and worms, in which, although the circulation is undoubtedly then ſtopped, they continue notwithſtanding to live and move, and are wrought on by their paſſions as uſual, appearing to be ſtill ſubject, to and ſenſible of, the wants of life.

I have found many animals, inſects, and worms, in which there is certainly no kind of circulation in the veſſels; there are others, in which it is only imperfectly carried on in ſome particular parts of the body, and not at all in the extremities. I purpoſe to give all theſe particulars to the publick, in a work I have been ſome years buſy in getting ready, entitled, *Sur les Animaux Microſcopiques (on Microſcopical Animals)*.

This errour has ſpread itſelf amongſt philoſophers, by the help of a falſe analogy they have ſuppoſed betwixt animals with warm blood, and thoſe with cold; a very dangerous mode of reaſoning in phyſicks, and belied at every ſtep by obſervation and experiments. A function has been obſerved to be executed in a certain way in animals with warm blood, and it has been immediately concluded to be the ſame in all others. Thus are general laws made, and propoſitions on ſo extenſive a ſcale advanced, merely becauſe nature has not been ſufficiently conſulted. We have needed a *Tremblei* and a *Bonnet* to tell us of theſe general axioms, and of the idea of a neceſſary and common law in the generation of all animals.

I can-

I cannot forbear mentioning in this place the ſingularity of motion of a ſmall microſcopical animal, which Lewenhoeck has named *rotifer* (wheel-polypus). All the obſervers, even the moſt modern ones, that have ſucceeded him, have believed that this animal has real wheels (*a*); but to be certain of the contrary, it is only neceſſary to place it betwixt two pieces of glaſs, and then obſerve it with an excellent microſcope. 'Tis a ſmall gelatinous worm, commonly found in the earth or ſand collected by rain in the tops of houſes. I have likewiſe found it in other earths, as well as in waters that have been ſometime ſtagnant, and more frequently again in thoſe that have a very gentle current, and are filled with *conferva* and other aquatick plants. This worm is divided towards its head into two pretty large trunks, which appear like two wheels or ſtars, from the number of ſmall, extremely ſharp, and ſhort, branches that are attached to their circumference. They really appeared to Lewenhoeck to be wheels of a rare mechaniſm, and every one would judge the ſame, on ſeeing the creature put

(*a*) Great care ſhould be taken not to confound what we imagine, with what is pointed out to us by obſervation. Indeed there have been authours who, either guided by analogy, or puzzled to explain ſo ſingular a motion, have ventured to aſſure us that theſe wheels are not real; they have luckily ſaid the truth. It muſt however be agreed that we ought to obſerve, and not to divine, the phenomena of nature. Whoever gives himſelf up to reſearches of this kind, without the faithful guide of obſervation, runs the greateſt riſk of falling into error.

them

them in motion. But a more exact obſervation at length convinced me that they are not wheels, but compoſed of a quantity of ſmall moveable arms, formed like pointed cones, and planted all round the two trunks. It lets fall theſe moveable arms or rays ſucceſſively, and afterwards raiſes one after the other with ſo much celerity, that the eye fancies they are turning round like the ſpokes of a coach-wheel, or rather, like the branches of a wheeled fire-work. It never moves theſe two wheels, except when it ſwims or wiſhes to eat, and theſe two ſtates are invariably the ſhorteſt of its life. In ſwimming, it ſtrikes the water with theſe arms or branches with great celerity, reſts itſelf at different periods, and thus tranſports itſelf from one place to another. When it eats, it, on the contrary, fixes its tail in ſome ſubſtance, and afterwards turns its two wheels, giving ſuch a motion to the water, that it directs the courſe of it towards its head, ſo that it preſents to its mouth all the ſmall corpuſcles with which it is filled. The velocity of the motion of its arms or wheels is incredible; but what is ſtill more aſtoniſhing, is the motion of its heart. This viſcus is ſeen very diſtinctly with a microſcope, and can never be confounded with any other part of the animal whatever. It is abſolutely immoveable when the worm does not play its wheels; but no ſooner are they in motion, than the heart moves too, and its action becomes ſtronger in proportion to the quickneſs with which the wheels are agitated, ſo that their motions are always in an exact proportion. I do

not

not take upon me to deny, but that it ſometimes happens (although very rarely, and that at very long intervals) that the heart is in motion even whilſt the wheels are at reſt; and as the motion of the wheels is always at the diſpoſition of the animal, ſo likewiſe is that of the heart. The heart then is a voluntary muſcle, depending on the will of the animal; a circumſtance which is at the preſent time *unique*, having never been obſerved in any other caſe. The wheel-polypus paſſes the greateſt part of its life then without any motion of its heart, and conſequently, without a circulation of blood, or of a fluid which receives motion from this muſcle. This does not, however, prevent it from moving during the other intervals, when it creeps and trains itſelf, as worms do, amongſt the bodies that ſurround it.

An objection may be ſtarted here, that this organ of the wheel-polypus is not the heart of the animal, but rather its ſtomach, ſince it is obſerved to move when the creature eats; and that it is altogether extraordinary to ſuppoſe, that the heart is a muſcle ſubmitted to the will, whilſt it is not ſo in any other animal. It muſt be confeſſed, that this is not impoſſible, but it is not, on that account, very probable; and even though it ſhould be true, it would be likewiſe true, that an organ ſuch as the ſtomach exiſts with a voluntary motion, which, any more than in the other caſe, is not obſerved in any other animal.—Thus the difficulty I encounter is of no weight, ſince it muſt always be agreed,

that

that a muſcular organ exiſts in this animal, which, in oppoſition to thoſe of all other animals, is ſubordinate to the will. This is preciſely what I wiſhed to prove by my obſervations, and my diſcovery, therefore, ſtill remains ſuch. It is likewiſe to be obſerved, that the rotifer puts this ſingular organ in motion, even when it does not eat; that is to ſay, at a time when it can make no uſe of it, provided it be its ſtomach. This happens every time it ſwims, or wiſhes to paſs rapidly from one place to another: it has then occaſion to move its two wheels, and this organ moves in conſequence. Hence we ſee, that the animal does not move it to eat, but that the motion of it neceſſarily takes place when it plays its two wheels, whatever may be its motive for ſo doing.

But ſince it is certain that the voluntary motions of the muſcles of animals with cold blood, do not depend more on the circulation of humours, than does the irritability of the fibres, which ſeems to be the ſource and principle of life and motion in the animal, it follows that, in animals, life conſiſts in the action of their muſcles and parts; for the moment this motion ceaſes, the animal ceaſes likewiſe to live; and its body then, as to life, differs no longer from the ſtate of any foſſil or vegetable ſubſtance whatever; and all this aſſemblage of veſſels, ſo many different organs, and this aſtoniſhing ſtructure of its parts, are no longer of any uſe to the animal, and ſhould be regarded as if no part of them any more exiſted: motion being once at

an end in the machine, ſenſation and life are ſo too. The animal will return to life as ſoon as its parts regain their former motion, inſtead of which, it dies for ever, when, as happens to man, its parts not only loſe their actual motion, but likewiſe the faculty of recovering it in the ſequel. Thus the microſcopical eels that are found dry and withered in ſmutty wheat, recover motion and life as ſoon as they are wetted with a little water, and again become lifeleſs and dry, whenever they are no longer moiſtened. I have repeatedly aſſured myſelf of this with an extreme pleaſure. Thus, then, do they preſerve the power of reviving and reſuſcitating effectually, by the ſimple preſence of the water with which they are moiſtened.

The celebrated Bonguer, in his work on the ſhape of the earth, relates to us, from the teſtimony of Father Gumillo a Jeſuit, and alſo of the Indians of Peru, that a large venomous ſnake is found in thoſe countries, which being dead, and dried in the open air, or in the ſmoke of a chimney, has the property of coming again to life, on its being expoſed for ſome days to the ſun, in a ſtagnant and corrupted water.—It were to be wiſhed, that ſuch a naturaliſt and philoſopher as Monſieur Bonguer, could have verified by his own proper obſervation, a fact ſo important in itſelf, and rendered ſtill more ſo by the ſize of the animal.

I have dried the worm called *ſeta equina*, or, according to Linnæus, *gordius*, ſeveral times in the open air, without leaving it there too long: it had

had loſt almoſt all its bulk and weight, and was become like a bruiſed and dry ſtraw: its ſkin had ſhrunk ſo as to leave no ſenſible cavity, and it had no longer any ſign of life or motion. I returned it into water, where, in leſs than half an hour, it recovered its bulk and weight, and ſoon afterwards diſcovered unequivocal and permanent ſigns of life.

The wheel-polypus I have ſpoken of above, likewiſe loſes, when dried, every kind of motion and life, and recovers both one and the other when again put into water. Laſtly, I left it, by way of experiment, in a very dry ſoil, and expoſed, during the ſummer, to the whole heat of the ſun, for the ſpace of two years and an half. I afterwards returned it again into water, where, at the end of two hours, it recovered life and motion. I put one of them on a bit of glaſs, which I expoſed, during a whole ſummer, to the noon-day ſun: it there became ſo dry, that it was like a piece of hardened glue. A few drops of water did not, however, fail to reſtore its motion and life.—I have ſince found a number of other ſmall animals, either on the tops of houſes, in earths, or in water, which, in the ſame way, alternately loſe and recover the uſe of their organs, on being dried, and afterwards returned again into water. I purpoſe ſpeaking of theſe little prodigies in a work apart, to be entitled, On the Life and apparent Death of Animals *(a)*.

But it is not the ſame as to the loſs of irritability in the muſcles of the animals poiſoned by the viper.

(a) De la Vie et de la Mort apparente des Animaux.

Theſe remain flaccid, and their motion is loſt for ever. It appears very certain, that the venom of this animal differs but little from opium as to its effects; and that its mode of action on the fibres comes very near to that of this vegetable juice. Both of them excite violent convulſions and vomitings. Each of them conveys an univerſal debility into the organs. They render the muſcles paralytick, make the animal heavy, and finally bring on a ſpeedy death, by deſtroying the irritability of the fibres. The heart alone in both caſes ſtill preſerves its irritable quality for ſome time after the death of the other parts. It avails nothing here to animals with cold blood, that they are endued with an obſtinate life, and are capable of preſerving that, as well as motion, after they are cut in pieces. If either of theſe poiſons attacks the principle of their motions, and deſtroys the irritability of their muſcles, they die ſpeedily, all motion is annihilated in them, and their parts will no longer give any ſign of life. Their body, it is true, will preſerve its organization; but an organized body that has loſt its motion, is truly a body without life.

It is evident, then, that neither of the numerous hypotheſes naturaliſts have invented, a great part of which I have taken care to relate, explain in a reaſonable way the death of the animals poiſoned by the viper; but that its venom kills in no other way, than by deſtroying the principle of motion, the only ſource of animal life, in the different parts. I am the more attached to the opinion that the venom of the viper acts in no other manner than by deſtroying

ſtroying the irritability of the muſcular fibres, from having already ſhown, in a memorial printed in the Acts of the Academy of Sienna, that the nervous fluid is by no means the true, the efficient cauſe, of muſcular motion; and even though I ſhould be of another opinion, and ſhould regard the animal ſpirits as the cauſe of irritability, and the true principle of all the motions in the animal economy, my diſcovery of the proximate cauſe of the death of animals that have been bit by the viper, would loſe no part of its importance; for whether it operates immediately on the nervous fluid, or on the muſcular fibres, it is not leſs true that this venom kills by depriving the animal of all motion, and the muſcles of the power of contracting.

Unleſs I am deceived, I have now, I think, happily terminated the controverſies that have ſo long kept people at variance, on the mode of action of the venom of the viper. I believe I have explained how it is able, in ſo ſhort a time, to kill even the animals that are the moſt obſtinate in dying. When once this poiſon is introduced into the blood, it deſtroys the irritability of the muſcular fibres, the ſource and principle of all the motions, not only during the life of the animal, but alſo after its death. I call every animal dead in which there are no longer any exteriour ſigns by which we can ſay that it lives; and, in truth, it is only according to our ſenſes, and the information they give us, that we can judge of the true death of animals; that is to ſay, of the preciſe inſtant when they ceaſe to

exiſt, and are no longer alive. Indeed, how can we conceive a living being, without the idea of ſome motion in its organs? We ſhould otherwiſe introduce into phyſicks a ſenſeleſs pyrrhoniſm, and caſt trouble and uncertainty on the moſt certain and moſt received notions and ideas. A principle of corruption penetrates into, and ſpreads itſelf in, the ſolids and fluids, relaxes and decompoſes the muſcular fibres, and cauſes them to loſe the power of contracting. It is to this general law of putrefaction, then, and to this univerſal principle of diſſolution and death, that the entire action of the venom of the viper on organical bodies is reduced; and to this we muſt confine ourſelves, ſince in effect what is called the ſcience of nature has its bounds, and ſince it is not permitted us to go beyond them. Whatever this ſcience may be, if it is true that a putrefaction exiſts in nature, and that it effects the deſtruction of all organized bodies, it is likewiſe certain that we are entirely ignorant of its mechaniſm. Indeed when theſe bodies are ſubmitted to its action, who is capable of informing us what is its mode of working on them, with what power it operates, and laſtly, what changes and what revolutions it cauſes them to go through? The prodigious number of ſmall movements that are exerciſed on parts of an infinite minuteneſs, are too obſcure for us, and eſcape our ſenſes. It is, however, enough, to ſee that a general principle of putrefaction and deſtruction, which decompoſes organical bodies, and gives them up to death, reigns throughout

out nature. To ſeek an acquaintance with nature, we have nothing more to do, than to aſſemble the effects or particular accidents of bodies, and to compare them with other more general effects, that are called *principles, or laws, of nature.* This alone is what the great Newton did, when he ſubmitted all the celeſtial motions to the general law of gravitation. Of what conſequence is it beſides to an aſtronomical obſerver, to know the cauſe of the reciprocal tendency of the bodies moving in the ſky? Such a knowledge would be rather an object of curioſity to man, than a real advantage to aſtronomy.

Theſe were my ſentiments when I publiſhed this Firſt Part in Italian, thirteen years ago. I have at preſent made but very few changes and additions, becauſe the ſucceeding parts are, rigorouſly ſpeaking, merely a ſupplement, ſerving to correct what is previouſly advanced; and becauſe order would then have obliged me to give conſequences, which could not eaſily have been comprehended, till after a general idea was formed of the ſubject.

Deſtroyed irritability in the living machine was what I moſt conſtantly obſerved at that time. It is for this reaſon that I have reduced the action of the venom of the viper to this general principle, and have entirely excluded the nervous ſyſtem. I muſt confeſs, however, that the number of my experiments was then very limited, and that I had not varied them ſo much as I have ſince done. I was likewiſe ignorant of the faculty of the poiſon *Tithymalus*; and alſo of the ſurprizing effects of the oil of

the Cherry Laurel, for the greater part unknown to other obſervers.

I have likewiſe, in this Firſt Part, paſſed in a very curſory way over ſeveral other ſubjects, and have even given into a few ſuppoſitions, which I ſhall diſcuſs in my *Microſcopical Obſervations (a)*. The principal objects of theſe obſervations will be, the figure and properties of the globules of blood; the animals that are capable of dying and returning to life, which will give me occaſion to enter into a complete hiſtory of the celebrated eels found in ſmutty wheat; and, laſtly, the cauſe of the death of animals ſurrounded by artificial and unrenewed airs.

(a) Obſervations Microſcopiques.

PART II.

CHAPTER I.

On the Source of many Errours.

THE *ignorance* of a truth in phyſicks may hide from us the cauſe of a natural phenomenon; but *errour*, when it ſupplies the place of truth, ſtops the progreſs of ſcience, and ſubſtitutes dreams and chimeras to facts and nature. It is at all times a misfortune to be ignorant of a truth; but when we are ſenſible of our ignorance, we may ſtill hope to come at a knowledge of it. The book that of all others would be the moſt uſeful to mankind, is ſtill wanted. It is one which would at once determine what we really know, and what we do not know, although we have perſuaded ourſelves that we do. Our reaſonings would no longer have hypotheſis and errour for their baſis, and inſtead of fabricating ſyſtems, we ſhould endeavour to prepare materials. Nature would be more conſulted; we ſhould reaſon leſs, and know more.

There are errours and truths which concern man more eſſentially than others do. Theſe are, parti-

cularly, thoſe which relate to the preſervation of his ſpecies.

Man is naturally ſubject to certain diſeaſes; whilſt others again are accidental to him. Medicine employs itſelf in each claſs, and in ſeeking the remedies, renders itſelf uſeful to ſociety.

Thoſe who have diſtinguiſhed themſelves in this reſearch, cannot be ſufficiently praiſed. Poſterity will do juſtice to their labours, and immortality is ſecured to them. But, on the other hand, who does not ſee the miſchief that a remedy propoſed in very violent diſeaſes may be productive of, if, inſtead of being ſalutary, it is entirely uſeleſs, or even hurtful? To paſs ſlightly over ſo important a matter, would be expoſing mankind to the worſt of miſchiefs. The more certain we think ourſelves of the remedy, the more we deſpiſe the danger, and do not endeavour, ſo much as we ought to do, to guard againſt it. The diſeaſe attacks us, we neglect the aſſiſtance of art, and frequently become the victims of our own credulity, and of the ignorance of others.

The perſuaſion we are in that a diſcovery is made, blunts the ſtimulus that would have puſhed our reſearches a greater length, and we remain for ages in a pernicious errour, which the hope of reward, and the aſpiring to fame, would have reſcued us from. The hiſtory of human diſcoveries is filled with examples of this. We owe every thing to theſe two great ſprings of human actions, intereſt and ambition. But when a perſuaſion of knowledge

is

is entertained, all inveſtigation is laid aſide; and far from any further diſcoveries being made, the very hopes are loſt of knowing any thing more. Such was the deſtiny of Europe when in a ſtate of barbariſm and ignorance; and ſuch are ſtill the ideas of the ſavage.

It is now more than ten years ago that I publiſhed a work *On the Venom of the Viper*, in Italian. It is this work which forms the firſt part of the preſent treatiſe. I then in a manner engaged myſelf to the publick to give a ſecond part, in which I propoſed not only to ſpeak of the remedies againſt this venom, but likewiſe to treat of ſeveral other intereſting points altogether new. I had neither time nor convenience to finiſh all the reſearches I had then in view. I was deſirous of certain and evident conſequences, and it was neceſſary to multiply experiments infinitely, and to vary them a thouſand ways. But the little ſucceſs I met with in ſeeking a ſure remedy againſt the bite of the viper, particularly occaſioned me to defer the publiſhing the Second Part for ſo long a time. Not but that I tried a great number already known, and ſeveral others which either imagination or chance threw in my way; but they all appeared to me more or leſs inefficacious, and I found none of them certain. It is natural to ſuppoſe, that, amongſt other remedies, I did not forget that moſt famous one of all, *Eau de Luce*; (which indeed is nothing more than the *fluid volatile alkali*, joined with a little oil of amber, that does not at all alter its qualities) I tried it, but the ſucceſs

ceſs by no means anſwered my expectations, ſo that I at length abandoned it, as I had done all the reſt.

A new work has lately awakened the attention of the publick on the advantages of *the volatile alkali* againſt the venom of the viper *(a)*. It is announced in this work, in a tone of aſſurance and perſuaſion, to be *the true ſpecifick* againſt this dangerous poiſon, as well as againſt almoſt all diſeaſes, however dreadful. In reading this work, I apprehended that I had been miſtaken from beginning to end. Indeed when I reflected on the experiments I had made in Italy, I knew no longer what to believe, and was at times even led to think, that the vipers of France are either leſs venomous and deſtructive than thoſe of Italy, or that they are of a different kind; ſo true it is, that ſelf-love does not allow us to acknowledge our errours till the laſt moment!

But what ſurpriſed me ſtill more, was to ſee the errours of Redi, on the uſe of the bag that covers the canine teeth of the viper, which were refuted more than thirty years ago by Mead, again make their appearance in modern publications; to ſee likewiſe the errours of Mead on the acidity of the venom of the viper, which he himſelf had abjured; and, laſtly, to find thoſe of the ſame authour on the

(a) This work is entitled; *Expériences propres a faire connoitre que l'Alkali volatil fluor eſt le remede le plus efficace contre les aſphyxies. Paris.* The Authour is Mr. Sage of the Academy of Sciences.

ſaline

ſaline nature of this venom, which have been refuted in Italy for more than ten years *(a)*.

If on one hand I could not perſuade myſelf that I had been deceived on ſo many points and queſtions, which I had, however, examined without prejudice, and with a wiſh to diſcriminate nicely; on the other hand, it was impoſſible for me to conceive, how certain authours could advance ſo many matters as facts, with ſo much aſſurance, and without having previouſly convinced themſelves by certain and repeated experiments. I could not comprehend any more, why the authours of theſe new publications had neglected to demonſtrate clearly the ſource of the errours into which the writers poſterior to Mead had fallen, who had flattered themſelves that they had ſhown with the ſtrongeſt evidence, both by obſervation and certain experiments, how Redi and Mead had each of them been led into errour.

As the publick are perſuaded, that in phyſicks, ſubjects are ſubmitted to experiment, and not to authority, theſe gentlemen ought to have expoſed

(a) There is however, nothing aſtoniſhing in this, when we conſider the mode that is generally adopted by our modern writers. More than two hundred authours may be named who have copied from each other on this ſubject, and have given us groſs errours for demonſtrated facts. One might reaſonably exclaim to them, "Modern parrots, the copyiſts of other parrots, ceaſe to deceive us, and for once conſult nature. Had you employed the time you have ſpent in copying each other, in making experiments, how many errours, and how much time, would you have ſpared to poſterity!"

experiments

experiments to experiments, and obſervations to obſervations, and to have developed the errours into which we are fallen. But they have not made any ſuch attempt. They have ſubſtituted their authority to experiment, and their name to obſervation. This method is altogether pernicious, and neceſſarily tends to perpetuate errours amongſt men, and to render diſputes eternal. When we know that two obſervers do not agree on a fact, or on an experiment, to which of the two muſt we truſt, provided they are both of an equal merit? We remain in an abſolute uncertainty, and can only have acquired, on their peruſal, a reaſonable pyrrhoniſm.

But is there no touch-ſtone to enable us to judge where the miſtake lies betwixt them, and of two contradicting experiments, to diſtinguiſh the true one from the falſe?

The difficulty of judging betwixt two authors, even in matters of ſimple fact, has been the occaſion of many errours and hypotheſes having laſted a long time, even after their falſehood has been demonſtrated; and many truths have been rejected, merely becauſe experimenters have not been able to repeat the experiments that proved them, in the ſame way in which they were firſt made.

For my part, I think it the duty of the lateſt obſerver, not only to repeat faithfully the anteriour experiments that contradict his, but likewiſe to give his own in ſuch a way, that they cannot leave the ſhadow of a doubt in the mind of the reader. Without this proviſo, he will loſe the aim he propoſed

poſed to himſelf in writing—that of being believed; which he will not deſerve, although he may, by accident, have ſaid the truth.

There are three principal methods of avoiding this inconvenience, which perpetuates errours, and ſtill keeps us in a very dangerous ſcepticiſm.

The firſt is, to multiply the experiments exceedingly. It is almoſt impoſſible, in repeating them ſo many times, that fortuitous caſes do not occur to vary them, and that the final reſult of ſo many of them is not certain and conſtant.

The ſecond is, to vary them in a thouſand ways, changing the circumſtances as the nature and ſpecies of them may require, and giving them all the preciſion and ſimplicity they are capable of. This method ſuppoſes much greater talents and genius in the obſerver than the firſt, and there are few of theſe, even amongſt the moſt ſkilful, who can boaſt of having invariably put it in practice.

The third method is, not only to ſucceed in making experiments, deciſive by their number, variety, and ſimplicity; but likewiſe to attain to a diſcovery of the ſource of the errours that others have fallen into.

It is a fault, then, in thoſe who write the laſt, not to enter into a very minute detail of their experiments, and to endeavour to demonſtrate their ſuperiority and exactneſs, in compariſon with thoſe of their predeceſſors. It, however, is particularly incumbent on them to trace the origin of errours, and to ſhow how the former obſervers have been deceived.

ceived. Without this, all their labour is a pure loſs, and they are by no means worthy of confidence.

From all theſe conſiderations, I have deemed it neceſſary to return to the ſubject of the preſent work, and to treat it in as particular a way as my circumſtances will allow me. The importance of the ſubject requires this of me, ſince it regards a very dangerous and mortal diſeaſe, which impreſſes with fear thoſe who have the misfortune to be attacked by it, and creates uneaſineſs in families.

Perſuaded that a perfect knowledge of the venom of the viper cannot be acquired unleſs by a ſearch into all its properties, which are, in a greater or leſs degree, unknown, I wiſhed that neither of them ſhould eſcape me, without ſubmitting it to a rigorous, and, at the ſame time, impartial, inveſtigation; and that nothing which related to the ſubject ſhould be wanted, was deſirous of examining afreſh the ſuppoſed acidity of this venom, and the ſalts of which ſome people will have it to be compoſed.

Any errour whatever that relates to this ſubject, may, in time, become dangerous in its tendency. Authours, perſuaded by a miſtake of Mead, that they were acquainted with the true nature of the venom, have been ready to fabricate ſyſtems to explain the way in which it acts, and how and by what mechaniſm it is that it produces ſo ſpeedy a death. They have afterwards invented remedies that relate to the ſuppoſed nature of this poiſon,

and

and what is ſtill more ſtrange, have found them efficacious. They have ſhouted victory, both on occaſion of the theory and the remedy, and have ſhown how the one ſerved as a guide to the attainment of the other. In a word, they have pretended that all is done, and that nothing more remains to be known of the viper's venom; maintaining, that they are acquainted with its nature, its mode of action on the animal machine, and laſtly, with the remedies capable of deſtroying its effects.—But let us leave theſe authours with their ſectaries, to applaud themſelves on knowing ſo many things, and on having divined nature. I believe, on the contrary, that we as yet know nothing about it, and that this matter is altogether new. My experiments will ſhow this, in the courſe of the preſent work.

A great part of theſe experiments required the aſſiſtance of ſeveral perſons, and I have reaſon to congratulate myſelf on this neceſſity, ſince, amongſt others, it procured me the preſence of two men of rare talents; Dr. Troja, Member of the Royal Academy of Naples, author of ſeveral excellent tracts on animal phyſicks, who happened to be at Paris at the time I made my experiments (*a*); and M. Jean Fabroni, of Florence, a fellow traveller, and attached to the cabinet of natural hiſtory of the Grand Duke of Tuſcany, a well inſtructed, and very promiſing

(*a*) M. Troja viſited me almoſt every day, to obſerve my method of making experiments on various ſubjects in phyſicks.

young man *(a)*.—I name theſe gentlemen here with the greater pleaſure, ſince, in thus publickly teſtifying to them my gratitude and eſteem, I give my experiments a greater degree of authenticity.

The firſt queſtion I now undertake to examine, and which has been the principal occaſion of my enquiries, is, whether the *fluid volatile alkali* is a certain remedy againſt the bite of the viper; that is to ſay, whether it reſcues from death an animal that would otherwiſe have periſhed by it. This firſt reſearch is clearly very intereſting, and deſerves to be examined with all poſſible attention. I have multiplied my experiments on this firſt point, in a way that more than one of my readers will deem unneceſſary. But I know of what weight the prejudice for a favourite hypotheſis and the authority of a celebrated writer, are. Errour and truth ſeem to meet with the ſame difficulty and reſiſtance from mankind; one in unrooting, the other in eſtabliſhing, itſelf. The Newtonian ſyſtem was combatted for a whole age before it was received, and it required as long a time to abandon that of Deſcartes. What is very certain is, that ſo many errours have not been ſpread abroad, as to the nature of the venom of the viper and its remedies, but becauſe too few obſervations have been made, and experiments too little diverſified.

(a) M. Fabroni was likewiſe preſent at the experiments I made in London and on my return into Tuſcany, and willingly charged himſelf with the deſigns of the plates in this work.

Mead

Mead himſelf was not exempt from this fault, as I ſhall ſhow in examining the remedies he has propoſed againſt the bite of the viper. The uſe of the volatile alkali itſelf was only introduced in conſequence of a falſe theory on the nature of the venom, and was only ſupported with ſo much prejudice and obſtinacy, for want of the making of a ſufficient number of experiments. 'Tis on the ſame account that the diſputes on animal phyſicks, which would have terminated at their birth if experiments had been much more multiplied than they were, ſtill exiſt. But the art of experimenting is ſlow and painful, inſtead of which it coſts but little trouble to follow the authority of another. It is eaſier to reaſon than to make experiments; and this art, invariably long and difficult, is not within the reach of every one.

Other readers will find, that the number of my experiments, however great it may be in itſelf, is not ſufficient to decide all the queſtions I examine in this work, nor to terminate all the reſearches I make into the venom of the viper. I have nothing to oppoſe to theſe laſt, and, likewiſe, I do not take upon me to ſay, that all the conſequences I have deduced from my experiments are certain. Perhaps a number of experiments twice as great, would ſcarcely be ſufficient for this. Thoſe who are acquainted with the difficulties that are met with in experimenting on living animals, and who know how much the circumſtances betwixt one animal and another vary, (which rigorouſly ſpeak-

ing, are never the ſame) will agree with me on this head.

Let all that has been written on the irritability and ſenſibility of the animal fibres be examined, and the ſame inconveniences, the ſame difficulties, will be diſcovered. It is true, that a very great number of experiments have been made in a few years, and that an infinite number of animals have beeen ſacrificed to philoſophy, or publick utility; but much remains yet to be known, preciſely becauſe the number of experiments is not yet ſo conſiderable as it ought to be.

I muſt likewiſe confeſs, that I have wanted both time and patience to do more. Nothing but the idea of publick utility can ſupport the horrour of ſeeing ſo many animals, ſenſible of pain like ourſelves, ſuffer under our hands; and to view them expoſed to a thouſand kinds of torments. I leave the purſuit of this career to thoſe who are more courageous than myſelf. The road is open to obſervers, and I ſhall rejoice to ſee them embrace with ardour, the ſearch of truths that are advantageous to the human ſpecies.

CHAPTER II.

Whether the Volatile Alkali is a certain Remedy againſt the Bite of the Viper.

I DEEMED it neceſſary to examine this firſt queſtion in the moſt circumſtantial way, and therefore multiplied the experiments extremely, and diverſified them very much. This is the only method that could lead to demonſtration, and I flatter myſelf that my readers will be freed from all doubt.

The animals I had bit by vipers were of three different kinds. I employed birds and quadrupeds with warm blood; and frogs, which have the blood cold.

Amongſt birds, I almoſt always employed ſparrows, pigeons, and fowls; amongſt quadrupeds, rabbits, guinea-pigs, cats, and dogs.

An animal may be bit by a ſingle viper, and by ſeveral. It may be bit once or more; in a ſingle part, or in ſeveral.—All theſe caſes may make great variations in the diſeaſe and effects of the venom; it was therefore neceſſary to diſtinguiſh them from each other.

Animals bit by a ſingle Viper, and only once.

The leg is the part of the animal I conſtantly had bit by the viper, in all the experiments contained in this chapter. By *leg*, I mean the muſcular part of the foot, that is betwixt the femur and tarſus. The facility of having animals bit in this part by the viper, made me give it the preference. There is likewiſe another advantage, the eaſe with which the remedies are in this caſe applied.

In the experiments of this chapter, as well as thoſe of the following one, I employed no other remedy againſt the bite of the viper than the *fluid volatile alkali*, to be found in every apothecary's ſhop. Some that I made uſe of, I made myſelf. Its compoſition has been long known, and is deſcribed in all the pharmacopeias. I employed it by having it ſwallowed, and by applying it to the part. When I wiſhed to treat the part bitten, I dapped it a long time with a piece of linen, well moiſtened with the volatile alkali, and laſtly, covered it with the ſame linen, to keep it wet the longer. That which was ſwallowed, as will be ſeen hereafter, was diluted with a quantity of water. On many occaſions, I repeated it ſeveral times, and likewiſe made freſh applications of it to the part. There are animals which live ſo ſhort a time after they have been bit, that I thought it ſuperfluous to make repeated applications of the volatile

latile alkali to the bite. When I ſay, ſimply, that I treated the part bit, or that I treated the animal, it muſt be underſtood, that the volatile alkali was not given internally, but only applied to the part.

I had a dozen ſparrows bit by as many vipers, a ſingle time each. I took them from the cage, one after the other, without any choice. The firſt that was bit was immediately treated; the ſecond was not; the third was treated, and the fourth not; and ſo on as to the others, each having a thread tied to its foot, with knots, to diſtinguiſh them from each other. The feathers had been previouſly cut from the legs with ſciſſars. The animal was ſcarcely bit by the viper, when it was treated; ſo that there was an interval of not more than five or ſix ſeconds betwixt the bite and the application of the *volatile alkali*.

The ſparrow firſt bit, which was treated, at the end of two minutes could no longer ſupport itſelf on its feet, and died at the end of fifteen.

The ſecond, not treated, began to reel after three minutes, and died at the end of the thirty-fifth.

The third fell on its belly after ſix minutes, and died at the end of thirty-eight.

The fourth fell after four minutes, and died at the end of twenty.

The fifth after five minutes, and died at the end of twenty-ſeven.

The ſixth after ſeven minutes, and died at the end of thirty.

The ſeventh was ſtiil living at the end of three hours, and did not appear to have at all ſuffered.

The eighth fell after two minutes, and died at the end of ſeven.

The ninth fell after three minutes, and died at the end of eleven.

The tenth fell after two minutes, and died at the end of fifteen.

The eleventh fell after a minute and one-third, and died at the end of two and an half.

The twelfth fell after ſix minutes, and died at the end of thirty-two.

The ſparrow that was bit the ſeventh, as I have juſt ſaid, was ſtill living at the end of three hours. I examined its leg, and found it perfectly in its natural ſtate, without lividneſs, without ſwelling, or any apparent wound. The legs of the other ſparrows were very much changed, even immediately after they had been bit; whence it was eaſy to conjecture, that either the ſparrow above alluded to had not been bit by the viper, or that this laſt was without venom.

To diſcover which of theſe two hypotheſes was the true one, I had this ſparrow bit by the ſame viper, in the ſame leg. A little blood flowed from the wound, which I immediately treated. It fell after two minutes, and died at the end of four; a proof that the viper was provided with venom, but that the leg had not really been entered by the teeth. I had, however, no ſuſpicion of this at firſt, as the creature bit in the uſual way.

I wiſhed

I wiſhed to repeat the ſame experiment on twelve other ſparrows, with the ſame order and circumſtances. But I made the ſix that were treated likewiſe ſwallow a few drops of water, in which I had put a proportion of the volatile alkali, of about an hundredth part.

The time of the death of theſe animals is expreſſed by the following numbers, repreſenting as many minutes elapſed after the bite; 10, 7, 8, 9, 6, 7, 3, 7, 15, 18, 5, 37. The ſix firſt numbers ſhow the time the ſparrows lived, that were treated with the volatile alkali.

From the preceding experiments the following conſequences may be deduced:

I. That the vipers I employed were ſufficiently provided with venom to kill ſparrows.

II. That the venom is ſcarcely introduced into the leg of the animal, when it ſwells in a ſenſible degree, changes its colour, and becomes ſomewhat livid.

III. That it is not ſufficient to enable the venom to inſinuate itſelf, that the viper ſeizes an animal betwixt its teeth and that it cloſes its mouth, and preſſes with it.

IV. That the fluid volatile alkali does not preſerve the lives of the ſparrows bit by the viper.

V. That the volatile alkali given internally to ſparrows, may even be hurtful to them. The ſpeedier death of thoſe that ſwallowed it, may at leaſt lead one to ſuſpect ſo.

But the number of the experiments is not yet ſufficient to render the conſequences I have juſt deduced certain; 'tis a multiplicity of them alone that can effect this.

I had twelve other ſparrows, equally lively, bit in the leg as above, each by a ſingle viper, and only once. I treated only ſix of them with the volatile alkali. They all died. In all of them the leg that was bit became livid, and ſwelled in a greater or leſs degree, in leſs than two minutes.

The ſix treated died in 3, 4, 6, 11, 30, 33, minutes. The ſix that were not treated in 4, 4, 7, 11, 18, 35.

To obtain ſtill more certain conſequences, I had twenty-four others bit. I treated twelve, and made them ſwallow the volatile alkali. All the twenty-four died. The following numbers ſhow the minutes the twelve that were treated lived, 2, 3, 3, 5, 5, 5, 7, 7, 10, 15, 15, 22; and theſe again indicate the minutes that thoſe ſurvived on which no remedies were tried, 4, 6, 6, 6, 7, 7, 9, 9, 9, 10, 15, 20.

It is a truth then, eſtabliſhed by experiments, that the fluid volatile alkali is altogether uſeleſs, whether it is ſimply applied to the part bit by the viper, or ſwallowed by the animal at the ſame time. We may even ſuſpect it to be hurtful, to ſparrows at leaſt.

However evident it may appear, that the volatile alkali is not an efficacious remedy in this caſe to a ſmall animal like a ſparrow, it is not on that account demonſtrated,

demonſtrated, that it may not be ſo to a much larger animal, and of a different ſpecies.

The venom introduced into the body of a larger animal, ſhould be conſidered as diminiſhed in quantity. Its effects ſhould certainly not be ſo violent; and indeed this is the caſe with all the poiſons that we know of. What is a remedy to a large animal, or one of a full ſize, may be a poiſon to a ſmaller animal, or to one ſtill young.

We muſt therefore again have recourſe to experiment, and ſee the effect the bite of the viper has on other animals.

Experiments on Pigeons.

I had a pigeon bit in the leg by a viper, and inſtantly treated the part. At the end of a minute it fell forward, and could no longer ſupport itſelf. In twenty ſeconds more it died.

I had another pigeon like the firſt bit in the ſame way, but did not treat it. At the end of two minutes it fell forward, and in two minutes more it died.

I had two other pigeons bit in the leg; one was treated, and the other not. The firſt fell at the end of three minutes, and died at the end of the twentieth. The other fell at the end of a ſingle minute, and died likewiſe after the twentieth.

Of two other pigeons bit in the leg, I treated only one. The one treated died at the end of forty hours, the other at the end of an hour.

I had

I had ſix other pigeons bit in the uſual way. Three were treated, and three not. Thoſe that were treated died at the end of 6, 22, 40, hours. The other three died at the end of 1, 2, 10, hours.

I had two others bit in the leg in the uſual way; one I treated, the other I did not. The treated one died at the end of eight minutes; the other at the end of two hours.

The intervals at which pigeons die that are bit by the viper are ſo different, that they ſcarcely allow a reaſonable conjecture. It ſeems, however, that two truths may already be deduced. One, that the *volatile alkali* does not preſerve from death the pigeons bit by the viper. The other, that birds larger than ſparrows live longer in the ſame circumſtances; or, if you will, that pigeons die much later than ſparrows.

But experiments muſt be multiplied, and the circumſtances attending them examined more attentively.

I do not conceive very well how it was, that of two animals of the ſame kind, bit once in the ſame part, one died at the end of two minutes, and the other at the end of 40 hours.

I likewiſe obſerved ſomething ſimilar to this in the ſparrows; and therefore determined at length to have a very large number of both kinds bit. I did not treat any of them; but, in return, I marked carefully all the circumſtances that attended the experiments. I ſhall not enter into a detail of them here, on account of the very great number of them;

but

but think it ſufficient to deduce the following truths:

I. That other circumſtances alike, the larger the viper, the more violent the diſeaſe, and the more ſpeedy the death.

II. That the diſeaſe increaſes likewiſe in violence, in proportion as the viper is more enraged.

III. That it likewiſe augments in proportion to the time the viper compreſſes the animal it has bit betwixt its teeth.

IV. That the diſeaſe of the part bitten ſeems to be greater in proportion to the time the animal ſurvives.

V. That in ſome animals black and livid blood flows from the wound, as ſoon as it is made.

VI. That in others, on the contrary, it flows red, and preſerves that colour.

VII. That the animals from which the red blood flows, die later than thoſe from which it flows black and livid.

VIII. That the venom likewiſe, which preſerves its colour and its qualities, ſometimes flows out with the blood. In which caſe, the animal not only ſurvives, or is much longer in dying, but ſometimes does not appear to have had any complaint.

Theſe conſequences, the fruit of an infinite number of experiments, diverſified in every poſſible way, and in which all the circumſtances that accompanied them were rigorouſly examined, form ſo many principles, which explain how it is that of two animals

mals bit in the ſame part, one dies ſuddenly, and the other ſurvives, or does not die till very late.

There is likewiſe another reaſon, which I have ſince diſcovered, and which may vary the effects, in animals that have been bit, very much. This is owing to the viper itſelf. I have ſometimes, tho' very rarely, found vipers that had no venom in either of the two veſicles, and more frequently, that only had it in one.

What led me at firſt to ſuſpect that the veſicles did not conſtantly contain venom, was obſerving it to be to no purpoſe that I had a pigeon bit repeatedly by a certain viper; and that it not only did not die, but diſcovered no ſymptom of diſeaſe, notwithſtanding the canine teeth of the viper had pierced its fleſh through in ſeveral places.

Having had occaſion, in the courſe of theſe experiments, to cut of the heads of a great number of vipers, and to examine their venom, out of two hundred, perhaps, I found two that were entirely deſtitute of venom, and five that, inſtead of venom, had a white and opake viſcous matter in the veſicles. In two of theſe laſt, I found this white matter to be perfectly innocent. But in the other three it ſtill preſerved, partly at leaſt, its venomous quality, as I aſſured myſelf by introducing a ſmall quantity of it into the legs of pigeons, which had been bit ſuperficially, and which died at the end of a few minutes.

It is another eſtabliſhed truth then, that vipers are ſometimes found without any venom, and that

ſomewhat

ſomewhat more frequently a whitiſh humour is contained in their veſicles, which is not always venomous. But theſe caſes are invariably very rare, and only met with in examining a very great number of vipers; whence it follows that it is alſo true, that vipers have in general their veſicles filled with venom, and that this humour occaſions diſeaſes, and even death.

I obtained much more uniform conſequences, by introducing the venom into the body of the animal, inſtead of having it bit by the viper. This is the method I employed. I cut off the head of a viper with a pair of ſciſſars, and, after a quarter of an hour, opened the mouth, and with another pair of ſciſſars ſeparated the lower jaw. I then divided the upper part of the head in two with very ſtrong ſciſſars; each part being furniſhed with the canine teeth, and with the veſicle of venom. With a little courage and dexterity, which are acquired by cuſtom, it is eaſy to force the tooth of the viper, on which a compreſſion is made with the fore finger whilſt the veſicle is preſſed upon by the thumb, into the ſkin of an animal. A greater or leſs quantity of the venom may be introduced, by preſſing more or leſs on the veſicle; the wound may be made wherever one pleaſes; and, laſtly, the venom may be kept from being rejected, by letting the tooth remain a long time in the wound. A great number of experiments made in this way, prove that ſparrows die betwixt five and eight minutes, and pigeons in betwixt eight and twelve. There are very few that

die

die ſooner or later; whence it follows, that by purſuing this method, the periods of their diſeaſe are both ſhorter and more uniform.

I had a dozen pigeons bit in the uſual way, one after another, by as many vipers, and treated them all with the volatile alkali. They all died. The numbers 4, 10, 16, 52, expreſs the time in minutes in which four of theſe pigeons died; and the numbers 2, 4, 9, 15, 19, 22, 25, 36, expreſs the time, in hours, of the death of the others.

Theſe new experiments leave no doubt as to the inefficacy of the fluid volatile alkali againſt the venom of the viper.

To aſſure myſelf ſtill better of this, I had twenty-four other pigeons bit, each of them once in the leg, by a viper. I treated them all, but only twenty-two died. The time of their death is expreſſed in minutes, by the numbers 4, 4, 6, 6, 7, 8, 8, 10, 12, 14, 14, 20, 50, 50, 56; and in hours, by 1, 1, 2, 4, 7, 10, 18, 26, 30.

Two of theſe pigeons, bit in the ſame way as the others, appeared not to have ſuffered at all, walking about the chamber as before the operation. At the end of two hours, being deſirous of examining the ſtate of their legs, I could find no ſign of diſeaſe. They were neither ſwelled nor livid. I could only find in one of them a ſmall hole, and a ſmall red ſpot of blood, at the part where the tooth had penetrated. Since there was not the ſmalleſt mark of diſeaſe, it was eaſy to perceive that the venom had not introduced itſelf; or, if it had, that it had been

thrown out again, ſo as not to occaſion any complaint to the animal. After ten other hours, I had both pigeons bit once in the ſame leg by two vipers that had already been employed in the ſame way. At the end of three minutes there were ſigns of diſeaſe: one died at the end of an hour, the other at the end of two.

Not content with theſe experiments, I had twelve other pigeons bit in the uſual way. I treated them immediately, and made them ſwallow the volatile alkali. They all twelve died, at the end of 4, 4, 7, 10, 10, 10, 15, 18, 20, minutes; and of 2, 3, 3, hours.

Whilſt it is certain, on one hand, that the volatile alkali is of no effect in recovering pigeons bit by the viper; on the other hand it remains undecided whether it is in this caſe hurtful or not.

The periods at which theſe animals die are ſo various, that it is not poſſible to deduce any certain conſequences from them.

Experiments on Fowls.

It is not ſufficient to have demonſtrated the inutility of the fluid volatile alkali adminiſtered to pigeons, to allow us to conclude that it is uſeleſs to larger animals, that are more difficult to kill. The volatile alkali may have time to act againſt the venom of the viper, when the diſeaſe is not ſo violent, and the animal ſlower in dying.

There are certain remedies which, although efficacious,

cacious, require a certain time to act; and, indeed, almoſt all are of this deſcription.

I had a fowl bit once in the leg by a viper, and immediately treated it; at the end of ſix hours the fowl died. I afterwards had another bit once by a viper, and did not treat it. This one died in eight hours.

I had two other fowls bit once in the leg as uſual. One was treated; the other not. The firſt died in four hours; the other in ten.

I had ſix other fowls bit as above, each once in the leg by a diſtinct viper. The three firſt were treated with the volatile alkali, and died; one in ſix hours, another in eight, and the third in nine. The three others were not treated, and died in 7, 9, 20, hours.

Although the number of the experiments hitherto made on fowls, is not yet ſufficient to allow certain conſequences to be drawn, it however appears, that the following very probable ones may already be ſtated.

I. That it is very poſſible for fowls bit once in the leg by a viper, to die.

II. That they in general die much later than pigeons; and than ſparrows, which die again with much greater facility than pigeons.

III. That birds reſiſt death in proportion to their ſize.

IV. That the volatile alkali is not only of no uſe in curing fowls bit by the viper, but that it is probably even hurtful to them.

But

But experiments muſt be multiplied much more, to ſee if the conſequences juſt deduced are well or badly founded.

I had therefore ſix fowls bit ſeparately by ſix vipers, each once in the leg. I treated them all ſix, and made a freſh application of the volatile alkali to the part bitten, every two hours. Two of the fowls died in the ſpace of four hours, one in five, two in ſix, and one in ten. A moment after, I had ſix other fowls bit by as many vipers, each once in the leg, and did not treat either of them. Two died in two hours, two in ten, and two in twelve.

Twelve other fowls were bit by as many vipers, each once in the leg. I treated ſix, and made them ſwallow the volatile alkali. The other ſix were left to themſelves. Of the ſix treated, five died; the ſixth had ſcarcely any ſymptom of complaint. Its leg neither ſwelled, nor became at all livid. There was ſimply a hole in the ſkin, which was red and a good deal inflamed. The five I have juſt mentioned died in 3, 4, 6, 7, 10, hours. The other ſix died in 6, 10, 17, 22, 36, 36, hours.

Had the experiments I have related ſo far been more numerous, the abſolute inutility of the fluid volatile alkali againſt the bite of the viper would not only have been demonſtrated, but we might even have doubted its innocence, at leaſt to this ſpecies of animals.

The treated fowl that did not die, proves nothing in favour of the volatile alkali, as will be ſeen in the continuation of this work. It is one of the

caſes I remarked above, in ſpeaking of the pigeons and ſparrows, in which the venom was not communicated to the part bit, although the canine tooth had left ſome opening in it, either owing to the viper not having any venom, or to the rejection of it. Nothing is found in either of theſe caſes to favour the volatile alkali.

Having aſſured myſelf of the inutility of this remedy to the three ſpecies of birds I have ſubmitted to the experiment, I think it time to make the ſame trials on quadrupeds.

Experiments on Guineapigs.

I had a large guineapig bit once in the leg by a viper, and immediately treated it. In a little time the leg ſwelled, and became livid. At the end of ſixteen hours a wound of an inch in breadth formed itſelf at the part that had been bit and treated. In twenty hours the ſkin in this part was entirely eaten away. The wound continued open for more than twenty days, during which time the animal moved its leg with difficulty; the foot was greatly contracted, and the muſcles very much diſeaſed. The animal recovered however, but its leg ſtill remained in a degree contracted, and it could never recover the perfect uſe of it.

Another guineapig, almoſt as large as the former, was, in the ſame way, bit once by a viper in the leg,

leg, which was not treated. The animal died at the end of two hours.

I had four others, of ſcarcely a third the ſize of two preceding ones, bit in the above way. I treated each of them, and made them ſwallow the volatile alkali. They all died, one in two hours, another in three, the third in ſix, and the fourth not till the twentieth hour had elapſed.

That I might have a comparative experiment, I had four other guineapigs, entirely like the preceding ones, bit, and did not treat either of them. They all four died, one at the end of ſeven hours, another at the end of ten, the third at the end of thirty, and the fourth at the end of thirty-one.

From theſe experiments we may, I think, already draw the following inferences, which if not certain, are at leaſt very probable.

I. That the bite of the viper is capable of killing even the largeſt guineapigs.

II. That the ſmaller animals die ſooner than the larger ones of the ſame ſpecies.

III. That the volatile alkali is not a certain remedy againſt the bite of the viper.

It may be objected, that the firſt guineapig bit and treated, at length recovered, and that all which were not treated died. This is true, but proves nothing, ſince, as has been ſeen above, there are ſeveral circumſtances that may render the bite of the viper innocent; and, on the other hand, we have ſeen that the other four guineapigs died, although they were treated. Now if we conſider

that the four treated died in a much ſmaller ſpace of time than the five that were not treated, we may ſuſpect that the volatile alkali was more than uſeleſs, that it was hurtful.

To remove all doubt, I had twelve other guineapigs bit, all alike in ſize, and ſimilar to the eight preceding ones. Six were treated, and ſix not.

The firſt I had bit was the ſame I have ſpoken of a little above, and which, far from dying of the bite, had not even the diſeaſe of the venom. Although treated, it died at the end of thirty hours. The five others that were likewiſe treated, had all of them the diſeaſe of the venom, but only three died; two in leſs than twenty hours, the other at the end of twenty-ſeven. The two that ſurvived had each of them a large wound in the leg that had been bit, and this remained open for more than ten days.

Of the ſix that were not treated two only died, in leſs than ſixteen hours. Three others had deep wounds, which remained open for ſeven days, and then healed. The ſixth had not the ſmalleſt ſymptom of diſeaſe, and I could not diſcover in its leg any mark of the viper's tooth having penetrated.

The caſes ſo far related, ſeem to leave no doubt as to the inutility of the volatile alkali, when tried likewiſe on theſe animals; and they do not remove the ſuſpicion, that it may poſſibly be even hurtful to them.

We likewiſe ſee that the ſmaller and younger guineapigs die ſooner than the larger ones.

I had a dozen very ſmall ones bit, each ſcarcely weighing five ounces. Six were treated, and ſix not. Thoſe that were treated died in 30, 40, 50, minutes, and 1, 2, 3, hours. Thoſe that were not treated in 57 minutes, and 2, 3, 4, 4, 4, hours.

I afterwards had ſix guineapigs bit, three of the largeſt of which were treated; the other three were not. Of thoſe that were treated only one died, and only one again of thoſe that were not treated. All of them, however, were very much diſeaſed, and thoſe that were treated were the laſt to recover.

Experiments on Rabbits.

It remained for me to make the ſame experiments on rabbits, in purſuance of the plan I had propoſed to myſelf.

With this view, I had a large rabbit bit by a viper once in the leg, which I immediately treated with the volatile alkali, making the animal ſwallow the ſame diluted with water. At the end of an hour I repeated the application and the internal remedy. The rabbit died at the end of three hours, with very ſlight marks of diſeaſe in its leg.

I had another, perfectly like the former, bit in the ſame way once in the leg by a viper, and at the ſame time. It had ſlight ſymptoms of the diſeaſe of the venom, and the leg became ſomewhat ſwelled. At the end of thirty hours a wound two lines in breadth, and very deep, appeared on the ſkin at

the part where it had been bit. After five days more the animal was perfectly recovered.

The reſult of two experiments alone can be in no way certain, I therefore had recourſe to my uſual method.

I had a dozen rabbits of a middle ſize bit by as many vipers, each once in the leg. Six were treated, and ſix not. Only two died of thoſe that were treated, and three of thoſe that were not. Two of the four treated ones that did not die ſcarcely had any complaint. Their legs were but little ſwelled, and were not livid. The other two were very much diſeaſed, and had large wounds that were four days in healing. Of the two that died, one lived two hours, the other five. The ſix that were not treated were all of them very much diſeaſed, and had large wounds in their legs, which ſwelled violently, and became very livid. The three that died lived 14, 22, 47, hours; the others did not recover till the end of the ſeventh day.

It is a conſtant obſervation, that when the animal bit by the viper dies very ſoon, the bitten part is proportionably leſs changed, leſs ſwelled, and leſs livid. The change which takes place at the part where the poiſon has entered, I call the external diſeaſe, to diſtinguiſh it from the other, which is infinitely more violent and dangerous, and which kills the animal in a more direct way. I ſhall ſpeak more fully of this laſt in the fourth chapter of this ſecond part, in which I ſhall endeavour to account for this particular.

The

The few experiments hitherto made on rabbits may already make us ſuſpect the little efficacy of the volatile alkali, which we may be even tempted to believe hurtful. It is certain in the interim, that middle-ſized rabbits frequently reſiſt the venom of the viper.

I wiſhed to try the effects of this on much ſmaller ones, and for this purpoſe had a dozen bit in the uſual way. I treated ſix, and did not treat the others. All the twelve died; the treated at the end of 2, 3, 4, 6, 8, 9, hours; and the others at the end of 3, 5, 7, 9, 12, 13.

I repeated theſe experiments on twelve other ſmall rabbits, exactly like the foregoing ones. I treated ſix, and made them ſwallow the volatile alkali every hour. The others I did not treat. They all died; thoſe that were not treated, at the end of 1, 1, 2, 2, 5, 17, hours; the others in the ſpace of 1, 3, 3, 10, 16, 16, hours.

Theſe new experiments already ſhow very clearly the little efficacy of the volatile alkali againſt the bite of the viper, when tried on rabbits; they even lead me to ſuſpect it to be rather hurtful than otherwiſe.

We likewiſe ſee that ſmall rabbits die from the bite of the viper, whether they are treated or not; but that the larger ones frequently ſurvive its effects.

In conſequence of this, I had ſix of theſe animals, very large ones, bit each by a viper once in the leg. Three were treated, and ſwallowed the vola-

tile alkali. Two of theſe died at the end of twenty hours; the third was very much diſeaſed, and had a large wound, which remained open for twenty-three days. Of thoſe that were not treated, one died at the end of thirty-four hours; the other two had the diſeaſe of the venom, but recovered in leſs than ten days.

I repeated this experiment in the ſame way on ſix other large rabbits. Of the three that were treated, one died; and one likewiſe died of the three that were not treated. The other two of theſe laſt recovered in ten days; and the two treated ones that ſurvived, not till the end of eighteen.

I think there can be no longer any doubt of the inefficacy of the volatile alkali to theſe animals; on the other hand, inſtead of diminiſhing it, it ſeems to ſtrengthen and reinforce the diſeaſe.

It remains to try the effects of the bite of the viper on cats and dogs. The number of my experiments on the animals of theſe two ſpecies is much ſmaller than that of thoſe on the others. The difficulty of procuring them, the danger one runs in operating on them, and, ſtill more, the inconvenience of keeping them during the long continuance of the diſeaſe, and the unpleaſantneſs of ſeeing them ſuffer, have occaſioned me to do leſs in this inſtance than the ſubject may perhaps appear to have required.

Experiments on Cats.

I had two very ſmall kittens bit in the uſual way, each once in the leg. One was treated, the other not. The laſt died at the end of ſixteen hours. The treated one was exceedingly ill, and had a wound which remained open for five days, in its foot. It lived, however.

Three very ſmall kittens were brought to me, ſtill younger than the foregoing ones. I had them bit, as uſual, in the leg. I treated one, and made it ſwallow the volatile alkali. I did nothing to either of the other two. They all three died in leſs than ſix hours.

Theſe experiments are neither ſufficiently uniform, nor enough in number, to admit the drawing of certain conſequences from them. We ſee that, in general, the ſmaller animals of the ſame ſpecies, ſuch as cats, for inſtance, die much readier than the larger ones; and likewiſe that thoſe die which have been treated, and have ſwallowed the volatile alkali.

I had two other kittens bit, larger than thoſe I employed before. Each of them was, as uſual, bit once by a viper. One was treated, the other was not. Neither of them either died, nor was very ill. They had no wound; and both of them at the end of twenty-four hours ate very heartily. The leg, however, in each, was not yet very ſupple in its motions. I did not make the one I treated ſwallow the volatile alkali, on account of the difficulty

one

one meets with in attempting this, when theſe creatures are pretty large. They become extremely furious, and are very difficult to manage, at leaſt without riſk.

I had two other kittens of the ſame ſize of the preceding ones bit, and treated neither of them. They were each of them bit once in the leg. They both recovered, and had no perceptible wound. It was twenty hours indeed before they had any uſe of the leg that had been bit; however, they ſeemed perfectly recovered at the end of the third day.

Two large grown cats were, in the ſame way, bit in the leg. Neither of them was treated, and neither died. At the end of ſixteen hours they fed a little, and could already uſe their legs, although not very well. At the end of thirty hours they appeared to be perfectly recovered.

Scarcely has a cat been bit in the leg by a viper, when it can no longer make any uſe of the part. It lies down, and continues longer in this poſture in proportion to the violence of the diſeaſe. It neither eats nor drinks till the ſymptoms abate, and when that happens, recovers to a certainty.

Experiments on Dogs.

We are now to try the effects of the volatile alkali, which has been of no utility to the cats, on dogs that have been bit by the viper. The dog has a great affinity to man himſelf, and is, of all animals, the one the moſt ſuſceptible of the paſſions.

It

It is certainly much more ſo than the cat and the other animals, that have been bit in the courſe of theſe experiments. Dogs are to be met with of every ſize, even ſo large as not to differ much, in that reſpect, from an adult perſon.

The effects of the bite of the viper on dogs, may be of great uſe in judging of the bite of the viper in man himſelf.

I had two dogs of a middle ſize bit once in the leg. I treated one of them every two hours, and made it ſwallow the volatile alkali as often. Neither of them died, although the leg was ſwelled in each. The one not treated had no wound, and recovered at the end of four days; the one that was treated had a large wound, and did not recover till the cloſe of the tenth day.

I had two other much ſmaller dogs bit, and treated only one of them. They both died in leſs than three hours, with a degree of ſwelling and lividity in the part bitten.

Two large dogs were brought to me, and I conceived from their ſize, that they would recover although not treated. I had them bit in the uſual way, once in the leg. One ſcarcely had any ſenſible complaint; the other no perceptible wound. The leg of the laſt, however, ſwelled very much, and did not get well till the end of the ſixth day.

I had two other large dogs bit by a viper as uſual, each once in the leg, and did not treat them. One recovered in two days, the other in ſix.

From

From the experiments hitherto made on dogs, we may draw theſe concluſions:

I. That the ſmaller ones uſually die from the bite of the viper.

II. That large ones generally recover.

III. That of the middle ſized ones, ſome recover, and ſome die.

IV. That the volatile alkali ſeems to be neither a certain nor a uſeful remedy againſt the bite of the viper.

Experiments on Frogs.

It remained for me to try the effects of the venom of the viper on frogs. I had hitherto operated on animals with warm blood; it was likewiſe neceſſary to make ſome experiments on thoſe that have the blood cold.

I had a dozen frogs bit by as many vipers, each once in the leg. I treated ſix of them only. Two of theſe died at the end of twenty hours, and the legs of the other four ſwelled, and were a little livid; they recovered however. Of the ſix not treated, three died at the end of five hours. Of the three that ſurvived, one had a ſwelling and diſcolouration of its leg; the other two had no apparent complaint.

The conſequences were as yet too vague, and too few in number, to admit any certain concluſions to be drawn from them.

I there-

I therefore had a dozen others bit in the ſame way, and treated ſix of them only. To theſe I renewed the application of the volatile alkali every hour, making them ſwallow it at the ſame time. All the ſix, one of which did not ſurvive the twentieth minute, died in leſs than four hours. Of the ſix not treated, four died at the end of 6, 10, 12, 20, hours; the fifth had ſcarcely any complaint, and the ſixth recovered two days after.

I repeated this experiment on twelve other frogs, having them bit in the ſame way, each once in the leg by a viper. Six were treated every hour, and ſwallowed the volatile alkali as often. The other ſix were left to themſelves. Five of the firſt died; the ſixth had ſcarcely any ſymptom of complaint. Of the ſix not treated, three died, and the other three recovered at the end of two days.

After what has been ſaid, I think there can no longer be any doubt of the inutility of the fluid volatile alkali. It is very probable that, when given internally to frogs, it increaſes the diſeaſe cauſed by the venom, inſtead of diminiſhing it. It is at leaſt certain, that the animal dies the readier under theſe circumſtances.

CHAPTER III.

Of the Effects of the Bite of one or ſeveral Vipers, on the ſame Part of an Animal, or on two correſponding Parts of the ſame Animal.

I HAVE hitherto ſpoken of the effects of the venom on animals bit once by a viper, in the ſame part. It now remains to ſpeak of animals bit repeatedly by one or more vipers in different parts.

It is natural to conceive, that a viper which bites the ſame animal ſeveral times, muſt bring on a diſeaſe proportionably violent. After having ſeen in the firſt part of this work, that the venom of the viper is a humour ſeparated from the fluids of the animal, and ſecreted in a veſicle or gland; and that this humour is always venomous in itſelf when it is introduced by a wound into the bodies of animals, particularly of thoſe with warm blood; there can no longer be any doubt of this truth, nor of the abſolute falſehood of the hypotheſis of Monſ. Charas, who pretends that the venom of the viper is entirely occaſioned by the fury of the animal, which changes the ſaliva and other humours of its mouth, to ſuch a degree as to produce a powerful venom, ſuch as is obſerved in the foam of a mad dog.

The

The veſicle is moreover conſtructed in ſuch a way, that all the venom cannot flow out at once, at a ſingle bite, however forcible it may be, and however the viper may be enraged. A deſcription of this veſicle, with that of the gland, will be ſeen in the third part of this work. From the foregoing conſideration it was neceſſary to examine the effects and diſeaſes produced by ſeveral bites, although of a ſingle viper. There are ſeveral examples of perſons bit more than once by the ſame viper; and notwithſtanding this caſe is not one of the moſt frequent, it occurs however from time to time.

It is not only very important to examine the effects of the repeated bites of the ſame viper on the ſame part of an animal; but likewiſe to obſerve the action of the venom on the different parts of the ſame animal.

We know that an animal is formed of organs and parts, differently organized. There are parts that have veſſels and nerves, without having muſcles; and theſe are in different proportions, and differently diſtributed: there are others again that have no nerves, and, if they have any, have only a few very fine capillary veſſels. It is natural to ſuppoſe that the effects of the venom of the viper, on parts of an animal ſo very different from each other, muſt be altogether different; and that the ſame quantity of venom conveyed into a wound made in an animal, may produce either death, a ſlight diſeaſe, or none at all. In a word, it appeared to me, that nothing ought to be omitted in ſo important a matter.

There

There is likewiſe a caſe, although I think it a very rare one, in which ſeveral vipers together bite the ſame part, or different parts, of an animal. However rare this accident may be, it is not impoſſible for it to happen; and it is not an extraordinary thing to find, at certain times of the year, ſeveral vipers collected together. A man who may not have noticed this, may by treading upon them be in danger of being bit by more than one; and I knew a viper-catcher who was bit in the hand by two at the ſame time, and who might have been bit by more than two, ſince ſeveral of them were making their way at the ſame time out of a box.

Theſe examples of animals bit by ſeveral vipers may, however, agree very well, making ſome little allowance, with the caſes of the repeated bites of the ſame viper, whether on the ſame part, or on different parts, of an animal.

I ſaid above, that I had found by experience the effects of the venom to be much more uniform, when, inſtead of having the animals bit by vipers, the venom is conveyed into them, by preſſing with one finger the veſicle which contains it, whilſt with the other the tooth of the viper is forced into the part. I have frequently employed this method during the courſe of my experiments, particularly in thoſe on the ſparrows and pigeons. In this way I not only ſucceeded in wounding the ſame part of the animal over again to a certainty, but even the very fibre. I could likewiſe aſſure myſelf, if I was deſirous of it, whether the veſicle contained venom, or whether

whether the quality of the latter was ſuſpicious or changed.

The ſlighteſt compreſſion made on the veſicle is ſufficient to bring a very ſmall drop of venom to the point of the tooth; its tranſparent colour determines its activity and nature.

The firſt thing I thought it neceſſary to determine here, was to ſee whether the ſecond bite of the viper is as powerful as the firſt, the third as the ſecond, and ſo on as to the others; and how many times, one after the other, the viper can venom animals with its bite. I took a viper of a middle ſize, and very lively, and, without provoking it much, made it bite a pigeon once in the leg. The pigeon died at the end of twelve minutes. A moment after it had bit this one, I made it bite a ſecond, a third, a fourth, a fifth, a ſixth, and a ſeventh, in the ſame part. The ſecond died at the end of eighteen minutes, the third of ſixteen, the fourth of fifty-two, and the fifth at the end of twenty hours; the ſixth had ſcarcely any ſymptoms of complaint, and the ſeventh continued perfectly well.

I repeated this experiment ſeveral times, and the conſequences were ſomewhat various. I met with ſome vipers, particularly the largeſt of them, that could kill ten, and even twelve pigeons. If they are very much enraged during the firſt bites, the laſt, as I have aſſured myſelf by repeated experiments, are leſs dangerous.

It is an eſtabliſhed truth then, as I have ſeveral times experienced, that the firſt repeated bites of a

viper are almoſt equally dangerous; and that in proportion as a viper is enraged, the diſeaſe occaſioned by its bite is more violent.

This laſt truth may tend in ſome degree, to account for the treacherous experiments of Charas on the venom of the viper. In oppoſition to Redi, as has been ſeen above, he was of opinion that this venom conſiſts ſimply in the rage of the animal, and made a great many experiments to ſupport his hypotheſis.

It was natural to conceive, that the more a viper is enraged, the greater will be the diſeaſe produced by it, and *vice verſa*. But to draw a certain inference from this, it was firſt neceſſary to be aſſured, whether the degree of the diſeaſe, or intenſity of the venom, is in proportion to the rage of the animal: a very difficult experiment, and perhaps impoſſible to make well; and which would not probably have been yet ſufficient, ſince after all this might have been an accidental circumſtance, and not the true cauſe of what was obſerved.

Charas, who was ignorant of the true reaſon of the greater intenſeneſs of the diſeaſe in the caſes in which the viper is enraged, was miſtaken in his inferences. It is not ſurprizing that the naturaliſt ſhould here take that for the cauſe, which is the effect of the circumſtances that accompany it.

There are three reaſons why the bite of an enraged viper is more dangerous than that of one which is not enraged. The firſt is, that the more a viper is enraged, the deeper it forces its teeth into the ani-

mal; the ſecond, that it keeps them there a longer time; the third, that without letting go the part it has bit, it continues to contract the muſcles that compreſs the veſicle of venom.

When one has been ſome time accuſtomed to have animals bit by vipers, it is not difficult to perceive the truth of the firſt reaſon; and it is ſometimes even obſerved, that the tooth of the viper pierces the ſkin of the larger kind of quadrupeds with great difficulty, or only imperfectly and in part. All my experiments have ſhown me, that the diſeaſe is in general more violent, in proportion as the venom has introduced itſelf deeper into the ſkin and other parts of the animal.

The ſame obſervation likewiſe demonſtrates the truth of the ſecond reaſon. We frequently ſee that when a viper is violently enraged, it does not eaſily let go its hold; one might even ſay that it finds a difficulty in withdrawing its teeth. In theſe caſes it is eaſy to perceive, that during all this time the tooth not only prevents the venom from being thrown out again with the blood that naturally flows from wounds; but likewiſe, that it facilitates its union and mixture with the fluids of the animal.

The third reaſon is ſtill of greater weight than either of the other two. It has been ſeen, that ſeveral bites of the viper are neceſſary to empty the veſicle of the venom perfectly. It has been ſeen, that the firſt bites of the viper are nearly of the ſame activity, becauſe they are ſucceeded by the flowing

of nearly an equal quantity of venom. The cellular ſtructure of the veſicle does not allow it to be eaſily emptied, nor at once. When the viper keeps an animal a long time compreſſed by its teeth, and is very much enraged, it continues viſibly to contract the muſcles of its jaw. The muſcles which ſurround the veſicle alternately relax and contract without interruption, ſo that in theſe caſes we may reckon the bite of the viper, not as a ſingle one, but as ſeveral; and this may be carried to ſuch a length, that the viper, almoſt exhauſted of its venom, may not be capable of killing a ſmall animal.

It has been ſeen, that the firſt bites of the viper are all nearly of the ſame activity, and that it is only the laſt which exhibit a marked difference; and this I have accounted for.

After what has been ſaid, it is natural to conceive that the diſeaſe of the venom muſt be more dangerous, if the viper has bit the ſame animal ſeveral times. I have aſſured myſelf of the truth of this, by experiments the detail of which I ſhall not enter into here, as it would be tedious, and would not beſides anſwer any great purpoſe.

In inveſtigating this ſubject, I took care to employ animals of the ſame ſize and ſpecies, and had them bit by vipers ſimilar to each other. I more commonly availed myſelf of my uſual method, and the conſequences were ſtill more uniform. When only a few experiments are made, the conſequences may be equivocal, ſince it can ſcarcely hap-

pen

pen that the circumſtances will be perfectly the ſame. They may not only differ on account of the quantity of venom that remains in the animal's wound, which is ſubject to greater or leſs variations, but likewiſe becauſe it is very difficult to wound the ſame fibres, and the ſame veſſels. Theſe variations do not fail to occur; but in a great number of experiments, the circumſtances counterbalance each other, and ſo great a variety of conſequences preſents itſelf, that there is not the ſmalleſt danger of being miſled by them. Such has been at leaſt my opinion as to thoſe I have obtained.

A new enquiry to be made, was to know if the diſeaſe would be equal, whether a ſingle part was bit ſeveral times by a viper, or two different parts, provided the number of bites was the ſame.

This enquiry coſt me a vaſt many experiments, which I was obliged to make with the ſame circumſtances, only varying the part bit.

I not only tried it on birds, but on a great number of quadrupeds. I had them bit in the ſame part of their legs. I compared thoſe that were bit in both legs, with thoſe that were only bit in one, the total number of bites being the ſame in each animal.

Here again the conſequences were more or leſs conſtant. I was obliged to multiply my experiments till I conceived that I could advance, with great probability, the two following poſitions,

I. That an animal dies ſooner when bit a certain number of times in two parts, than when the ſame number of bites is confined to one.

II. That in this caſe the ſingle part is ſubject to a much more violent external diſeaſe.

By external diſeaſe, I mean the ſwelling of the part that has been bit, the livid and black colour of the ſkin and blood, and the wound that forms a ſhort time after the bite. Theſe ſymptoms are certainly more violent, when the part has been bit ſeveral times; although it is a fact, as will be ſeen hereafter, that the animals die much later, and that fewer of them die in proportion. It is however to be noticed, that this only happens when the animals live for ſome time after being bit, ſince otherwiſe the venom has not an opportunity of effecting any notable change in the external parts; in ſo much, that if the animal dies almoſt immediately after being venomed, there are ſcarcely any ſigns of local diſeaſe.

Before I examine the effects of the bite of the viper on the different parts of an animal, let me be permitted to relate the event of a great many experiments I made on animals of different ſpecies, which I had repeatedly bit, and by ſeveral vipers. In all theſe caſes I employed the fluid volatile alkali, either ſimply applied to the part bitten, or given internally at the ſame time. Theſe new experiments demonſtrate ſtill more the inefficacy of the volatile alkali, and how little dependence ought to be placed upon it.

I had

I had ſix fowls bit, each of them twice by a diſtinct viper. Three were ſimply treated, three were not. The three that were treated died at the end of 3, 5, 6, hours; the other three at the end of 3, 9, 12, hours.

I had ſix other fowls bit, each by two diſtinct vipers, once in each leg. I treated them, and made them ſwallow the volatile alkali. They were all dead before the expiration of ſeven hours; one of them died in leſs than twenty-ſeven minutes.

Twelve other fowls were bit, each of them twice in the leg, and by different vipers. Six only were treated, and ſwallowed the volatile alkali. Nine died in the whole; five of thoſe that were treated, and four of the others. Two of theſe laſt lived forty-three hours; the treated five that died did not ſurvive the ſeventh.

The reſult of the laſt claſs of experiments, although it does not agree with that of the two that precede it, is nevertheleſs given with preciſion. This ſhows how much experiments of this kind may differ from each other, from circumſtances which occaſionally vary, and which cannot always be aſcertained. Thoſe that are capable of influencing the moſt are, that vipers are not always provided with the ſame quantity of venom, and that they are more or leſs vigorous in biting, and in forcing this humour from the veſicle: to theſe may be added, the effect of a milder or ſeverer ſeaſon. I began my experiments in September, and continued them with more or leſs earneſtneſs till the cloſe of

the January following. I likewiſe made a few in February, March, and April, and found a ſenſible difference at theſe different times. During the ſevere froſt, the vipers were ſo weak, that it was with difficulty I could get them to bite; and their bites were in a very ſmall degree dangerous.

I cannot here paſs over an experiment I made in the month of January, and which at firſt made me ſuſpect that the volatile alkali might ſometimes be a remedy againſt the bite of the viper.

I had ſix fowls bit in the leg, each by three vipers, all of which bit three times ſucceſſively. I treated them ſeveral times, and made them as repeatedly ſwallow the volatile alkali. They all had the diſeaſe of the venom, but in a very ſlight degree, and recovered in a few days.

There remained, as chance would have it, in the ſame box, eighteen other vipers, perfectly like the eighteen employed in the preceding experiment. Perceiving at the end of fourteen hours that neither of the fowls was dead, and that they were all but ſlightly diſeaſed, I had ſix others bit in the ſame way, each by three of theſe vipers, and each viper biting three times. I treated neither of them, and only one died, at the end of ſix days. Two had ſcarcely any complaint, and the three others recovered on the third day. This experiment demonſtrates clearly, that the ſix treated fowls were not cured by the volatile alkali, but that their recovery was owing to the little vigour and activity of the vipers by which they were bit.

The

The fowl in the laſt experiment that was not treated and died, argues nothing in favour of the volatile alkali, ſince it is only one out of ſix, and ſince it did not die till the end of the ſixth day. This evidently proves, that if the venom had been in a ſomewhat ſmaller quantity the fowl would not have died. We have ſeen above, that a thouſand accidents may vary this greater or leſs quantity of venom, both in the viper that inflicts the bite, and in the animal that receives it.

On this very account, I have made it a maxim, in almoſt the whole courſe of this work, to form comparative experiments, and only to compare thoſe with each other, that were made at the ſame time and with the ſame circumſtances.

I muſt here inform my readers of what befel the vipers I employed laſt. The ſeaſon was very cold, and notwithſtanding the temperature of my chamber was twelve degrees above the freezing point, the vipers were very ſluggiſh and inactive. I conceived that I could give them a freſh vigour by additional warmth, and therefore, after keeping them in my laboratory for upwards of ſix hours, in a box pierced with holes, I at length placed the box on a ſand heat, the warmth of the ſuperficies of which was only twenty degrees. At the end of two minutes I found every one of the vipers dead. The ſame accident happened to me twice beſides, in the ſame month, and on an occaſion ſomewhat ſimilar.

Expe-

Experiments on Guineapigs bit ſeveral Times, and by ſeveral Vipers.

I had two very large guineapigs bit repeatedly in the leg by two vipers. One was treated, the other was not. They both died; the firſt at the end of two days, the ſecond of thirty-two hours.

I had four other guineapigs, preciſely of the ſame ſize, bit each in the leg by three vipers, three times by each. Two were treated, and ſwallowed the volatile alkali; the other two were left to themſelves. All four died in leſs than two days.

Again, I had four others of the ſame ſize bit in the ſame way. They were not treated. One only died, at the end of the fifth day.

Twelve very ſmall ones were bit in the ſame way. Six were treated, and ſwallowed the volatile alkali; the other ſix had nothing done to them. They all died in the ſpace of twenty minutes.

Two days after, I had twelve others bit, of the ſame ſize as the laſt, each receiving from two diſtinct vipers three bites in each leg. Six were treated, and ſix not. They all twelve died in two hours. One of the treated ones died in ſeven minutes, and two of thoſe that were not treated in fourteen.

Theſe experiments convince us at a glance of the inutility of the volatile alkali. They likewiſe ſhow, that in animals of this ſpecies the ſmaller ones

ones die ſooner than the larger, and that their deaths are ſpeedier and more certain, in proportion to the greater number of the viper's bites.

Experiments on Rabbits bit ſeveral Times, and by ſeveral Vipers.

I had four middle-ſized rabbits bit, each four times in the leg, by two diſtinct vipers. I treated two of them, which I made ſwallow the volatile alkali every two hours, repeating the application as often. They all four died; the two that were treated, in eighteen hours, the other two at the end of three days. In all of them the diſeaſe of the venom was very violent, and their legs were very much ſwelled.

I had four very large rabbits bit, each by two vipers, twice in the leg. Two were treated, and two not. The two that were treated, although they ſurvived, continued ill and with open wounds in their legs, for upwards of twenty days. One of the two that were not treated died on the third day; the other recovered on the tenth.

I had a dozen middle-ſized rabbits bit in the leg, each by two diſtinct vipers, and each viper biting three times. Six were treated, and ſix not. Four of the former died, and five of the latter.

Theſe conſequences not being either ſufficiently uniform, or in a ſufficient number, to enable me to

decide

decide as to the volatile alkali. I judged it neceſſary to have recourſe to new experiments.

I had twelve rabbits, ſomewhat ſmaller than thoſe employed in the laſt experiment, bit in the ſame way. Six of theſe were treated, and ſwallowed the volatile alkali; the other ſix were left to themſelves. All of the former ones died, and five of the latter; the ſixth had ſcarcely any perceptible complaint.

I wiſhed to ſee whether there would be a ſenſible difference betwixt the effects of the venom, on animals bit a greater or leſs number of times, by a greater or leſs number of vipers. For this purpoſe, I had ſix middle-ſized rabbits bit, each once in the leg, by a diſtinct viper. I had ſix others bit in the leg each by two diſtinct vipers, each of which made two bites. I had ſix others bit in the ſame part, each by two diſtinct vipers, each of them biting four times; and ſix others again, each of which was bit by three diſtinct vipers, four times in the leg by each.

Of the ſix of the firſt claſs, three died; the other three had moderate complaints. Of thoſe of the ſecond, five died, and the other had a violent attack of the diſeaſe. All of the third claſs died in in leſs than forty-three hours; and thoſe of the fourth in leſs than twenty.

Expe-

Experiments on Dogs, bit ſeveral Times, and by ſeveral Vipers.

I had two ſmall and young dogs bit in the leg, each by two diſtinct vipers, and twice by each. One was treated, and ſwallowed the volatile alkali; the other had nothing done to it. They both died in the ſpace of thirteen hours.

I had two dogs, larger by one-half than the preceding ones, bit each by two diſtinct vipers, and twice by each. One was treated, the other not. Both recovered; the treated one in twenty-ſix days, the other in ten.

I had four very large ones bit, each by three diſtinct vipers, three times by each. Two were treated, and two not. One of thoſe that were treated died at the end of the ſixth day. The other three were exceedingly ill, and had each of them a large wound in the leg that had been bit.

Two very large dogs were brought to me in excellent order. I had each of them bit in the leg by four well-irritated vipers, each viper biting at leaſt four times. I did not treat them, on account of the difficulty of doing it effectually without the riſk of being bit. Both of them recovered in leſs than ten days. They had wounds, tumour, and lividity, in the part bit. At the end of two days they began to drink, and ate at the end of the third.

Scarce'y

Scarcely have animals of any kind, and particularly dogs and cats, been bit by a viper, and are at liberty, when they lie down on the part oppoſite to that which has been bit, and in this ſtate continue very quiet till they recover. Whenever they begin to drink and to eat, 'tis an almoſt certain ſign that they will get the better of their complaints. Cats are leſs deſirous of food than dogs; I have met with ſome that did not eat till after they had been ill ſeveral days.

That the number of my experiments on dogs might be competent to the purpoſe, I procured ſix ſmall ones, of the ſame ſize, ſpecies, &c. I had them all bit in the leg, each by three vipers, and each viper at three bites. Three were treated, and three not. The three firſt died, and only two of the others; the third was exceedingly ill, had a large wound, and did not recover till the end of the fifteenth day.

Not perceiving that the volatile alkali had any good effect againſt the bite of the viper, when given to dogs, I thought it proper to purſue my experiments on other kinds of animals.

Experiments on Cats.

This animal makes a very ſtrong reſiſtance to the bite of the viper. This is not becauſe the venom is innocent to it as it is to ſome other animals, but becauſe it is very hard to kill.

I had

I had a middle-ſized cat bit in the leg by two vipers, each viper biting twice. I did not treat it. Its leg ſwelled, but not violently. It lay reclined on its belly during the whole time of its illneſs; it drank at the end of thirty-ſix hours, ate at the end of fifty-two, and on the fourth day was perfectly recovered.

I had it bit in another leg by three vipers, each viper biting twice. Here again I did not treat it. It vomited ſeveral times after the ſixth hour, and again after the thirtieth. It drank at the expiration of forty-two hours, and ate at the cloſe of the third day. On the fifth day it was recovered.

I made choice of another cat of the ſame ſize as the former one, and had it bit in the leg by four vipers, each viper biting four times. I did not treat it. It ſwelled very much, vomited ſeveral times, and did not eat till the cloſe of the ſixth day.

Two days after I had it bit by four freſh vipers in another leg. It was very much diſeaſed, and had frequent vomitings. It ate at the end of five days, and on the eighth was quite recovered.

I had another cat, larger than the former ones, and very wild, bit by ſix well-enraged vipers, ſeveral times by each. One of them could not let go its hold, and was diſengaged with ſo much difficulty, that its teeth were broken and left in the fleſh. The cat was in a violent rage, but became tranquil on being ſet free. It reclined itſelf on its belly, as the others had done, vomited from time to time, and did not eat any thing till after the fifth

day. It continued ill two days more, and at length recovered.

It was quite unneceſſary to give the volatile alkali to the cats, ſince, as we ſee, when they are of a certain ſize, they do not die of the diſeaſe of the venom. Kittens are, however, known to die of it; and it is likewiſe certain, that grown cats would die too, provided they were bit by a greater number of vipers.

The bite of the viper produces a true diſeaſe in this animal, and this diſeaſe is more violent in proportion to the greater number of bites. I cannot, however, preciſely ſay, how many vipers it would require to kill a ſtrong cat of the largeſt ſize. Ten or twelve would, perhaps, be ſcarcely ſufficient.

CHAPTER IV.

Of the Effects of the Bite of the Viper on different Parts of an Animal.

I HAVE hitherto ſpoken of animals bit by one viper, or by ſeveral, either once or repeatedly, but only in a ſingle part; that is to ſay, in the leg, or in two legs at moſt. We are now to ſee the effects of the bite of the viper on the other parts of an animal. It is eaſy to conceive that the conſequences will be ſomewhat different from thoſe that have already been obſerved, and that there muſt be parts in the ſame animal, more or leſs ſuſceptible of the venom; ſeveral of theſe, on having them bit, have afforded ſingular and unforeſeen appearances.

Experiments on the Skin.

The part of an animal which is firſt pierced by the canine tooth of the viper, and which feels before the others the action of the venom, is the ſkin. I have confined my experiments to the ſkin of guinea-pigs and rabbits, harmleſs animals, that are managed without riſk. I have not employed birds, as their ſkin is too delicate for theſe experiments.

Wounds made in the ſkin may be very ſlight, and altogether external; they may be more or leſs deep; and laſtly, they may pierce the ſkin through and through. I have obſerved all theſe caſes in the courſe of my experiments on the bites of the viper. I have ſometimes ſeen the viper's tooth ſtrike the ſkin ſo obliquely, that it was either not cut at all, or only ſuperficially. The firſt caſe I mentioned, happens frequently, from the viper, when it is enraged, biting at every thing that is preſented to it, in any way, and under any form whatever. The ſecond caſe is much leſs frequent; and that in which the bite is made without piercing the ſkin, ſtill rarer.

Theſe two laſt caſes may happen to man, whoſe ſkin may be more or leſs injured by the canine teeth of the viper.

This reſearch, beſides its being curious, may likewiſe be uſeful in practice, by aſſiſting to make the quality of the venom well underſtood in theſe caſes. Such an inveſtigation, well handled, may likewiſe ſerve, as will be ſeen in the ſequel, to explain the action of the venom of the viper on animals in general.

Superficial Wounds of the Skin.

I ſat out by making the following experiments. I cut the hair with ſciſſars from the ſkin of a part of the leg of a guineapig, and rubbed a portion of this, of about half an inch in length and breadth,

ſeveral times, with a ſmall file. The ſkin became red, and an almoſt imperceptible quantity of blood exuded from it, which could not, however, form itſelf into entire drops. Having wiped it well, I poured on it with a large drop of venom, which, to make it flow eaſier, and to extend itſelf over the whole ſurface of the raſped ſkin, I had united with a drop of water.

The animal did not appear to ſuffer in the leaſt, and there was ſcarcely any perceptible mark of cicatrice. On the following day, obſerving it to continue ſound and vigorous, I had it bit twice in the foot by a viper. It died at the end of twenty-four minutes. This experiment I repeated twice, with nearly the ſame reſult; both guineapigs died after being bit.

I ſhaved the hair with a razor, from the external lateral part of a guineapig's leg. The ſkin was red, and a little moiſture exuded from it, which was likewiſe of a reddiſh tinge. I put two drops of venom on this part, the ſize of which was about two thirds of an inch. The animal did not ſuffer the ſmalleſt inconvenience, and the ſkin dried without eſchar or cicatrice. On having it bit in the foot the next day, it died at the end of twenty-ſix minutes.

I removed the hair with boiling water from a portion of the back of a guineapig, and made two very ſmall, but very deep, inciſions in it, wiping away the blood that flowed from them. I applied two drops of venom, unmixed with water, to the inciſed ſkin, which was eaten away for half its thickneſs,

by a wound that formed over the whole ſurface the venom had touched. This wound diſcharged pus, and the next day was covered with an eſchar. The animal was perfectly recovered in ſix days, and on the ſeventh, on having it bit by a viper once in the foot, it died at the end of forty minutes.

I repeated this experiment, with the ſame circumſtances, as nearly as I could judge, on two other guineapigs. The effects were exactly the ſame; wound, ſkin conſumed for half its ſubſtance, pus, eſchar, and recovery. On having them afterwards bit in the foot, they both died in leſs than an hour.

I likewiſe wiſhed to make a ſimilar experiment on an animal with a ſkin much thicker than that of a guineapig. I choſe a very ſmall rabbit, and removed the hair with a razor, in ſuch a way, that there was a ſenſible diſcharge of blood. I applied to this part, about half an inch in length and breadth, two drops of venom. A true wound formed, and the ſkin was entirely conſumed, and covered with a great deal of pus. The rabbit notwithſtanding did not ſeem to ſuffer much, and at the end of ſeven days was perfectly recovered. I had it twice bit in its leg by a viper, and it died at the end of ſix hours. I repeated the ſame experiment on two other rabbits, with the ſame ſucceſs.

The following concluſions may, I think, be drawn from the above experiments:

I. That the venom of the viper, applied to the ſkin of guineapigs and rabbits, ſlightly ſcraped or punctured, is not mortal.

II. That

II. That it produces but a ſlight diſeaſe in the ſkin of guineapigs, and a ſomewhat greater one in that of rabbits.

III. That this diſeaſe is confined to the part of the ſkin touched by the venom.

I was deſirous of making a ſomewhat different experiment on the ſkin of guineapigs, and accordingly removed the hair with ſciſſars from a portion of the back of one of theſe animals, of about the breadth of half an inch. I then made an inciſion with a lancet, ſo as not to puncture it through, but only for about half its thickneſs, and applied two drops of venom. A wound, occupying the whole ſpace covered by the venom, formed, and ſuppurated very abundantly, and the ſkin was entirely conſumed, and covered with a ſcar. The animal did not appear to ſuffer otherwiſe, ate conſtantly, and recovered at the end of ten days.

This laſt experiment ſeems to indicate that when the wounds of the ſkin are deep, the effects of the venom, or its diſeaſe, are more conſiderable, although not mortal; and likewiſe, that the diſeaſe is entirely confined to the ſkin.

Wounds in the Skin, through its whole ſubſtance.

I pinched the ſkin of a ſmall rabbit's leg with my thumb and finger, and pierced it five or ſix times with a viper's tooth, from which the venom flowed. At the end of twelve hours, an encyſted tumour, filled with matter, formed in the ſkin, an inch below the

wounds. The cyſt was quite excoriated and bare of hair, and a little moiſture exuded from it. The rabbit died on the fifth day.

I repeated this experiment on a rabbit of the ſame ſize, pricking the ſkin ſeveral times with a venomous tooth. At the end of ten hours, the ſame kind of tumour formed in the ſame place; on the ſecond day the ſkin fell off; on the third the tumour burſt; and the rabbit died four hours after.

I treated two other ſmall rabbits in the ſame way, and the effect was perfectly the ſame. In both of them a tumour formed, and burſt; and both died.

I had the ſkin of a guineapig's back bit repeatedly by a viper, raiſing it with pincers, to prevent the muſcles beneath from being wounded. In leſs than two hours, the part that had been bit became livid, and the animal died at the end of thirty-two hours, without an open wound. The ſkin was gangrened, and the blood black and extravaſated in the adipoſe membrane, as far as the muſcles of the abdomen and breaſt.

I repeated this experiment with the ſame circumſtances on four other guineapigs, all of which died. Neither of them had any wound, but the adipoſe membrane had a gangrenous appearance, and was filled with black extravaſated blood. The extravaſation was extended to the adipoſe membrane which covers the pectoral and abdominal muſcles, and was in ſuch a quantity as to form a bag.

Experiments

Experiments on the Adipoſe Membrane.

The preceding experiments not only relate to the ſkin, but likewiſe to the adipoſe membrane. Whenever the tooth pierces through the whole ſubſtance of the ſkin, the venom muſt neceſſarily communicate itſelf to this membrane; and its effects, or the diſeaſe it occaſions, will be communicated to both parts. It was therefore neceſſary to have the adipoſe membrane wounded apart, to know what related to the ſkin in the above experiments. It is not very eaſy to do this with preciſion and nicety.

I made an inciſion in the ſkin of a guineapig, near the groin, and introduced a drop of venom without its touching the ſkin. It brought on a tumour of the groin, which increaſed for two days. The third day the animal died. On opening the tumour, I found it filled with a great quantity of black, diſſolved, and extravaſated, blood.

I repeated this experiment on two more guineapigs, one of which died, the other did not. This laſt had ſcarcely any tumour. The one that died had a large one; with the ſame ſymptoms as in the preceding experiment. Two days after, I opened the one which ſurvived, and which appeared ſound, and in good health. I found the adipoſe membrane ſomewhat bloody, and with ſome humours extravaſated in it; but all this in a ſlight degree. There was no appearance that could induce one to conclude, that the animal would afterwards have died

of the diſeaſe of the venom. It was vigorous, fed well, and ran about in good health, whilſt the other was in the heighth of its diſeaſe at the end of two hours after being bit.

Theſe experiments ſtill leave us in a doubt whether the venom might not have been communicated to the inciſed edges of the ſkin. To obviate this, I fell upon ſeveral modes of experimenting, but invariably met with difficulty in the attempts, and ſomething equivocal in the conſequences.

After ſeveral trials, I purſued the following method:—I cut away a large portion of ſkin from the back of a guineapig, dried the adipoſe membrane well, and applied to it two drops of venom. The circular piece of ſkin I removed was more than an inch in diameter. I ſpread the venom on the membrane for about three lines in circumference, and at equal diſtances at all ſides from the ſkin.

In leſs than ſix hours, the adipoſe membrane became black as ink, and at the end of twelve it was covered with an eſchar, which continued ſo long as twenty-two; the animal recovered notwithſtanding.

I repeated this experiment on ſix ſmall rabbits, and ſix ſmall guineapigs, and the conſequences were ſomewhat different from each other.

In the firſt place it muſt be remarked, that neither of theſe animals died. Six of them were very much diſeaſed, and recovered very late. Four had ſlight ſymptoms of the diſeaſe of the venom, and ſeemed to be perfectly recovered at the end of the ſecond day. The others had no certain ſymptoms

of diſeaſe. I think it may be ſaid, in a general way, that the venom of the viper is not mortal, if it penetrates no farther than the adipoſe membrane.

Experiments on the Muſcles.

I ſtripped the exteriour muſcles of a pigeon's leg of the ſkin and adipoſe membrane, without producing any ſenſible hemorrhage. I introduced into one of theſe muſcles a viper's tooth filled with venom. A minute after the pigeon fell forward, and died at the end of ten. The wounded muſcle was extremely livid, throughout almoſt the whole of its ſubſtance.

I repeated this experiment on four other pigeons, all of which in leſs than two minutes fell forward. One died at the end of eleven minutes; another at the end of ſeventeen; the third in a quarter of an hour; and the fourth not till four hours.

I ſtripped ſeveral muſcles of the leg of a middle-ſized rabbit of the ſkin and adipoſe membrane, and wounded them ſeveral times with venomous teeth, *(a)* in ſuch a way that they entirely entered the muſcles. I wounded them at the parts where there did not appear any conſiderable veſſels. There was ſcarcely any diſcharge of blood from the muſcles, which, notwithſtanding, very ſoon became livid at

(a) Theſe are viper's teeth detached from the animal, but ſtill adhering to the veſicle filled with venom. I have already explained the method I purſue in experiments of this kind.

the

the places I had wounded. The animal not only ſurvived, but diſcovered no ſigns of ſuffering any great inconvenience; and at the end of fifteen hours there was ſcarcely any diſcoloration in the wounded muſcles. At the end of thirty hours, nothing was to be ſeen but the mechanical wound of the ſkin, where the inciſion had been made to come at each of the muſcles.

I repeated this experiment, with the ſame circumſtances, on another rabbit. The muſcle became diſcoloured, but not much; and the animal, at the end of twenty-three hours, ſeemed to be free from all complaints, except that there ſtill remained a ſolution of continuity in the ſkin.

I entirely ſtripped ſeveral muſcles of a guineapig's leg of the ſkin and adipoſe membrane, and plunged a tooth, charged with venom, betwixt the fibres in ſuch a way, that few or no veſſels were divided. The muſcle became livid, but the animal recovered.

I repeated this experiment on the bared muſcles of ſeveral other animals, ſuch as guineapigs and rabbits, and found that in theſe caſes the venom of the viper does not fail to bring on a complaint, which, although it is frequently very violent, is never mortal.

The Venom of the Viper, when ſimply applied to the Muſcular Fibres, is entirely innocent.

I wiſhed to know what would be the effects of the venom, when ſimply applied to the muſcles, without cutting the fibres.

I ſtripped the muſcles of a pigeon's leg of the ſkin, and contrived in ſuch a way, that the uncovered fibres and veſſels did not bleed ſenſibly. The experiment ſucceeded ſo well, that the muſcles, ſtripped of the adipoſe membrane, appeared perfectly dry. On theſe muſcles I laid a large drop of venom, obſerving that it did not communicate itſelf to the adjacent parts. The pigeon had no complaint, and the wound I had made healed very ſoon.

I got ready another pigeon in the ſame way, but took care that the muſcles ſhould bleed a little; one vein in particular bled conſiderably. I applied the venom, and the pigeon died at the end of thirty hours, with a very ſlight change in the parts that had been wounded.

I repeated this experiment on four other pigeons, the muſcles of which did not bleed. Neither of them died, nor ſeemed to have any other complaint than that occaſioned by the inciſion in the ſkin.

When we know how ſmall a quantity of venom is capable of killing a pigeon, as it were, inſtantly, we cannot heſitate to pronounce, that the venom of

the

the viper, when ſimply applied to the muſcular fibres, is entirely innocent.

The Venom of the Viper does not loſe its deadly Qualities, even after it has acted on an Animal as a Poiſon.

I was deſirous of ſeeing whether the venom of the viper, after having communicated the diſeaſe to one animal, would act as a poiſon on another. To aſſure myſelf of this, I laid the muſcles of a pigeon's leg bare, and made ſmall inciſions in them, into which I introduced about a drop of venom.

I likewiſe got ready another pigeon, making ſmall inciſions in its muſcles, as I had done in thoſe of the firſt. At the end of four minutes I put the bared muſcles of the two pigeons in contact, and kept them in that ſtate for two minutes. Neither of the pigeons died: the firſt, however, was very ill; the ſecond had ſcarcely any complaint.

I laid the muſcles of two other pigeons bare, and made ſmall inciſions in them. I wounded thoſe of one with a venomous tooth, and at the end of four minutes put them in contact with thoſe of the other, keeping them together in this way for three minutes. The firſt pigeon died at the end of three minutes more; the ſecond at the end of an hour.

I repeated this laſt experiment on two other pigeons. The one that was venomed by the tooth

died—

died at the end of eight minutes; the other at the end of eighteen.

Conſequently the venom of the viper, as in all the caſes related above, continues to be ſuch, and does not loſe its deadly qualities, when it mixes with the blood of living animals, and excites in them the uſual diſeaſe.

Animals bit in the Breaſt.

I had a pigeon bit once by a viper in the breaſt. I treated it, and it died at the end of ten minutes.

I had another pigeon bit twice in the breaſt by a viper, and treated it. It died at the end of two hours.

I had ſix pigeons bit in the breaſt by as many vipers, each twice by a diſtinct viper. Three were treated, and three not. They all died; the three that were treated at the end of 10, 20, 50, minutes; the other three at the end of 17, minutes, and 2, 4, hours.

I had ſix others bit an equal number of times; three in the breaſt, and three in the leg. They all died; the three bit in the leg at the end of 10, 15, 20, minutes; the three in the breaſt at the end of 17, 50, minutes, and two hours.

Theſe few experiments on pigeons would lead one to ſuſpect, that bites in the breaſt are not more dangerous than thoſe of the leg; and that it may even be the reverſe. They are however not ſuffi-

cient

cient in number to admit of any certain conſequences being drawn from them.

I had a guineapig bit twice in the breaſt by a viper, and immediately treated it. It died at the end of two hours.

I had another guineapig, of a much larger ſize, bit twice in the breaſt by a viper, and treated. At the part where it was bit, a large wound, which continued open for upwards of fifteen days, formed; the guineapig at length recovered.

I had a very large guineapig bit twice in the breaſt by a viper, and treated it immediately. It had no ſymptom of diſeaſe. Two days after I had it bit afreſh by another viper, in the ſame place, and at the end of twelve hours it died.

The ſkin of guineapigs, particularly that part of it which covers the breaſt, is very tight, in conſequence of which the viper finds it very difficult to ſeize it betwixt its teeth. I was ſeveral times deceived by this, ſuppoſing the animal bit when it was not; and was therefore obliged to repeat the experiment.

I had a ſmall rabbit bit in the breaſt by a viper, and immediately treated it. At the end of thirty ſeconds it fell on its belly, and died in leſs than a minute.

I had another rabbit, of the ſame ſize, bit in the breaſt, and did not treat it. It had a ſmall wound, and recovered at the end of three days.

I had four rabbits bit in the breaſt, each twice by a diſtinct viper. Two were treated, and two not.

The

The two that were treated died; one at the end of an hour, the other at the end of ten. Of thoſe not treated, one died in an hour, the other had a very ſmall wound in the part bit.

I had a fowl bit twice by a viper in its breaſt, towards the right wing. I treated it, and it died at the end of twenty-four hours.

I had another fowl bit twice by a viper in the ſame place, and did not treat it. It died at the end of nine hours.

I had four other fowls, like the preceding ones, bit, and obſerved the ſame circumſtances. They all four died in eighteen hours.

I had four other fowls bit, two in the breaſt, and two in the leg. The two that were bit in the breaſt died in leſs than ten hours. One of thoſe bit in the leg died at the end of twenty-ſeven hours; the other was violently diſeaſed, but recovered.

Had the number of experiments been greater, we might have deduced from them, that to fowls the bite of the viper in the breaſt is more dangerous than that in the leg. This is contrary to what was obſerved in the rabbits and guineapigs.

Animals bit in the Belly.

I had a rabbit bit twice in the belly by a viper. At the end of eighteen hours a very large tumour formed in the part bit. This tumour encreaſed for four days, and the hair fell from the ſkin, which

was

was corroded and ulcered. The animal, nowithſtanding, lived twenty days.

I had another rabbit, of the ſame ſize, bit repeatedly in the belly by a viper. At the end of twelve hours a tumour formed, and the hair and epidermis came away. The tumour was moiſt and bloody, and burſt at the end of eighteen hours, when an ulcer formed, of two inches and an half in length, and more than an inch in breadth. The rabbit ſurvived, but it was more than twenty days before it recovered.

I had two others bit in the belly in the ſame way. Both of them had a tumour, which was ſucceeded by an ulcer that remained open for ſeveral days; and both recovered.

I had two other rabbits of the ſame ſize bit ſeveral times in the belly by two vipers. One died at the end of twenty-ſix hours; the other had a wound which covered the whole of the ſkin of the lower part of the belly, and continued ill twenty-ſix days.

Experiments on the Inteſtines.

I opened the belly of a rabbit, and had the *ileum*, at the diſtance of three inches from the *colon*, bit twice by a viper, binding up the part as well as I could. The rabbit died at the end of ſix hours. The inteſtine was inflamed, black, and contracted, more than ſix inches above and below the part that was bit; ſo that theſe changes had extended to the

colon.

colon. The *meſenterick veſſels* were black and ſwelled, and the blood curdled.

I repeated this experiment on four other rabbits, each of which I had bit in the inteſtines in the ſame way by a viper. The reſult of theſe experiments was perfectly analogous to that of the former one.

Experiments on the Liver.

Having opened the belly of a rabbit, I wounded the right lobule of the liver, in the inner part, with a venomous tooth. At the end of a few ſeconds, the creature began to cry and to writhe itſelf, and died in leſs than two minutes. All the veſſels of the liver were filled with black and clotted blood; the meſentery was in the ſame ſtate; and the heart and auricles were filled with black, but fluid, blood.

I wounded the outer lobule of the liver of another rabbit in two places with a venomous tooth. The creature drew itſelf together, but did not cry. It died an hour after.

I introduced a venomous tooth into the outer lobule of the liver of a third rabbit, and did not withdraw it immediately. This one cried, writhed itſelf, and died in leſs than a minute and an half. The blood was coagulated both in the liver and meſentery.

I introduced a venomous tooth in the uſual way into the inner lobule of the liver of two other rabbits, and kept it there for ſome time. Theſe ani-

mals, as uſual, cried out after a few ſeconds, and died in leſs than two minutes. The blood in the liver was black and coagulated; that in the heart and auricles was likewiſe black, but in a fluid ſtate. I did the ſame to the outer lobule of the liver of two rabbits, but withdrew the venomous tooth immediately after having introduced it. One began to cry and writhe itſelf after a few ſeconds, and died in two minutes. The other lived nearly two hours. The blood in the liver of the firſt was quite coagulated; as it was in a degree in the ſecond. In the former, the blood in the auricles and ventricles was fluid; in the latter it was coagulated.

Experiments on the Ears.

I had the ear of a middle-ſized rabbit, towards its extremity, or point, bit twice by a viper. The part was a little ſwelled at the end of ſix hours; the rabbit, however, ate, and was lively. At the end of four days it was perfectly recovered.

I had two other middle-ſized rabbits bit in the ſame way at the extremity of the ear, each twice by a diſtinct viper. The ears ſwelled a good deal, but the rabbits ate, and were lively. At the end of five days they were both recovered.

I had another rabbit bit in the right ear, towards its extremity, twice by a viper. I treated the part, in which there was a conſiderable ſwelling that did not ſubſide till after ſixteen days.

I had

I had a rabbit's ear bit twice by a viper, at a third of its length from the baſis. At every hole made by the teeth in the oppoſite ſides of the ear, a drop of blood appeared, and beſide it a ſmall drop of venom, which, although it was in contact with the blood, did not unite with it in the leaſt. There were four holes made by the teeth at each ſide the ear, ſo that the ſmall drops of venom were eight in number. The ear ſwelled a good deal, and the ſwelling did not ſubſide till after twenty days.

There is no difficulty in accounting for the ſmall drops of venom that appeared at the oppoſite ſides of the ear. We know that the venom flows from the point of the tooth. The ear of a middle-ſized rabbit is not ſo thick as the viper's tooth is long, which muſt of courſe pierce the ear through. When the viper withdraws the tooth, the venom has already reached the point of it; and from the elaſticity of the ſkin of the ear, which cloſes the hole it went out at, is forced to ſhed itſelf at the ſides of it. In finding its way to the part of the ear at which it entered, the tooth in the ſame way leaves the venom, which it continues to ſhed, at the edges of the hole at this ſide. I have ſince obſerved theſe ſmall drops of venom on each ſide the ear in almoſt all the rabbits I have had bit in this part, and find them in general to be larger at the part the tooth went out at, than at that where it entered; particularly if the viper is prevented from withdrawing its teeth too ſuddenly.

I had

I had a rabbit bit in both ears at a third of their length from the baſis. Each ear was bit three times by a diſtinct viper, and both of them ſwelled violently, for nearly eight lines towards the baſis. The rabbit was very much diſordered, and did not eat for ſeveral days, when it began to feed ſparingly. It was not perfectly recovered till the end of twenty days, and was then very much waſted.

I had two others bit repeatedly at the ſame part of the ear, by two vipers. At the end of the ſecond day, the ears were diſfigured by a ſwelling, which became ſo large, that in two days more they hung down on each ſide the neck. One of the rabbits died at the end of eight days, with its ears ulcered and ſphacelated; the other recovered, but not till the end of twenty-eight days.

I had a middle-ſized rabbit bit once in the ear by a viper. The ear bled a little, and two ſmall drops of venom appeared at the ſides of the two holes made by the teeth. I did not treat it. There was a degree of tumour and inflammation in the part, and at the end of thirty hours the rabbit was perfectly recovered.

I had another rabbit bit, of the ſame ſize as the preceding one. I treated it immediately, and made it ſwallow the volatile alkali. The ear ſwelled exceedingly, and became livid at the part where it was moſt ſwelled. The tumour continued ſix days, and in four more the animal recovered.

I had four rabbits bit in the ears by as many vipers. Two were treated, and two not. Neither of them

them died. The ears in all of them ſwelled conſiderably, and they all recovered at the end of three days.

Having aſſured myſelf that the bite of the viper in the ears of rabbits is not very dangerous, I thought of having theſe animals bit by ſeveral vipers, in different parts of the two ears. For this purpoſe I choſe a dozen middle-ſized rabbits, and had them all bit repeatedly in ſeveral parts of each ear, and each by three vipers. The parts ſwelled exceedingly, and continued in that ſtate for upwards of twelve days. Three of the rabbits had an enormous bag or tumour in the fore part of the neck, larger than the head itſelf. Theſe tumours were filled with a humour, and yielded to preſſure. At the end of two days they increaſed in ſize, and the ears became ulcerous. The rabbits recovered in ſixteen days.

Experiments on the Pericranium.

I laid bare the pericranium of a pigeon, by removing a good portion of the ſkin, and made ſeveral ſmall inciſions into it with the point of a lancet. I poured venom upon it, but in ſuch a way that it did not reach to the adjacent parts of the integuments that had been cut. The pigeon did not ſeem to be at all diſordered by it, and recovered in the ſame ſpace of time as another did, which I had prepared by way of compariſon, and to the pericranium of which I had not applied the venom.

I repeated this experiment on four other pigeons, with the ſame ſucceſs. Neither of them died, and in neither was the attack of the diſeaſe of the venom perceptible.

On the Bones and Periosteum.

I laid bare the cranium of a pigeon, ſtripping off a good part of the pericranium. I made ſmall wounds in the cranium with a lancet, taking care not to pierce the whole ſubſtance of it, and introduced a conſiderable quantity of venom, preventing it as uſual from communicating to the neighbouring parts. The animal not only ſurvived, but did not appear to have ſuffered the ſmalleſt inconvenience.

The conſequences of three experiments on pigeons, treated in the ſame way, were the ſame.

Having laid bare the *tibia* of two pigeons, and freed it well of the cellular membrane, I wounded both perioſtæum and bone in ſeveral places with the point of a needle, and poured the venom copiouſly upon them. The pigeons had not the ſmalleſt perceptible complaint, and recovered in the ſame time as two others did, that I had treated in the ſame way, but without applying the venom, to ſerve as a compariſon.

I repeated this experiment with the ſame circumſtances on two other pigeons, and the reſult was exactly the ſame. Neither of them died, nor had the ſmalleſt ſymptom of the diſeaſe of the venom.

I laid bare the perioſtæum of the *tibia* of ſix other pigeons, and having pierced it in ſeveral places with a needle, moiſtened it with venom. Neither of the pigeons died, nor had any complaint.

Dura Mater and Brain.

I removed a portion of the cranium of a pigeon, taking care to lacerate the dura mater as little as poſſible. I wiped this membrane, which was well expoſed, as gently as I could, with dry lint, and applied to it a drop of venom. The pigeon had no ſymptom of the diſeaſe of the venom, and recovered within the ſame ſpace of time as another did, that I had prepared in the ſame way, but without applying the venom, as a compariſon.

This experiment, on two other pigeons treated as above, terminated in the ſame way.

I removed a portion of the cranium of a pigeon, and made inciſions in the dura mater all round, introducing at one of the apertures a drop of venom The pigeon recovered, without having had any ſymptom of the diſeaſe of the venom.

After having removed the dura mater of another pigeon, I made a ſlight inciſion into the brain, and introduced the venom. The animal recovered in the ſame way with the preceding one.

A third pigeon, on which I made the ſame trial, died at the end of four hours.

Marrow of the Bones.

I cut the *tibia* in two pigeons, towards the lower extremity, and introduced lengthways into the marrow two ſmall bits of wood covered with venom. Neither of the pigeons died, nor had any ſymptom of diſeaſe.

I cut the *tibia* in the ſame part, of two other pigeons, and introduced into the marrow two ſmall bits of wood, well covered with venom, keeping them there ſix minutes. Neither of the pigeons had any apparent ſymptom of the diſeaſe of the venom.

I repeated this experiment with the ſame circumſtances on four other pigeons. Each of the trials ended the ſame way, the pigeons all recovering within the ſame ſpace of time that two others did, which I employed as a compariſon, without venoming them.

The Venom applied to the Tranſparent Cornea.

I pierced the tranſparent cornea of the right eye of a large rabbit with a venomous tooth. The aqueous humour flowed out. I then, with another venomous tooth, firſt ſcratched, and afterwards pierced, the tranſparent cornea of the other eye. At the end of an hour I found the right eye filled with the aqueous humour, and perfectly ſound. At the

end

end of eighteen hours, a ſmall white ſpot formed in the tranſparent cornea of the other eye, but without any inflammation about it. At the end of three days, the white ſpot in the left eye was raiſed above the cornea.

I ſcratched the cornea of another rabbit with a tooth well dried, and at length pierced it. At the end of fourteen hours a dark ſpot appeared, and two days after the cornea was raiſed up in the form of a pearl.

I poured a drop of venom into the eye of a large rabbit, which I examined every hour. At the end of eighteen hours, the *membrana nictitans* ſeemed ſomewhat redder than uſual.

I poured two drops of venom into the eye of another rabbit, and this was not ſucceeded by any inflammation.

I made the ſame experiment on the eye of a third, which continued in its natural ſtate.

I repeated it on three other rabbits, neither of the eyes of which became ſenſibly inflamed.

I moiſtened the eyes of a large rabbit ſeveral times with a conſiderable quantity of venom, and likewiſe applied ſeveral drops to its lips and tongue. At the end of three hours the *membrana nictitans* appeared a little red, but at the end of eighteen hours returned to its natural ſtate.

I put ſeveral drops of venom on the tongue of another rabbit, and ſmeared it on the lips and palate with a bruſh. There was no ſwelling in any part

part of its mouth, neither was the rabbit at all diſordered.

This experiment repeated on two other rabbits was attended with the ſame conſequences. No part of the mouth was either ſwelled or inflamed.

CHAPTER V.

Experiments on the Comb, Gills, Noſe, and Neck, of Animals.

My next purſuit was that of examining the effects of the venom of the viper on the comb, gills, noſe, and neck of animals. My experiments on theſe parts have been attended with unexpected and intereſting conſequences; and therefore I have thought it proper to treat them apart, in an extenſive way.

Experiments on the Comb of Fowls.

I had the comb of a fowl bit twice by a viper. There was a conſiderable hemorrhage from the wounds made by the teeth. At the end of three hours the gills were ſwelled, and in ſix a large tumour or bladder was formed. The fowl died at the end of four days, without having either eat or drunk.

The

The tumour of the gills, which united them into one monſtrous body, was filled with a watry fleſh-coloured humour, and with an heap or web of filaments and veſſels.

I had a ſmall cock bit once in the comb by a viper, and treated it immediately. It died at the end of ten minutes.

I had another cock, of the ſame ſize, bit once in the comb by a viper, and treated it. At the end of two hours both gills had already ſwelled; at the end of twenty-two this ſwelling was very much abated; and in thirty-ſix there were only ſome little remains of ſwelling in one of them. In forty hours the cock was perfectly recovered.

I had the comb of a large cock bit three times by a viper. It was branched, pointed, and more than a third of an inch in thickneſs. It bled a little, and there were ſome ſmall drops of venom beſide the holes made by the teeth. I made a ſmall wound in the comb with the point of a lancet, and introduced a ſmall quantity of venom. The cock had no ſymptom of complaint. Two days after I had it bit twice in the comb by another viper. At the end of two hours the part appeared ſomewhat livid towards its baſis, and perhaps a little ſwelled. At the end of three hours the gills were very much enlarged, and at the expiration of twenty, were become of a monſtrous ſize, and livid for their whole extent. At the end of twenty-three hours they burſt, and the cock died very ſoon after.

There

There cannot be the ſmalleſt doubt but that the venom in the firſt caſe was thrown out by the blood, and this happens not unfrequently. It is much more difficult to account for the tumour that, notwithſtanding the cock was bit in the comb, formed in the gills. However I have frequently ſeen ſomething ſimilar happen in other animals. The bite made in the leg of a rabbit frequently cauſes a tumour, or an obſtruction of the humours in the moſt inferior parts of the ſame leg. But the experiments muſt be continued.

I had the comb of a fowl bit by two vipers, by each twice. At the end of two hours one of the gills only began to ſwell. In twenty they were both very much ſwelled, and united in ſuch a way that they formed a ſingle body. At the end of thirty-ſix they were enormouſly ſwelled and very livid. In ten days the fowl recovered. On the fourth day of the diſeaſe it reſpired with difficulty, and with a hiſſing noiſe; the glottis was inflamed and open, and the trachea arteria ſwelled.

I had the comb of another fowl bit ſeveral times by two vipers. At the end of three minutes the part next the head was livid, and appeared a little ſwelled. In an hour the livid colour and tumefaction ſeemed to be ſubſided, but on the other hand, the gills were enlarged. In three hours one of the lower eye-lids exuded blood from all its ſmall orifices. The gullet and palate were black. In twelve hours the fowl was in a dying ſtate, the gills being

livid

livid and of an enormous ſize. It died at the end of thirty-three hours.

I had the comb of a fowl bit ſeveral times by a viper. One of its gills ſwelled a little. At the end of thirty-ſix hours this ſmall degree of ſwelling had diſappeared, but the fowl reſpired with difficulty, and in doing this made a great noiſe. The wind-pipe was ſwelled, and very much enflamed, even at the end of ſix days. The animal was perfectly recovered in ten.

All theſe experiments ſhow that there is an immediate communication of veſſels and humours, betwixt the comb and gills of fowls. I do not give a detail of more than ten experiments beſides that I made on fowls, ſince they terminated in the way with the caſes juſt related.

Experiments on the Gills of Fowls.

I was deſirous of knowing what would be the conſequence of having fowls bit by vipers, not in the comb, but in the gills only; that is to ſay, whether the bite would be equally dangerous, and whether the tumour would fly up to the comb without forming in the gills, or would form both in the gills and comb.

I had the gills of a fowl bit repeatedly by two vipers. At the end of two minutes they had already ſwelled, and become livid. There was a great flux of humours in the eyes, which were

cloſed

cloſed by the enlargement of the membrana nictitans. In leſs than an hour the gills were of an enormous ſize, and livid all over. The fowl died at the end of the fifth hour.

I had the gills of a ſecond fowl bit twice by a viper. They ſwelled in leſs than four minutes, and in two hours were extremely large and livid. The comb appeared a little dark towards its points and edges. The fowl died at the end of three days.

Theſe trials may induce one to ſuſpect that wounds made in the comb are leſs dangerous than thoſe made with the ſame circumſtances in the gills.

To come at the truth of this hypotheſis, I made the following experiments. I had ſix fowls bit, each twice by a diſtinct viper; three in the comb, and three in the gills. One only of the former died, and two of the latter.

On repeating this experiment on ſix other fowls, the reſult was ſomewhat different. Only one of thoſe bit in the comb died, and all thoſe that were bit in the gills.

Theſe new experiments led me to think that my conjecture was very probable; that is to ſay, that the bite of the viper in fowls is more dangerous when made in the gills, than when it was made in the comb.

The accident which ſupervenes in the fowls the comb of which has been bit by vipers, is very ſingular. The action of the venom, or its diſeaſe, is conveyed

conveyed to a remote part that has not been bit; but when the gills are bit, the venom does not fly up to the comb, nor does the diſeaſe communicate itſelf to that part, and yet the ſtructure both of the gills and comb is the ſame, and the veſſels and nerves are common to both.

This circumſtance ſtruck me ſo forcibly, that I thought it deſerving an analyſis of ſome kind, and of being ſearched into by ſtill further trials.

I began by having the comb of a fowl bit once, and at the end of fifteen ſeconds cut off both gills. The fowl not only recovered, but there was no change in the gills, nor any appearance nor ſymptom whatever of the diſeaſe of the venom.

I had another fowl bit once in the comb, and at the end of fifteen ſeconds cut it entirely away. The gills did not ſwell, neither had the fowl any ſymptom of the diſeaſe of the venom.

I had the gills of a large cock bit repeatedly by a viper. In ſix hours they were both very much enlarged. On the following day they were ſtill more ſo, and were beſides livid. The cock recovered at the end of thirteen days.

I had the gills of another cock, a very large one, bit ſeveral times by two vipers. At the end of ten minutes I cut them off. On the following day it ate, and appeared in health, and after three days was perfectly recovered.

I repeated this experiment on the gills of ſix other cocks, each of which I had bit repeatedly

by two diſtinct vipers. I cut off the gills in all of them, but at different intervals; at the end of 1, 2, 4, 8, 16, 32, minutes. They all recovered, and had no other complaint, than that produced by the cutting off of the parts.

I had a large cock bit ſeveral times in the comb by two vipers, and after eight minutes cut off its gills. It died at the end of three hours.

I had another cock, a very large one, bit repeatedly in the comb by two vipers, and after four minutes cut off its gills. It died at the end of twenty-ſeven minutes. It was ſcarcely bit by the firſt viper, when it could no longer ſupport itſelf, or hold its head erect. It opened its beak, from which a glutinous humour flowed, and breathed ſhort, and with difficulty.

I repeated this experiment on ſix other cocks, each of which I had repeatedly bit in the comb by two diſtinct vipers. I cut off the gills in each of them at the end of four minutes. Three died in leſs than twenty hours; the other three were very much diſeaſed, and did not recover till the end of ten days.

Experiments on the Neck of Animals.

I had a ſmall guineapig bit twice by a viper in the back part of its neck. I treated it. It died at the end of forty minutes.

I had

I had a middle-ſized rabbit bit twice by a viper in the upper part of the neck, and treated it. It died at the end of twenty-four hours.

I had two guineapigs bit in the neck, each twice by a diſtinct viper. One was treated, and the other not. Both of them died; the one treated at the end of an hour, the other at the end of four.

I had two ſmall rabbits bit in the neck, each repeatedly by a diſtinct viper. I treated one, and made it ſwallow the volatile alkali ſeveral times, and did nothing to the other. Both of them died; the firſt at the end of four hours, and the other at the end of twenty-two.

I had a large guineapig bit twice in the neck by a viper. In an hour the part of the neck that had been bit became ſwelled and livid. At the end of twenty-three hours a large wound appeared. At the end of the ſecond day the humours which formed the tumour, had extended to beneath the chin, and formed a large bag or bladder. In four days the tumour had ſwelled to ſuch a degree, that it almoſt covered the breaſt. The ſkin had loſt its hair and epidermis, and a ſlightly-coloured humour exuded from it. At the end of ſix days the ſwelling began to diminiſh, and the guineapig recovered at the end of fifteen.

The diſeaſe in this animal, or the matter which deſcended from the upper to the lower part of its neck, and which even reached to the breaſt where it formed a cyſt or bladder, bears a ſtrong analogy to the circumſtances that were obſerved on having

the fowls bit. There is only this difference, that in fowls this effect is more frequent, and is oftener the caſe than otherwiſe; whilſt, on the contrary, it happens very rarely in quadrupeds bit in the neck, at leaſt in guineapigs. Of twenty-two animals bit in the ſame way, of which eleven were treated and eleven not, I found five in which this tumour deſcended below the neck, and formed a bladder. Of theſe five, three were treated, and two not. The number of deaths, which conſiſted of four in the whole, was equal on both ſides.

It is however certain, that having had ſome others bit, but each of them by ſeveral vipers, and ſeveral times by each, the tumour or cyſt formed in the interiour part, in a greater number of them, and that almoſt all of them died.

The conſequences were analogous, on trying the ſame experiments on rabbits. The cyſt ſometimes forms beneath the chin of theſe animals, although they have only been bit in the neck; and this happens more frequently when they have been bit by ſeveral vipers, in which caſe they die much readier.

Experiments on the Noſe of Animals.

It remained for me to examine the bite of the viper on a part that is held the moſt ſenſible, and the moſt likely to occaſion death when it receives an injury, in ſome particular animals.—This is the

noſe.

noſe. It appears, that the cat, an animal very obſtinate in dying, periſhes as readily as others, if ſtruck in this part.

Mead reckoned it ſo ſenſible and ſo dangerous in dogs, that, wiſhing to aſſure himſelf of the efficacy of a remedy againſt the bite of the viper, he had a dog bit on the noſe, and applied the remedy. The dog lived, and this was ſufficient to give the remedy the reputation of a true ſpecifick; ſo ſtrong was the opinion, that a bite of the viper on the noſe was mortal.

I ſhall not relate here all the experiments I made on this part, but only a ſmall number of them, which will be ſufficient to give a clear idea of the fallacy of ſome popular opinions.

We ſhall ſee what we ought to think of the bite of the viper on the noſe, and how abſolutely neceſſary it is to conſult nature by facts and experiments. Nothing is more dangerous and more uncertain in reſearches of this nature, than a vague analogy, or a ſeducing and probable reaſoning. Nature is not to be divined, and *prophets* in the ſcience of phyſicks are not to be believed.

I had a ſmall rabbit bit twice on the noſe by a viper. In two minutes the part was ſenſibly inflamed. In three hours a tumour was formed in the neck, beneath the chin. In ſeven hours this tumour was become very large.—The animal recovered however.

I had another rabbit, ſomewhat larger than the former one, bit on the noſe by a viper, and treated

it. It was bit twice, but one of the bites was made on the upper lip, at the ſide of the noſe. In the ſpace of two minutes the noſe was ſwelled, and a very large tumour formed under the chin. At the end of twenty hours this tumour burſt, and diſcharged a great deal of matter. The rabbit recovered at the end of ſix days.

I had a third rabbit, of a middle ſize, bit twice on the noſe by a viper. In a very little time the part ſwelled and inflamed. In two hours a tumour formed beneath the chin, which at the end of ſeven diſcharged blood, and was very large. In thirty-ſix hours the tumour and ſkin began to dry, and the animal recovered at the end of the ſixth day.

Six other rabbits were bit in the ſame way. Neither of them died, and the effects of the bites were pretty much the ſame as thoſe related above. The bite of the viper on the noſe of rabbits, contrary to what one would naturally have thought, ſeems to be leſs dangerous than that in other parts. The diſeaſe it produces, as to the ſeat of it, is very ſimilar to that in the comb of fowls. Here again a tumour forms, in a part where the animal has not been bit, and beneath the ſeat of the bite, in which, in moſt caſes, the venom ſcarcely occaſions a real and ſenſible complaint. The only eſſential difference is, that the tumour in rabbits is of a greater extent, reaching ſometimes to the middle of the breaſt.

We are now to ſee, whether the ſame thing happens in animals of other ſpecies.

I had.

I had a large guineapig bit on the noſe by a viper. In two hours the part was very much ſwelled; at the end of three the ſwelling was diminiſhed, but in its place a large tumour formed beneath the chin. At the end of fifteen hours, the tumour broke, and diſcharged a great deal of blood and ſerum. In thirty-ſix hours the diſcharge ceaſed, and the opening in the ſkin dried up. The animal was perfectly recovered at the end of four days. It was never very much diſordered, ſince it ate during the whole time of its illneſs.

I had another large guineapig bit twice on the noſe by a viper. The noſe and mouth ſwelled very much, but this ſwelling diminiſhed in proportion as a tumour formed under the chin. After twenty-two hours the tumour, which had broke an hour before, began to dry up, and at the end of thirty-ſix ſeemed perfectly dry. At the end of two days the animal was recovered. During the whole courſe of its complaint it ſuffered but little, and ate conſtantly.

I had a large guineapig bit on the mouth by two vipers, each biting twice. The noſe ſwelled in leſs than three minutes, and ſtill more ſo at the end of ten. Two hours after, a tumour formed beneath the chin, when the ſwelling of the noſe diminiſhed, and in a ſhort time entirely ſubſided. At the end of twenty-three hours the tumour beneath the chin was ſo large as almoſt to cover the breaſt, and in two hours more it burſt. In the ſpace of five other hours the animal recovered.

I repeated this experiment on another large guineapig, which I had bit by three vipers, each viper biting three times. The noſe and mouth both ſwelled very much, but continued in that ſtate only four hours. At the end of the ſecond hour a large tumour appeared under the chin, which in twenty-three hours was become enormous, and reached to the breaſt. This tumour broke at the end of thirty hours, but the animal was not perfectly recovered till the eighth day. The bones of the noſe were laid bare, and the ſurrounding ſkin all conſumed.

I made the ſame experiment on two other guineapigs, but ſmall ones. One died at the end of twelve hours; the other had the uſual tumour, was exceedingly ill, but did not die.

The bite of the viper on the noſe ſeems to produce pretty much the ſame effects on guineapigs as on rabbits; and it appears that the venom is leſs dangerous in this part than in any other. The ſame effects are here conſtantly obſerved as to the ſeat of the diſeaſe; but are they the ſame in all other animals?—I ſhall relate what I met with in dogs and cats, creatures that enter into the plan of my preſent reſearches, and this will ſhow how little analogy alone ought to be truſted, and that the ſame cauſe produces very different effects, on ſimply changing ſome circumſtance, which one would not have ſuppoſed capable of influencing a great deal.

I had a ſmall dog bit repeatedly on the noſe by two vipers. Both noſe and mouth ſwelled, and the

dog

dog died at the end of eight hours, without any ſymptom of diſeaſe in any other part.

I had another dog, twice as large as the preceding one, bit repeatedly on the noſe by two vipers. Its mouth ſwelled to ſuch a degree, that the lips were very much enlarged twelve hours after. It vomited ſeveral times, and continued ill for three days, when it began to drink. On the fourth it ate, and on the fifth was perfectly recovered.

I had another dog, ſtill larger than the one laſt mentioned, bit on the noſe by three vipers, each viper making three bites. In a little time its noſe, mouth, and lips, were ſwelled ſo as to become hideous. It vomited a great many times, ate and drank on the fourth day, and recovered on the fifth.

I had another dog, of the ſame ſize as the preceding one, bit on the noſe by four vipers, each biting three or four times. It had a bite at the ſide of its noſe, and another on one of its lips. It vomited frequently, neither ate nor drink till after the third day, and recovered on the fifth.

I had another large dog bit on the noſe by ſix vipers, each of which bit three or four times. Its noſe and mouth ſwelled enormouſly; it vomited a great many times, ate after the fourth day, and recovered on the ſixth.

Laſtly, I had another dog, of the ſame ſize as the three preceding ones, bit on the noſe by ſix vipers, by each three or four times. The part ſwelled vio-

lently, and the animal did not eat till after the fifth day. It vomited frequently, and recovered at the end of ſeven days.

Rabbits and guineapigs bit on the noſe, uſually have the diſeaſe beneath the chin, and not in the part bit. It is quite the contrary with dogs, in which the diſeaſe is entirely confined to the noſe, the part that received the bite. They therefore form an exception to the caſes related prior to theirs.

It is likewiſe ſingular, that as the action of the venom is confined to the noſe, it does not produce incurable wounds and gangrenes in that part. We, however, find it to be quite otherwiſe;—bites on the noſe in dogs are very rarely attended with a wound in the part, and the animal not only makes a ſtrong reſiſtance to the diſeaſe, but the latter at the ſame time appears to be very ſlight, ſince the recovery takes place in a few days.

Experiments on Cats bit on the Noſe.

We have ſeen above, that the cat makes the ſtrongeſt reſiſtance of any animal to the bite of the viper, although the venom conſtantly produces in it a diſeaſe. We may therefore conjecture, that this bite on the noſe of cats will not be productive of mortal effects. But we know, on the other hand, that mechanical percuſſions on the noſe are dangerous

dangerous to theſe animals, and that they ſoon die, if they fall from a height on this part.

From theſe conſiderations, I wiſhed here again to have recourſe to experiments, which can alone determine the truth.

I had a middle-ſized cat bit repeatedly on the noſe by a viper. Its mouth ſwelled for a conſiderable extent. It ate on the ſecond day, and recovered on the third.

I had another, of the ſame ſize, bit repeatedly on the noſe by a viper. A few minutes after, the part ſwelled. The cat vomited twice, ate on the ſecond day, and was perfectly recovered on the third.

In this ſecond cat, the diſeaſe of the venom was ſo very ſlight, that the animal appeared to ſuffer but little during its continuance.

I had a third cat bit repeatedly on the mouth by a viper. One of the bites fell on its upper lip, which bled a good deal, and the whole of its mouth ſwelled very violently; however it ate on the ſecond day, and on the third was recovered.

I had a large cat bit repeatedly on the noſe by a viper. The part bled very much, and ſwelled a few minutes after. At the end of twenty hours it was ſtill more ſwelled, notwithſtanding which, the cat appeared but little diſordered. It recovered at the end of forty hours.

I had another cat, of a middle ſize, bit repeatedly on the noſe by a viper, which likewiſe bit it on the mouth and lips. The mouth ſwelled at the end of five minutes, and at the end of five hours the

the cat vomited ſeveral times. In thirty-ſix hours it was perfectly recovered.

I had another cat, of a middle ſize, bit on the noſe, and on the mouth both below and above. After ſeven hours it vomited ſeveral times. Its noſe and mouth were but little ſwelled, and at the end of twenty hours it recovered.

Another cat, of a middle ſize, was bit by three vipers, each of which bit three times, or upwards, on the noſe, mouth, and even within the palate, from which there was an hemorrhage. Some minutes after, its mouth ſwelled, it vomited ſeveral times, but the palate did not ſwell at all. It ate at the end of three days, and at the end of the fifth was perfectly recovered.

I had another cat, ſomewhat larger than the preceding one, bit by four vipers. Each viper bit ſeveral times, on the noſe, mouth, and lips, and in the palate, inſomuch that the cat, feeling one of the bites within its mouth very ſenſibly, ſeized the viper betwixt its teeth, and almoſt ſevered its head from its body. The noſe and mouth in this cat ſwelled very much, it vomited ſeveral times, ate on the fourth day, and recovered on the ſixth.

I repeated theſe experiments on three other cats, which I had bit repeatedly in the noſe by a viper, and the effects were pretty much the ſame. We may therefore, I think, conclude, that the bite of the viper on the noſe is not very dangerous to dogs, and that it is ſtill leſs ſo to cats.

At

It is, however, very ſtrange, that in both theſe animals there is no tumour beneath the chin, and that the local diſeaſe is confined to the part bit; whilſt, on the contrary, the diſeaſe in rabbits and guineapigs is not in the part bit, but in another part of the animal beneath it.

It is clear, that this difference can only depend on the different organization and nature of theſe animals; and it is preciſely this diverſity that we are ignorant of.

I muſt here obviate a difficulty that may be made by thoſe who are not accuſtomed to ſuch experiments.

Theſe may oppoſe, that bites in the noſe probably become leſs dangerous from the animal's licking the part. This is never done by rabbits and guineapigs, notwithſtanding they are bit. I have aſſured myſelf of this particular in ſuch a way, that I have not the ſmalleſt ſuſpicion of having been deceived.

More than two-thirds of both dogs and cats that I had bit in the noſe, never licked the part, although they could eaſily have done it. I obſerved them myſelf, and had them obſerved, for whole hours. It is true, that thoſe which bled a good deal licked themſelves if they could; but it was evident, on obſerving them, that they only endeavoured with the tongue, to free themſelves from the blood which tickled them in flowing down, and that this was no ſooner effected, which happens in a moment, than they ceaſed to do ſo.—In the

expe-

experiments I made on dogs and cats, I prevented ſome of them, when they bled at the noſe, from licking the part, and ſuffered others to do it. The diſeaſe was the ſame in all; and it is, therefore, certain, that ſimply licking the noſe, whether in dog or cat, does not at all diminiſh the effects of the venom of the viper on that part.

CHAPTER VI.

Experiments on the Tendons.

SEVERAL modern phyſiologiſts have thought that the tendons are not endued with ſenſation. It is certain that it has not yet been proved clearly, that a tendon receives nerves either from the muſcle, or from the tunica vaginalis which covers it. Neither is it more apparent that it has blood-veſſels, at leaſt in any number, and ſenſible ones. It is therefore natural to ſuſpect, that the bite of the viper on a tendon cannot be of any great conſequence, and that the venom cannot act on this part. I wiſhed nevertheleſs to conſult experiment once more on this point.

In having the tendons bit by vipers, I was more than once on the point of being deceived; and if I had

had not multiplied my experiments, and varied them in ſeveral ways, as I did, I ſhould certainly have been ſo. I ſhall be circumſtantial in relating ſome of the trials I made on the tendons, to ſhow that it is eaſy for any one, even for an obſerver, to be deceived, if he only follows ſimple experiments, ſince the reſult of them may vary, although there appears to be no variety in the circumſtances with which they are made.

My experiments were made on rabbits, of which I employed the largeſt I could find, ſome of them weighing ten pounds and upwards.

Having removed the ſkin from the *tendo achillis* of a rabbit, and perfectly ſtripped it of its tunick, for a ſpace of ſix lines in length, I paſſed under it ſeveral folds of fine linen, to prevent the venom from communicating to any other part. I wounded the tendon in ſeveral places with a venomous tooth, and afterwards covered it with bits of linen in ſuch a way, that it did not ſeem poſſible for the poiſon to communicate to the neighbouring parts. The rabbit died at the end of thirty-ſix hours. The tendon was livid throughout its whole ſubſtance, but the parts about it were not ſenſibly changed.

I opened by an inciſion, the ſkin that covers the *tendines achillis* of another rabbit, and ſtripped the tunick from both. The tendons were ſmooth, ſilver-coloured, and free from veſſels. I paſſed ſeveral folds of linen beneath them, and had them bit ſeveral times by two vipers, covering them with linen in ſuch a way, that the venom could not glide elſe-

where.

where. The rabbit died at the end of thirty-eight hours. The blood in the auricles, in the ventricles, and in the large veſſels of the lungs, was black and coagulated. There were ſeveral livid ſpots in the lungs. The muſcles adjacent to the tendons were a little inflamed, and likewiſe had livid ſpots in ſeveral places.

I repeated this experiment on two other rabbits, with pretty much the ſame reſult. Both of them were dead at the end of thirty-ſeven hours.

Although it clearly reſults from the experiments I have juſt related, that rabbits die after having been bit in the *tendo achillis* by vipers, I could not however conceive, that the death of thoſe I have mentioned, was occaſioned by the introduction of the venom, and its ſubſequent diſeaſe.

It did not appear poſſible to me, that a part endued with ſo few vital principles as the tendon, which is not at all ſenſible, and which may be cut both in men and animals with impunity, could be ſuſceptible of the action of a venom that has no influence either on the mouth or ſtomach. I ſuſpected that theſe animals died from ſome other cauſe or circumſtance I could not diſcover.

In conſequence of this ſuſpicion I determined to multiply my experiments, and to diverſify them as the caſe might require.

Having removed the ſkin from the *tendo achillis* of a rabbit, and ſtripped it of its tunick above and below, ſo that it appeared ſmooth and white, I wounded it with the point of a large and ſharp needle,

dle, which pierced it through. The needle was well covered with venom, and I had put ſeveral folds of linen beneath the tendon. I wiped the tendon ſeveral times, removed the linen, and left the part expoſed. I then introduced into the aperture made in the tendon, a bit of wood well covered with venom, and having withdrawn it, poured in a drop of pure venom. At the end of twenty-four hours, the tendon ſeemed diſcoloured at the part wounded. The rabbit, however, ate conſtantly, and at the end of fifteen days was recovered.

In another rabbit, I removed a large portion of the ſkin that covers the joint of the knee, and ſtripped the ligament that binds this part of the adipoſe membrane. I wounded it obliquely with a venomous tooth, in eight places, at each of which a ſmall drop of venom appeared. I made ſmall inciſions into each puncture, with the point of a lancet, which penetrated into the ſubſtance of the ligament without piercing it through, and conveyed the venom within. The rabbit recovered in eight days, and ſeemed not to have had any internal complaint. It ate conſtantly, and continued lively and active.

Having ſtripped the *tendo achillis* of another rabbit of its tunick, and put folds of linen beneath it as uſual, I had it bit ſeveral times by two vipers. I then covered it with linen, and removed that which was beneath. The rabbit for the firſt few days ſeemed to have no complaint, but the wound in the tendon never healed perfectly. At the end of ten

days

days its belly appeared to ſwell, and, on its dying at the end of ten days more, I found it to be dropſical.

Theſe experiments ſeem to oppoſe the former ones, and to render it a matter of doubt whether the bite of the viper on a tendon produces the diſeaſe of the venom, or not. The latter caſes ſeem to indicate that it does not; but they are contradicted by the former ones. Now, as one of the principal reſearches I propoſed to myſelf to make, at ſetting out on my experiments, was to diſcover what are the parts acted on by the venom of the viper, and to obſerve its different effects on the different parts of an animal, I was determined to continue my experiments on the tendons with a degree of obſtinacy, and to ſee if I could ſucceed in clearing up this point.

Wiſhing to obſerve a greater degree of preciſion in my experiments, and ſuſpecting that the venom might perhaps communicate to the neighbouring parts in which the inciſion had been made, and that it might penetrate by degrees through the linen, how much ſoever the latter might have been folded, I conceived the idea of putting betwixt the folds, a piece of thin and pliable lead.

Having ſtripped the *tendo achillis* of a rabbit of its tunick, I paſſed beneath it a piece of linen folded eight times, in the middle of which I had put a bit of lead, ſuch as I have juſt deſcribed. I pricked the tendon in ſeveral places with two venomous teeth, and covered it in ſuch a way that it was quite

encloſed, having a bit of lead both above and below. The animal died at the end of thirty-two hours. The tendon was black at the parts where it had been wounded, the muſcles near it were a little inflamed, and the blood about the heart in a diſſolved ſtate.

All theſe precautions, as we ſee, could not prevent or retard the death of the animal. As this was, however, but one ſolitary caſe, I did not think it proper to ſtop here.

I repeated the experiment on the *tendines achillis* of four other rabbits, well ſtripped of their tunicks. I applied the linen and bits of lead as uſual, wounding the tendons with venomous teeth, that the venom might be more collected, and ſpread as little as poſſible on the tendon. In a word, I omitted nothing that could make theſe experiments deciſive ones. The rabbits all died in leſs than forty hours. In ſome of them the blood was coagulated about the heart, but not in others. The lungs were ſpotted in all of them. The muſcles in the vicinity of the tendons were a little inflamed, and in two of the rabbits livid.

Theſe new trials did not clear up my doubts. If on one hand they rendered the action of the venom on the tendons probable, on the other hand I could not conceive how a part, that is neither ſenſible, nervous, vaſcular, nor muſcular, could either receive the diſeaſe of the venom, or communicate it to the animal, ſo as to occaſion its death. I reflected again, that I had employed large rabbits; that I had neither applied much venom, nor made uſe of

many vipers; and that, on other occaſions, I had found a large rabbit to die late and with difficulty, although it had been bit by ſeveral vipers, and died with large wounds, and with the moſt aſſured ſymptoms of the diſeaſe of the venom. This made me fall upon a new ſpecies of experiments, from which I flattered myſelf that I ſhould draw ſome kind of information.

I prepared the *tendo achillis* of a rabbit as above, and paſſed beneath it a piece of linen folded ſixteen times, with a bit of lead in the middle. I pierced the tendon in the uſual part with a venomous tooth, and introduced a drop of venom collected at the orifice, into the ſubſtance of the tendon, by a longitudinal inciſion three lines in length, made with the point of a penknife. The inciſion did not penetrate through. I left the tendon, venomed in this way, during a ſpace of ſix or ſeven minutes, and then ſoaked up the venom with dry lint, and by the means of ſmall pincers, waſhed the wounded part of the tendon ſeveral ſucceſſive times. In proportion as the linen became moiſt, I took hold of one end of it, and drew it by degrees from under the tendon. It was impoſſible in this way for the water to ſoak through the linen, and communicate the venom to the adjacent parts. As I waſhed the tendon upwards of twenty times, it was not poſſible for an atom of venom to remain within it. The rabbit died at the end of thirty-two hours; the tendon was almoſt in its natural ſtate, its colour being ſcarcely deepened at the part where the wound was made.

I re-

I repeated this experiment on two other rabbits, uſing the ſame precautions. They were both dead in leſs than thirty-ſeven hours.

It occurred to me, that the linen left above and beneath the tendon till the death of the animal, might perhaps bring on ſuch a change in the neighbouring parts, as to occaſion a mortal diſeaſe.

Having removed the ſkin from the *tendo achillis* of a rabbit, and ſtripped it of its tunick, I put linen beneath it as uſual, and wounded it with a venomous tooth. I wiped the tendon with lint, and waſhed it a little, taking care that the water did not touch the adjacent parts. I then removed the linen, and applied freſh, to the part. The rabbit died at the end of thirty-ſix hours. The parts about the tendon were in a natural ſtate.

I prepared the Achilles' tendons of another rabbit in the uſual way, and wounded them with a venomous tooth. I left them in this ſtate for two minutes, and then threw on them a great deal of water, repeating the ablution till I conceived that they were perfectly cleanſed in every part, and that the venom was either totally waſhed away, or diluted ſo effectually, that it could not convey its action to the adjacent parts. I had found from former experiments, that when any other part of an animal was bit, or wounded by a venomous tooth, the throwing of any quantity of water on it was ineffectual, and did not prevent the animal from dying, and from having the uſual diſeaſe of the venom in

the part bit. The rabbit, the ſubject of this experiment, died at the end of thirty-two hours.

Another rabbit treated in the ſame way, not only recovered, but ſeemed to have no other complaint than that occaſioned by the inciſion of the ſkin, and other parts that cover the tendon.

Theſe reſpective caſes, conſidered circumſtantially, began to perſuade me, that the venom of the viper is perfectly innocent to a tendon. To be certain of this, I thought of varying my experiments ſtill more, and of making them in ſuch a way, that they ſhould at length become deciſive.

Having removed the ſkin, and laid bare the *tendo achillis* of a rabbit, I bound it very tight with a piece of packthread, at both extremities of the tendinous ſubſtance. The ligatures were made in ſuch a way, that it was not poſſible for any communication either of humours or ſenſation betwixt the tendon and the animal to take place. I put the uſual folded linen under the tendon, which I wounded in ſeveral places with a venomous tooth, betwixt the two ligatures. I covered the tendon with linen, and the rabbit died at the end of thirty-two hours.

I repeated this experiment on another rabbit, the tendons of which I tied in the way above, and had it bit betwixt the two ligatures. I waſhed the wounds with a great deal of water, which I threw on with force, and then removed the linen from beneath. This rabbit died at the end of thirty hours.

Another rabbit died in twenty-ſeven hours, after having been treated pretty much in the ſame way,

with

with the preceding one, with only this difference, that inſtead of throwing a great deal of water on the tendons, I waſhed them by degrees, applying clean and dry linen, after removing that which I employed at firſt.

It ſeems at length pretty clear, that the venom of the viper is not the cauſe of the death of the rabbits, in the caſes in queſtion, and that it has no action on the tendons. A doubt ſtill remained, however, which it was neceſſary to clear up. I had obſerved that ſeveral muſcular fibres had found their way into the tendinous portions that form the *tendo achillis*, and conceived that the venom of the viper might firſt communicate itſelf to them, and from them to the other parts of the animal. Notwithſtanding there was but little probability in this conjecture, I wiſhed to inform myſelf on the ſubject by experiment.

Having removed a portion of the ſkin from the *tendo achillis*, and ſtripped it of its tunick, I deſtroyed the muſcular fibres that deſcend from the crural muſcles, and implant themſelves betwixt the three portions of this tendon. I paſſed ſeveral doubles of linen betwixt theſe tendinous portions, in ſuch a way that one of them was ſeparated from the other two, and encloſed in the linen. I wounded this portion with a venomous tooth, and covered it ſo as to prevent the venom from touching any of the adjacent parts. The rabbit died at the end of thirty-two hours, with its heart and veſſels filled with black and clotted blood.

I repeated this experiment on the tendons of another rabbit, which died at the end of thirty-two hours. The wounded portions of the tendons were dark throughout their whole ſubſtance, and thoſe which had not been wounded were ſtill much more ſo. The lungs were covered with livid ſpots, and the heart and large veſſels filled with black and clotted blood.

I made an experiment on another rabbit, in which, after deſtroying the fibres betwixt the portions of the tendon, I paſſed a folded linen under its whole ſubſtance, as I had done in the caſes related a little above, and wounded it, without ſeparating the parts, with a venomous tooth. The rabbit died at the end of thirty-three hours. The wounded tendon was become darker and redder in ſome places, and the blood in the heart, and in the veſſels that go out from it, was black, but fluid.

It appears ſtill more, that the venom of the viper is not the cauſe of the death of theſe animals, but that it depends on another cauſe, probably on the denudation of the tendon itſelf. The following experiments remove all doubts on the ſubject.

I got ready ſix very large rabbits, all alike in ſize, in two of which I laid the Achilles' tendons bare, as uſual, and wounded them with a venomous tooth, after having incloſed them well in linen. In two others I laid the tendons likewiſe bare, but pricked them with a needle in ſeveral places. In two others I ſimply laid them bare, without wounding or pricking them.

I

I covered all the tendons in the ſame way with linen. All the rabbits died; the two that had been venomed, died together at the end of thirty-two hours; of the two the tendons of which were pricked with a needle, one died in thirty hours, the other in thirty-two. The two in which the tendons were ſimply laid bare, died, one in twenty-ſeven hours, the other in forty.

The inferences I have drawn from the experiments on the tendons hitherto related, are as follows:

I. That a tendon is not ſuſceptible of the diſeaſe of the venom.

II. That when a tendon is ſtripped of its tunick, the animal almoſt invariably dies, without any intervention of venom.

This laſt inference is a very important one, and may be of ſome uſe in the punctures of tendons in man. It ſhows how dangerous it is to ſtrip theſe parts of the *tunica vaginalis*, and how much this membrane ought to be ſpared.

It remained for me to make one other experiment on a tendon, which I ſhall relate here, and which may throw ſome light on the nature and economy of tendinous ſubſtances, and of their nutrition. Having laid the *tendo achillis* of a rabbit perfectly bare, and deſtroyed the muſcular fibres that enter into it, ſo that there could be no longer any fleſhy fibres or veſſels in the tendon, I found that the rabbit ate a few hours after, and conjectured that it would

would therefore probably recover. In effect it lived, and at the end of thirty-four days recovered perfectly, the wound made in its ſkin healing up. I wiſhed to ſee what had happened to the tendon, and whether, as one would have ſuppoſed, it had dried up from a want of veſſels. All the veſſels about the tendon had been cut, and it was abſolutely ſeparated from every other ſubſtance throughout, except at its two extremities. I found it covered by a ſubſtance, partly ſpongy or cellular and partly callous, and ſprinkled with ſeveral veſſels. When I got to the tendon, I found it whitiſh, ſupple, and nouriſhed, as uſual, although it did not any where appear to receive veſſels.

Were repeated experiments ſimilar to this one to be made, important conſequences, and facts relative to the nutrition of certain parts, would perhaps reſult from them.

The multiplied and varied experiments I made on the tendons, have been of very great uſe to me in the purſuit of my reſearches. If any doubt had remained on the ſubject; if I had not aſſured myſelf to a certainty that the bite of the viper on this part is not attended with any conſequence; if I had apprehended that the venom could communicate itſelf to the animal through the medium of this ſubſtance; I ſhould have had a thouſand doubts as to the parts on which the venom acts, in an animal that has been bit. No ſubject in nature is abſolutely indifferent; and when ſuch rare and extraordinary effects are to be examined in the animal

mal body, nothing is to be neglected—nothing is to be deemed unneceſſary.

CHAPTER VII.

On the Nature of the Venom of the Viper. Deſcription of certain Parts of the Head of the Viper, that relate to the Venom.

BEFORE I examine the properties and nature of the venom of the viper, I think it incumbent on me to ſpeak of ſome other particulars, that relate to the canine teeth of this animal, to the bag or membrane with which they are naturally covered, and to the veſicle or receptacle of the venom, which the moſt modern writers continue to confound with the bag or ſheath of the teeth. I have treated of all theſe particulars in the firſt part of this work, but think it eſſentially neceſſary to introduce ſome figures here, which will give a juſter conception of what I have ſaid in the part alluded to, and of what I ſhall ſay in the ſequel.

I have judged it expedient to devote a chapter entirely to this ſubject, and to interrupt, as it were, the chain of my experiments on the effect of this poiſon, applied to the different parts of animals; ſince it is before all neceſſary, that the reader ſhould know the nature of the venom, and not be left any longer to bewilder himſelf in the erroneous opinions, and hypotheſes deſtitute of foundation, that have been ſpread by the writers who have employ-

ed

ed themſelves on the occaſion, both before and after the publication of my firſt experiments. Too much cannot be ſaid for this effect; for unfortunately, when the mind is prejudiced in favour of any opinion whatever, eſtabliſhed by authority, and generally adopted, it ſeems to deny itſelf to even the evidence of fact.

In Mead's work on Poiſons, a deſcription is found of the head of the viper, the parts of which are repreſented by figures. Theſe figures of Mead, or rather of Nicholls, who is the real author of them, are ſo imperfect, that I have been obliged to ſubſtitute others I have had purpoſely made. I have found the former ones out of all truth and nature, and whoever will take the trouble to confront them with the parts from which they were drawn, will find no difficulty in agreeing with me.

Fig. 1. of plate I. of this work (ſee the concluſion of the ſecond volume) repreſents the two canine teeth of the viper of one ſide of the upper jaw, partly covered by a membrane in the form of a bag or ſheath, open, as it is ſeen, to give paſſage to the teeth. Mead pourtrays this bag as if it was fringed at its edges. It is indeed ſometimes found in this ſtate, but is oftener without fringe or indentation, and ſuch as I have repreſented it. The canine teeth are elevated and laid a little bare, as they appear when the viper is on the point of biting; when it depreſſes them, they enter entirely into the bag or ſheath. It is eaſy to ſee, that if this bag were the receptacle of the venom, the latter would naturally flow out at the opening in it, and would

paſs

paſs continually into the mouth of the viper. This errour is copied from Redi, who believed that the venom was contained in this ſheath that covers the teeth, and that it was ſecreted in a ſmall gland ſeated under the eye.

Fig. 2. repreſents this bag or ſheath, *s s* opened with ſciſſars as far as its baſis, and likewiſe on the bone of the upper jaw. An elliptical hole *n. e.* with rounded edges, is ſeen at the baſis of each of the canine teeth, and a longer and narrower hole, towards the point of each tooth, *r. a.*

At the ſide of the teeth a bladder is found, *m.* reſembling a ſhepherd's purſe, which pierces the ſheath by a long canal ending in a ſmall orifice *o* betwixt the two teeth. The venom contained in the purſe or bladder paſſes through this canal, and conveys itſelf to the tooth, entering at the hole ſituated at its baſis, and going out by that at its point.

Fig. 3. repreſents the bladder or purſe ſeen with a lens. It is not formed of a ſmooth even membrane, but is on the contrary full of plaits, as if it was a compages of inteſtines, or of wrinkles and ridges. It is of a triangular ſhape, and has a much greater width than depth. If it is cut tranſverſely, and examined with attention, it is found to be of a ſpongy ſubſtance, and compoſed of cells deeper than they are broad. Every thing concurs to the belief that it is not a ſimple bladder or receptacle of venom, but rather a true gland, very voluminous and of a particular ſtructure, which ſeparates the venom from

from the blood of the viper, and in which it is reſerved for the purpoſes it is deſtined to by nature, undoubtedly for the animal's advantage.

The cellular ſtructure of this ſingular gland does not permit the viper to expreſs with facility all the venom it contains. I have found a difficulty in forcing it out by a very ſtrong preſſure on the gland with my fingers; and indeed we have ſeen, that a viper is capable of killing ſix or ſeven pigeons, one after the other.

The two figures, 4, 4, repreſent the receptacle of venom in its natural ſize, ſeen at its anteriour, and at its poſterior part, and united with its excretory canal.

Fig. 5, ſhows a tranſverſe ſection of the above, ſeparated by many ſmall partitions, *s c*, &c. and filled with the venom, which flows out drop by drop, as at *r a*, &c. It appears in this way when obſerved with a lens.

Fig. 6, repreſents a canine tooth of a viper, with all its internal cavities, and its two external openings.

s s, are the elliptical hole at the point of the tooth.

c a, the opening of the hole at its baſis.

i i i, are the internal canal of the tooth, which opens at the baſis *c a*, and at the point *s s*.

There is a large opening, *e*, which forms the baſis of the tooth, and the ſection of which is repreſented by *m*.

r o

r o, of the figure at the ſide, are the two openings, *i e*, of figure 6, which are diſcovered by a ſection of the tooth, as at *a b*.

r, repreſents the ſhape of the longitudinal hole of the tooth.

o, repreſents the opening of the hole at the baſis. This ſecond canal of the tooth does not communicate with the firſt, and only extends as far as *r*.

Fig. 7, repreſents two canine teeth on one ſide, having at their baſis ſeveral other teeth, more or leſs formed, *a c r*. Theſe teeth are moſt frequently ſix in number, and are ſituated in the ſheath, and covered with a very fine cellular web, which binds them, and unites them together. They are placed one over the other, thoſe that are uppermoſt, or neareſt the canine teeth, being the largeſt. The others decreaſe in proportion, and the two that are neareſt the canine teeth are perfectly alike in ſize. The points of all them, even of the ſmalleſt, are pretty hard, and well formed;—they are channelled, and end by the uſual hole at the point.

When theſe teeth are ſeven in number, the ſeventh is always the ſmalleſt of the whole. It is ſituated below all the others, and in the middle. The baſis of theſe teeth is not yet formed, and merely conſiſts of a flexible, tranſparent, and whitiſh jelly. They are not only deficient at their baſis, but likewiſe want the oval hole;—the principles of it are ſometimes ſeen in the largeſt of them.

Although the matter at the baſis of theſe teeth appears a ſimple jelly, even when it is viewed with the common lens, the naturaliſt would be very much

much miſtaken, if he ſuppoſed it to be non-organical. Stronger lens than the ordinary ones have ſhown me, that it is compoſed of a very fine webbed membrane, filled with extremely ſmall round corpuſcles. This membrane folds over itſelf, and ſeems to ſhow, even the holes, and the form that the baſis of the tooth is one day to take. I have, however, ſometimes thought I could diſtinguiſh this. Be that as it may, it is certain, that the gelatinous part of the tooth is organized, and that it exiſts in a ſtate of organization a long time before the tooth is entirely formed and in a perfect ſtate.

Of the Nature of the Venom of the Viper. It is examined as to its Acidity.

The acquiring a perfect knowledge of the nature of the viper's venom may be of the greateſt importance to animal phyſicks, and, at the ſame time, very uſeful to the human ſpecies. Vague and ſuperficial notions on this point, have given birth to hypotheſes, to theories, and laſtly, to remedies.

The volatile alkali in a great meaſure owes its reputation, to the opinion of the venom of the viper being acid.

The ancients were ignorant of what it conſiſted in, and of the part of the animal in which it reſided. François Redi was the firſt to eſtabliſh theſe points. He found it to be a humour ſimilar to the oil of ſweet almonds, which the viper conveys with its tooth

tooth into the wound it makes in biting. But he was miſtaken in almoſt all he ſaid beſides on the ſubject of this venom. He believed that it reſided in the bag, or plaited membrane, that covers the canine teeth. He could never diſcover that it entered into the tooth itſelf, and flowed out of it; and thought that the ſmall gland ſeated under the eye of the viper, ſerved to ſecrete this humour, into the nature of which I do not find that he ever made any reſearch.

Before the time of Redi, there were none but very vague and confuſed ideas on the venom of the viper. We owe to this celebrated Italian naturaliſt the firſt advances into a ſubject, which he found in its infant ſtate, filled with hypotheſes and vulgar errours. Theſe errours were proper to the time he lived in, and it required a genius like his to combat them, and to open a new road to truth. It ſeems as if we only throw off our ignorance to plunge ourſelves into errour, and that it is at this criſis that the man of genius gives us ſome glimmerings of light. We ſet out at ignorance, which leads us to errour, and from errour we at length arrive at truth. This is the uſual progreſs of human intelligence, and through theſe gradations the moſt enlightened nations have paſſed.

Mead is the firſt who in any way examined the nature and qualities of the venom of the viper; but from a fatality to which even the moſt diligent obſerver is oftentimes ſubject in his endeavours to make the earlieſt opening to truth, Mead found

that this venom was acid, and that it changed the dye of the turneſol red, and even gave a reddiſh tinge to the ſirop of violets.

A few years after, Mead himſelf, in a ſecond edition of his work on poiſons, retracted all he had advanced on the acidity of the venom of the viper, and confeſſed, as a candid and ingenuous man, that it neither changes the dye of the turneſol, nor the ſirop of violets red, and that it is neither acid nor alakaline. Doctor James, who aſſures us that he repeated the experiments of Mead, has latterly found this venom to be acid; he however does not ſpeak of the poſterior experiments of the above cited authour; neither does he inform us how, ſuppoſing him right at firſt, he was deceived on the ſecond occaſion. This manner of publiſhing ones ideas, or ones experiments, neceſſarily tends to perpetuate doubts and hypotheſes, ſince, after all, the authority of one man is of as much weight as that of another, and ſince we cannot gueſs which of the two is in the wrong. Another writer, ſtill more modern than Doctor James, has received it as a truth, that the venom of the viper is acid; ſupporting his opinion on the bare authority of Mead, without telling us, that this authour has ſince denied its acidity.

It was natural to conceive, that experiment itſelf had demonſtrated to theſe writers, that Mead was miſtaken the ſecond time, and that he was right in his firſt trials, when he found the venom acid; and this conſideration obliged me to examine the

matter

matter afreſh. I hope that no doubt will any longer remain, and flatter myſelf that I have diſcovered the errour into which Mead fell when he firſt examined this venom; an errour againſt which Doctor James was not able to guard.

I have ſometimes, but rarely, found, that the venom of the viper gives the dye of the turneſol a light-red colour. This circumſtance, inſtead of inducing me to believe the venom acid, excited me rather to examine once more into the cauſe of it, which might be accidental. I obſerved that in theſe caſes it was not very pure, and on examining it with a microſcope, diſcovered globules of blood floating in it. I then examined the mouth of the viper, and found the two bags or ſheaths which cover the teeth ſlightly inflamed. It is not uncommon to meet with vipers that are naturally in this ſtate, and it is ſtill more frequent to find theſe bags reddened after the vipers have bit. We likewiſe frequently ſee the venom ſtained with blood, if its receptacle is too ſtrongly compreſſed. All theſe caſes may happen, and in all theſe caſes the dye of the turneſol may become red, without ſuppoſing an acidity in the venom. It is, therefore, not unlikely that Doctor James has been deceived in the ſame way with Mead. It is certain, that in the few caſes in which I have found the dye of the turneſol reddened, the venom was not pure, but was mixed with blood.

Aware of all theſe accidents, I took the utmoſt precaution in collecting the venom. I generally

cut off the head of the animal at a blow. Some hours after, when the muſcles had loſt their motion, I opened the mouth carefully, and contrived that the points of the canine teeth ſhould be free from their ſheaths. I then made a gentle preſſure on the receptacle of the venom, and received the latter in a glaſs, as it flowed out at the point of the tooth. In this way it is uſually ſo pure, that it appears, when viewed with a microſcope, like an oil, more or leſs yellow. No extraneous matter is obſerved in it; and when I accidentally thought I perceived corpuſcles floating in it, I did not employ it in the experiments of which I am about to give a detail.

When the venom was drawn in this way from the tooth, I never could perceive that it changed the dye of the turneſol red, however often I made the experiment, and I repeated it very many times. In moſt caſes, I began with uniting a drop of venom with thirty drops of the dye or tincture, and not finding the colour of the latter to be changed, I added another drop of the venom, proceeding in this way to a tenth, or one-third the quantity of the tincture, which never either reddened or changed its colour, but only appeared not quite ſo clear as before. I repeated this experiment too often to apprehend that I was miſtaken.—I not only tried the venom with the tincture, or dye, of the turneſol, but likewiſe made the ſame experiments on the blue juice of radiſhes, a liquor very ſenſible to the action of acids, even of the weakeſt of them. It continued blue as before, without my being able to obſerve

the ſlighteſt change in it. I likewiſe had paper well tinged with the juice of radiſhes, and let fall large drops of venom upon it;—the venom ſoon dried, and, except a yellowiſh tinge it gave where it fell, I could perceive no change in the colour of the paper.

On ſeveral other occaſions I diluted the venom with water, but could find no greater change in the paper on which I dropped it, than when I tried it pure.

I cannot deny but that I ſometimes obſerved a weak reddiſh tinge on the blue paper, when I made the experiment in the following manner:—I covered a large ball of cotton with the paper, and on forcing the viper to make a ſtrong bite at it, perceived this very pale tinge of red, at the parts the animal had pierced with its teeth. I did not, indeed, multiply my experiments ſufficiently to be able to ſay with certainty whence this tinge proceeded in theſe circumſtances. We may ſuſpect that it was owing to a ſmall quantity of blood from the mouth, blending itſelf with the venom; it is, however, very certain, that that taken from the veſſels, neither changes the die of the turneſol, nor the juice of radiſhes, red.

But even though we ſhould agree that the venom of the viper may be capable of giving a red tinge to theſe liquors, does it follow of neceſſity, that the volatile alkali is a certain remedy againſt this venom, and that the latter occaſions death preciſely becauſe it is acid?

The rock that men uſually ſplit upon, a rock that the moſt circumſpect philoſophers have not always been able to avoid, is that it is ſufficient for them to find a circumſtance which accompanies the effect, to be too readily perſuaded that it is the cauſe of it.

The innate deſire we have of knowing every thing, makes us ſtrive to explain every thing. If we ſee an effect produced after the application of any given ſubſtance, we immediately endeavour to ſee if there may not be ſomething in that ſubſtance that may ſerve in ſome way to explain the effect; giving ourſelves very little trouble to examine whether the cauſe we have diſcovered is proportioned or not to the effect produced. This errour ſeems to have been committed by two men of the firſt talents, Mead and Juſſieu. Mead, when he publiſhed the firſt edition of his work on poiſons, perſuaded of the acidity of the venom of the viper, judged that it muſt neceſſarily kill animals, becauſe it coagulates the blood as acids do.—Juſſieu, perſuaded likewiſe of the acidity of the venom, from the authority of Mead, immediately found a ſpecifick againſt it in the volatile alkali (a).

(a) Juſſieu was not the firſt after Mead to recommend the uſe of the volatile alkali againſt the bite of the viper, but as he made a brilliant cure, it is to him that this remedy owes its greateſt reputation.

The

The venom of the viper, as well as many other ſubſtances, is formed of ſeveral principles we are ſtill ignorant of. All the qualities we find in bodies do not conſtitute their real nature;—ſome of theſe qualities are accidental, others are not ſo. The acidity, even though it ſhould be conſtantly obſerved in the venom of the viper, may, nevertheleſs, be nothing more in it than an accidental quality; and the venom, in ceaſing to be acid, may not ceaſe to be a poiſon. Chemiſtry furniſhes us with a thouſand ſimilar examples. It is, therefore, improper to deduce the cauſe of the death from the acidity, and to deduce from the ſame acidity the uſe of the volatile alkali, as a remedy; for even ſuppoſing the venom to be conſtantly acid, and that this acidity cannot be ſeparated from it, does this enable us to ſay, that it kills becauſe it is acid, and that the volatile alkali is its ſpecifical remedy, becauſe it is capable of ſaturating it? The venom of the viper may likewiſe have ſeveral other qualities that we are unacquainted with, and may occaſion death by each of them ſeparately, or by all of them together. Why then are we to ſuppoſe, that it derives its noxious qualities from its acidity? There are arguments that demonſtrate the contrary.

Water abſorbs about its own bulk of fixed air, and conſequently a cubick inch of water can contain but very little more, if it does contain more, than a cubick inch of this air. It is not yet proved that a cubick inch of fixed air weighs an entire grain. A cubick inch of water weighs about 373 grains,

and conſequently the fixed air contained in a cubick inch of water, cannot be more in weight that its 373 part. Now a cubick inch of water impregnated with fixed air, is capable of giving a red tinge to 60 cubical inches of the tincture, or dye, of the turneſol, that is to ſay, to 22380 grains. Whence we ſee that the $\frac{1}{22380}$ part of a grain of fixed air is capable of beſtowing a ſenſible tinge of red on a grain of the dye of the turneſol. Now granting this hypotheſis, there cannot be at moſt in a grain of venom more than the $\frac{1}{22380}$ part of acid matter, and ſince the thouſandth part only of a grain in weight of the venom is capable of killing a ſparrow, as will appear by and by, we muſt ſuppoſe, that the $\frac{1}{22380000}$ part of a grain of acid can kill an animal ſimply as an acid principle.

Who does not now ſee, that even though it ſhould be granted that the venom of the viper gives a red tinge to the dye of the turneſol, it would not, on that account, follow, that it would kill becauſe acid? Its acidity would be ſo inconſiderable, that it would produce no ſenſible change in the animal body. And where is that violent acid, or any other principles of bodies, which is active to ſuch a degree, that in diminiſhing its quantity it does not at length become innocent?

Let any one ſuppoſe, if he will, that the acidity of the venom of the viper is as great as that of the *glacial* oil of vitriol (oil of vitriol concentrated to the conſiſtence of *ice*) itſelf. If the mortal effects of the former depended on its acidity, the glacial vitri-

vitriolick acid, thrown on a wound, although in a very ſmall quantity, would occaſion the death of animals. Glacial oil of vitriol applied to a wound, may indeed render the ſtate of it worſe, and may corrode the fleſh, but will not kill the animal on which it is tried. Very little of it can be introduced into the circulation of animals, and the little that is introduced is then weakened by the blood with which it mixes. It is true, that, as well as the venom, it may kill, if injected in a ſmall quantity; but this only happens becauſe it is not yet mixed with the humours, and weakened by them. Both the venom of the viper and the oil of vitriol may be abſorbed by the veſſels, and notwithſtanding the former is abſorbed in a very ſmall quantity, and very much diluted by the blood, it will kill an animal which will not be killed by the oil of vitriol. The venom of the viper does not therefore occaſion a very ſudden death from its acidity, but from other principles as yet unknown to us.

Mead, who changed his opinion as to the acidity of the venom of the viper, never wavered however in his ſentiments in regard to its ſuppoſed ſalts. He has always remained in the perſuaſion of having obſerved them floating in the yet fluid venom, ſoon after having taken it from the animal; and not only believes in the exiſtence of theſe floating ſalts in the venom, but aſſerts that the venom itſelf changes to a ſimple ſaline network, of a very beautiful ſtructure, which he compares to a ſpider's web. He ſpeaks of the ſolidity and firmneſs of theſe ſalts,

which he deſcribes minutely, and even gives a ſeparate drawing of them. He adds, that he has diſcovered here and there in theſe ſalts, ſmall circularly-formed knots, which are extremely ſolid, and never loſe the ſhape they have at firſt taken.

This ſubject, which appeared to me extremely intereſting, I examined very extenſively, in my work publiſhed in Italy, which forms the firſt part of this publication. I even flattered myſelf, at that time, that I had not only demonſtrated the errour of Mead in an inconteſtible way, but had likewiſe diſcovered the ſource of it. To refute an errour in phyſicks in a deciſive manner, nothing can be more effectual than the recurring to its origin. But even this does not ſeem ſatisfactory to certain authours, who perſevere in maintaining, after the authority of Mead, that the venom of the viper is a maſs of ſalts; notwithſtanding it is more than twelve years ſince Mead was refuted on this point. I demonſtrated at that time, that this venom is an homogeneous fluid, which, when taken pure from the tooth, is never found mixed with ſalts floating in it, nor with other heterogeneous particles; and that theſe floating corpuſcles, when they are to be found, are merely accidental, and are by no means ſalts. The ſmall knots ſeen by Mead, are nothing more than ſmall bubbles of air interſperſed in the venom. Theſe ſmall air-bubbles are never ſeen when the venom is taken immediately from the veſicle, and may be made to appear at pleaſure, by taking it blended

blended with the ſaliva of the animal, from the mouth of the viper *(a)*.

The ſaline net-work, which Mead ſays he obſerved, and which has been deſcribed by many authours after him, is no other than the fragments of the dried venom, which, when taken from the tooth, and put on a bit of glaſs, very ſoon dries, and whilſt it is drying cracks in different parts, preſenting pieces and fragments very different from real ſalts. The Count de la Garaie made *ſalts* of the ſame kind, by thoroughly drying his extracts on earthen plates, the glazing of which gave the hardened fragments a kind of ſhining ſaline appearance.

If a drop of the venom of the viper, put on a bit of glaſs, is examined with a microſcope, the ſubſtance of it will be ſeen to crack gradually at the circumference, where it dries ſooneſt. The fiſſures in

(a) To have demonſtrated the falſehood of any opinion whatever, is not a ſufficient cauſe for its being laid aſide, if it is generally adopted by authours. Nothing leſs is needed for this effect than the renewal of the entire generation, to the end that it may flatter itſelf, that it cannot be reproached for rejecting an error, ~~that it~~ has not committed. It required half a century to eſtabliſh the circulation of the blood, and the attraction of Newton, amongſt philoſophers. Man, always filled with a ſecret pride, thinks that he is humbled if he diſcovers himſelf liable to err; and the vulgar, never to be truſted in their deciſions, are of the ſame opinion. We have unfortunately too many examples of this kind, not to perceive that the love of truth is by no means the firſt ſpring of human actions.

this

this part are ſmaller and more crooked than elſe-where; but, by continuing to obſerve the venom, larger, broader, and deeper ones, which advance towards the centre of the drop where they end and meet, are ſeen at every part of the circumference. Theſe crooked lines are obſerved very diſtinctly with a microſcope, running to the centre, and lengthening in ſuch a way that one might miſtake them at firſt ſight for ſmall ſnakes, writhing themſelves from the circumference of the venom to the centre. After all the fiſſures are formed in this way, they enlarge ſtill more in proportion as the venom becomes drier, and occupies a leſs ſpace on the glaſs.

I do not know any microſcopical obſervation more certain and more evident than this, and in regard to which one may aſſure ones-ſelf with better grounds, that circumſtances are thus, and not otherwiſe. But that not the ſmalleſt doubt may remain, even in thoſe who may not have an opportunity to repeat my experiments, I have thought it incumbent on me to to repreſent, by ſeveral figures, a drop of venom in the act of deſiccation. It will be ſufficient to give a glance at theſe figures to be ſatisfied of the truth.

Fig. 1. of plate II. repreſents a drop of venom at the moment of its beginning to dry on a bit of glaſs. The fiſſures that are the moſt curved, at the circumference of the drop, are already entirely formed, the venom beginning to dry at the circumference. The others are ſeen becoming ſtraighter, lengthen-

lengthening, and approaching to the centre, where the venom dries the ſloweſt. When it is perfectly dry, the firſt figure changes to the ſecond, (ſee figure 2.) in which the fiſſures appear carried on to the centre, after having taken different curvatures. The fiſſures in the centre are broader, becauſe the venom, which is there in a greater quantity, ſeparates more on that account in drying.

Fig. 3. repreſents ſeveral fragments of the dried venom, in which the fiſſures are deſcribed by ſpiral lines. Theſe ſpires, as at *a*, are formed particularly, when the venom is dried in a conſiderable quantity, and when it is pretty thick on the glaſs of a watch. The fragments, which in this caſe are pretty large, open in the middle, and the opening, as I have juſt ſaid, is of a ſpiral form. The letter *e* repreſents a cleft that ſeparates the fragments from each other.

In Fig. 4. a drop of venom is repreſented, taken from the mouth of the viper, and dried. The ſmall balls, or knots, of Mead are ſeen in it, as at *o*. Theſe ſmall balls are real bubbles of air, which are made to diſappear with the point of a needle, as all air-bubbles are that are produced in fluids. Letter *m* repreſents a cleft that ſeparates the fragments, as above.

It is an errour then, founded on ill-contrived experiments, that there are ſalts floating in the venom of the viper; and the regarding the fragments of this venom, when it is dried, as ſalts, is another errour. It is equal and homogeneous through-

throughout, and nothing of this can conſequently be obſerved in it.

Mead, who regarded the venom of the viper as a maſs of ſalts, likewiſe believed it to be cauſtick and acrid when put on the tongue. He quotes himſelf and ſeveral of his friends, as having taſted it. He likewiſe obſerves, that when the viper bites, and when the venom begins to find its way into the wound, the animal cries out, writhes itſelf, and and exhibits other manifeſt ſigns of pain. Without pretending to decide at all on this queſtion, which I have likewiſe examined in the firſt part of this work, I ſhall obſerve here, that the experiment on dogs, which howl when they are bit, is not a certain and evident proof of the cauſtick nature of the venom. Perhaps when it is united in theſe caſes with the fluids of the animal, it is decompoſed, and acquires qualities it did not poſſeſs a moment before. It is true, that this howling which is mentioned, is ſometimes obſerved, but not always, and may be occaſioned by its frequently happening that a nerve is pricked by the teeth of the viper, in which caſes the venom may cauſe the ſame pain as any other body, or ſimple fluid, applied to the nerve itſelf.

If Mead taſted the venom, and found it cauſtick, I have taſted it likewiſe, and have made others taſte it, and we have neither found it cauſtick nor acrid. According to my ſentiments, it has no kind of taſte when put on the tongue, and is neither perceived to ſting or heat the part. It is true

true that a ſenſation is felt ſoon after, which may have made thoſe who believed it compoſed of ſalts, and who waited for ſome extraordinary change, ſuſpect that it was cauſtick and hot. The ſenſation it leaves when taken by the mouth, is that of a torpor or ſtupefaction in the part it touches. The tongue particularly ſeems numbed; it even appears to be grown larger, and its motions are ſlower and more difficult. This is certainly extraordinary, but appears very different from the effects occaſioned by cauſtick and acrid ſubſtances, when put on this part.—Laſtly, Mr. Troja wiſhed to taſte it himſelf, and aſſured me that he found it neither hot nor cauſtick, but that this ſenſation of torpor and ſtupefaction was the conſequence of it in the mouth. I can likewiſe take upon me to ſay, that I put five or ſix drops at a time into the mouth of ſmall animals, ſuch as rabbits, guineapigs, &c. without even having been able to obſerve any ſwelling or redneſs. Theſe experiments, when made on man, cannot be obſerved without a degree of repugnance, ſince, after all, a ſmall excoriation in the mouth, or on the tongue, may cauſe them to be too dearly paid for by the obſerver. I conceived that I could aſſure myſelf as to this particular in another way, and on a part even more ſenſible than the tongue itſelf; that is to ſay, on the eyes of different animals.

I put ſometimes one, and ſometimes ſeveral drops of venom on the eyes of a cat, and kept its eyelids open by force. I let it fall into the eyes of ſeveral

ſeveral rabbits without their perceiving it, and did the ſame thing to dogs. It was ſeen running over the tranſparent cornea and opake cornea, and getting within the eyelids. I could not perceive in any of theſe caſes that it acted as a cauſtick or acrid ſubſtance.

If Mead was miſtaken when he believed the venom of the viper to be compoſed of ſalts, he was not miſtaken, however, when he aſſerted that it was neither acid nor alkaline, ſince, in effect, it neither effervesces with alkalies or acids.

It is needleſs, after the experiments recited in the firſt part of this work, to enter here into a detail of thoſe I was induced to repeat upon this occaſion, and which can no longer leave any doubt in the minds of thoſe who are ſkilled in obſerving. It is an eſtabliſhed truth that the venom of the viper does not effervesce with any of the mineral or vegetable acids, nor with any kind of alkali we at preſent know of. I have repeated theſe experiments too often to have any doubt of having been miſled by them.

But it is not ſufficient to have ſatisfied ourſelves that the venom of the viper is neither acid nor alkaline; that it is not compoſed of ſalts; and that it is not corroſive to the palate; to inſtruct us in what it really is. I do not know with what other ſubſtance that is better known, it may be made to agree. It is principally to this point that the efforts of obſervers ſhould be directed, ſince it is certain that we are not thoroughly acquainted

with

with the true nature of any ſubſtance, although we are more or leſs acquainted with the properties of certain ſubſtances.

When the venom of the viper is yet liquid, it unites in a greater or leſs degree with acids. But we muſt likewiſe examine it when dry.

I put ſeveral drops of very pure venom into the concave part of the glaſs of a watch; as it dried, it became yellow, and full of cracks. I poured oil of vitriol on it, but no viſible ſolution followed. I raiſed from the bottom of the glaſs, with a capillary tube, ſeveral fragments of the venom, which floated in the oil of vitriol without diſſolving. At length, after ſome time, they ſeemed to begin to divide a little, and though they were indeed reduced to a kind of liquid paſte, ſtill preſerved their natural colour. There did not appear to be a true and perfect diſſolution of them, at leaſt during the time I obſerved them.

The marine acid, when poured on the dried venom, acts pretty much in the ſame way as the oil of vitriol. The fragments of venom do not appear, in a ſtrict ſenſe, to be diſſolved by this acid, although they are ſoftened by it.

The nitrous acid ſeems to have no greater power to diſſolve the dried fragments of venom, although it at length ſoftens them. Notwithſtanding the venom is rendered flexible by this acid, it ſtill preſerves a certain conſiſtence or tenacity which keeps it together, and it becomes yellower. If examined

in

in this ſtate, it appears to be compoſed of an infinite number of very ſmall ſpherical corpuſcles.

Thus then it appears that the ſtrongeſt acids have but a very ſlow and weak action on the dried venom of the viper, and that the diſſolution they at length occaſion is but a very imperfect one.

Vegetable acids, however concentrated they may be, do not diſſolve this venom better than do the mineral ones; and alkaline ſubſtances have no greater tendency to this effect.

I was likewiſe deſirous of knowing whether eſſential oils would diſſolve it, and on trying them did not find them to poſſeſs that property.

The hepar ſulphuris makes no greater impreſſion on it.

Theſe experiments, which I varied ſeveral ways, made me ſuſpect by degrees, that the venom of the viper might be either a gummy or a lymphatick ſubſtance, ſeparated from the blood of the animal. I had obſerved a long time before, that the dried venom appeared to be tenacious, like one of the ſtrongeſt gums, when broke betwixt the teeth. Freſh experiments were neceſſary, however, to be certain that it poſſeſſed the nature of a gum.

Chemiſts know that gums neither diſſolve in ſpirit of wine, nor in oil; but that they diſſolve very readily in water. This kind of examination might without doubt be ſatisfactory, but it was firſt neceſſary to prove that it was not of the ſame nature as animal lymph, or the white of an egg. We know that theſe ſubſtances coagulate in warm water,

ter, inſtead of diſſolving, as gums do. I got ready for this trial a great quantity of venom, which I kept in a ſmall capſular glaſs till it became perfectly dry. On this venom I threw at once about half an ounce of boiling water, by which it was inſtantly and effectually diſſolved, inſtead of being coagulated. On repeating this experiment ſeveral times, the conſequence was invariably the ſame. The water, after having been thrown into the glaſs, ſtill preſerved upwards of fifty degrees of heat.

Having thus, by direct experiments, excluded the hypotheſis of a lymphatick animal matter, I proceeded to the experiment of the ſpirit of wine.

I had a good quantity of venom dried as uſual in a ſmall glaſs, and poured on it half an ounce of highly rectified ſpirit of wine. I left it in an undiſturbed ſtate for upwards of two hours, when I found the venom undiſſolved at the bottom of the glaſs. I broke it into ſeveral ſmall bits with the ſharp point of a ſmall glaſs tube, and ſhook the whole together for ſome time. There was, however, no diſſolution, the ſmall pieces of venom continuing whole, hard, and of their uſual colour. This experiment will always ſucceed in the ſame way, if the ſpirit of wine is good; but if it ſhould contain too much phlegm, the venom may be partly diſſolved by it. Even this laſt circumſtance proves that the venom of the viper is a gummy ſubſtance, ſince gums are very readily diſſolved in water, which likewiſe diſſolves the dried venom, as I have aſſured myſelf an infinite number of times.

If the venom is perfectly pure, the water does not loſe any part of its tranſparence. Diſtilled water is the beſt calculated for theſe experiments.

I have frequently held the dried venom to the fire, and have increaſed the heat by degrees, but it has never melted. If it is thrown on a live coal, it ſwells and puffs up, but does not begin to take fire till it has aſſumed the appearance of a coal.

Another experiment now remained to be made, to render this matter deciſive.

All chemiſts know that gums diſſolved in water are precipitated by ſpirit of wine; and that in this trial, the water in which they are diſſolved becomes very white.

I put equal proportions of water into two ſmall glaſſes, and added to one of them a quantity of the venom of the viper, and to the other an equal quantity of gum arabick. The ſolution of gum arabick, which was made by heat, being reduced to the temperature of the liquor in the other glaſs, I poured ſeveral drops of ſpirit of wine into each of the glaſſes. The number of drops thrown into each was pretty much the ſame, when I began to perceive a whitiſh cloudineſs, which diſappeared a moment after in both ſolutions, at every drop of ſpirit of wine poured into them. On continuing to throw an equal quantity of ſpirit of wine into each glaſs, I ſaw the white cloud, inſtead of diſappearing, extend itſelf over the fluids, which became whiter and more opake, at every addition of the ſpirit. On ceaſing to throw it in, I perceived that the white matter began

gan to precipitate; and on adding a few drops of the ſpirit afreſh, found that there was no longer any ſeparation in either of the liquors. At the end of twenty-four hours the precipitation was complete, and there was at the bottom of each glaſs, pretty nearly the ſame quantity of an equally white, ſoft, and paſte-like, ſubſtance.

The venom of the viper, when diſſolved in water and precipitated by ſpirit of wine into the form of a white powder or meal, cracks in different parts when dried afreſh, and its fiſſures are of the uſual reticular form.

When a clear and tranſparent oil of vitriol is mixed with the venom, precipitated by ſpirit of wine, and dried in a glaſs, it becomes at the end of a certain time, of a dark vinous colour. The ſame changes are obſerved in the ſolution of gum arabick in water, precipitated by ſpirit of wine. This gum, in drying, likewiſe adheres to the glaſs and cracks, and if a few drops of oil of vitriol are thrown on it, they become in the ſame ſpace, of a dark vinous colour. The analogy betwixt the venom and the gum cannot be more perfect. They alike diſſolve in water; they are precipitated in the ſame way by ſpirit of wine; the precipitated powder or meal is of the ſame colour; both of them crack in drying; oil of vitriol does not ſoften them till after ſome time; and changes its colour in the ſame way with each of theſe ſubſtances.

I made another experiment on the venom of the viper, which though it does not prove any thing eſ-

ſential as to the internal nature of this venom, is ſtill a further proof that it has a great analogy to the gums.

I put ſix grains of very pure dried venom into a ſmall matraſs, and added to it fifty drops of nitrous acid, to throw off its airs. There came off from it, by the aſſiſtance of heat, as much air, or perhaps ſomewhat more, as the matraſs could contain. This was common air, a little changed in its qualities. I continued the fire, and a clouded air came off, which on examination I found to be compoſed, one third of fixed, and two thirds of phlogiſtick air.

Gum arabick, in the ſame circumſtances, likewiſe afforded fixed and phlogiſtick air, and the conſequences of both experiments were ſo perfectly ſimilar, that they might have been confounded together. It is true that gum arabick likewiſe affords nitrous air, but this only happens when it is in a conſiderable quantity. If the quantity is very ſmall, the little nitrous air it furniſhes, decompoſes itſelf, and unites with the common air in the matraſs.

It ſeems then to be demonſtrated, that the venom is in reality a gum; we at leaſt ſee, that it has all the properties and principal characteriſticks of ſuch.

This venom is found in an animal, is elaborated in its organs, and formed of its humours. It therefore ought to be conſidered as a true animal gum, particularly as the viper feeds on animals. Although we are unacquainted with any other animal gum, I do not think that the venom on that account

ſhould

ſhould be denied to be ſuch, ſince it has all the properties of a gum.' It ſhould therefore for the future, be inſerted in the catalogue of gums, and this diſcovery may perhaps induce naturaliſts to examine, whether a gummy ſubſtance may not likewiſe be found in ſome other animal.

Allowing the venom of the viper to be a real gum, this will not lead us to conceive what it is that conſtitutes it a venom, ſince it is a known truth that gums are not ſo, and that they may be employed with impunity. It would be ſuperfluous to relate the experiments I made on this ſubject, out of pure curioſity. I aſſured myſelf in a thouſand ways, that gum arabick is entirely innocent when applied to wounds. But ſuch is the condition of man, and ſuch is the nature of what we call ſcience. We at length arrive at certain bounds, to carry us beyond which all our efforts are to no purpoſe. The idea that the venom of the viper is a gum of ſome kind, does not ſerve in the leaſt to explain to us, how this gum brings on a violent diſeaſe in an inſtant, and how it is that, in ſo ſmall a quantity, it deſtroys life in ſo ſhort a time. Whatever the principle that renders it venomous may be, the proportion of it is ſo ſmall, that it does not at all change in it the uſual properties of a gum; and the ſmalleſt veſtige of this principle cannot be traced, whether the ſtrongeſt microſcopes are employed, or the venom obſerved in any other way. The moſt active ſubſtances are rendered ſuch by quantities of matter that cannot be traced. The

point of a needle that has touched a variolous puſtule, preſerves its activity for years, and brings about violent changes in the bodies of ſeveral perſons ſucceſſively pricked with it.

How far are we ſtill from penetrating the depths of this myſtery! Through how many difficult and unknown ways muſt we not paſs, in getting ſome inſight into a matter ſo obſcure and difficult as this! Happy at length, if all the pains that are taken, if all the efforts that are made, do not prove totally ineffectual.

This diſcovery, which enriches natural hiſtory with a new gum, ought not to be neglected by naturaliſts. It may in time lead to a better knowledge of the nature of the venom of the viper, and of the complicated effects it produces. It may perhaps be one day uſeful to us in enabling us to comprehend; why animals with cold blood are ſo long in dying of the bite of the viper; why there are ſome that are not killed by it; and why the venom, in whatever way it is introduced into its body, is altogether innocent to the viper itſelf. If the animals with cold blood that die late; if the others that do not die; if the viper to which the venom is not at all hurtful; had humours or parts of ſuch a nature, that they could be but little, or ſlowly, or not at all changed by this animal gum: we might then in ſome way explain a ſubject which is as yet very obſcure, and which does not ſeem capable of being cleared up, till after we have acquired a thorough knowledge of the venom itſelf, and of the moſt latent

rent principles and qualities of the animal bodies on which it acts.

On Bees, Drones, and Waſps.

In the firſt part of this work, I related a few experiments on the venom of the ſcorpion, and on the humour which flows from bees when they wound with their ſting. I have had occaſion ſince to make ſome other obſervations, not only on bees, but likewiſe on waſps, hornets, and drones. I do not know that any naturaliſt has examined in a proper manner, the liquor with which theſe animals are provided, that wound with a ſting. Indeed Mead ſays that he found the humour of bees, to be compoſed of very ſmall ſaline needles, or points. He aſſures us that he examined it with a microſcope, and found it filled with theſe pointed ſalts. I do not know whether this obſervation made by Mead, has been confirmed or not by other naturaliſts, but can for my own part take upon me to ſay, that I never have been able to find any thing ſaline in this humour, whatever attention I paid in inveſtigating it, and notwithſtanding I employed the ſtrongeſt lens for that purpoſe. I am perſuaded that Mead has been miſtaken in this particular, as he was in obſerving the venom of the viper. He aſſuredly ſaw particles floating in this humour before it

it was dry, and immediately perſuaded himſelf that they could be no other than floating points.

We may eaſily conceive that Mead only examined this humour, in an impure ſtate, and mixed with corpuſcles that were foreign to it; and that this was ſufficient to induce him to believe it compoſed of ſalts. He was deceived on this occaſion, as he was in his opinion of the venom of the viper, in which there is nothing to be met with of all that he fancied he ſaw; and he ſeems in both caſes to have erred exactly in the ſame way. The humour of bees, after the manner of the venom of the viper, cracks in drying, and preſents the uſual ſharp and regular fragments. This was ſufficient to perſuade Mead that it was a true ſalt.

I can venture to ſay, that when the obſervation is well made, nothing can lead to ſuch an opinion. But if, in expreſſing the liquor from the bee's ſting, the greateſt care is not taken to prevent the breaking and mixing any thing with it, it may eaſily be charged with other irregular bodies; and when it is put on the port-object, ſome ſmall degree of motion may likewiſe be obſerved in theſe bodies, which may float in a greater or leſs quantity. But this accidental motion, which is not proper to theſe ſubſtances, ſoon ceaſes altogether, when the humour is left undiſturbed. By degrees it dries, and, in drying, breaks, cracks, and forms angles and points.

When the venom of the viper and the humour of bees, are dried and obſerved with a microſcope, no ſenſible difference can be obſerved between them.

I have

I have only taken notice, that the humour of bees, expoſed to the open air on a bit of glaſs, is much longer in drying than the venom of the viper, and that the cracks or fiſſures in the former, are likewiſe formed much later than thoſe in the latter, ſuppoſing the degree of deſiccation in the two fluids alike.

Theſe two humours not only agree in the appearances their parts preſent in drying, but likewiſe in other qualities. If a bit of the dried humour of bees is ſtrongly compreſſed betwixt the teeth, it, as it were, glues them faſt together; and exactly the ſame thing happens on trying the venom of the viper, and all hardened gummy ſubſtances. The dried humour of bees likewiſe diſſolves in ſimple water, and reſiſts the action of ſpirit of wine, as the venom of the viper and gums in general do; ſo that I am almoſt inclined to believe that, as the venom of the viper is moſt aſſuredly a gummy ſubſtance, this humour is ſo too. Indeed the quantity one is able to collect of it is ſo very ſmall, that one can ſcarcely attempt to make any certain experiments on this ſubſtance; the conſequences, however, of thoſe I have made have been ſufficiently uniform to lead me to think that I cannot eaſily have been miſtaken in what I have conjectured.

I have met with the ſame ſucceſs in examining the humour of waſps and drones, and of the other flying inſects in general, that wound with a ſting, and are provided with a humour. In all theſe, the humour is bitter and acrid, and has all the appear-

ance of being of a gummy nature. When left to dry on a bit of glaſs, it cracks throughout like the venom of the viper, and when chewed, is tenacious, glutinous, and elaſtick.

But it muſt not therefore be thought to be the ſame as the venom of the viper, and that it has all the other qualities of this poiſon. The venom of the viper neither has any ſenſible taſte when taken into the mouth, nor is ſufficiently acid to give a red tinge to the tincture of turneſol, or juice of radiſhes. The humour of bees, and of the other analogous inſects, the moment it is applied to a piece of paper that has been previouſly ſtained with the juice of radiſhes, gives it a ſlight red tinge, which afterwards changes to a pale yellow, ſo that one would conjecture that this humour deſtroys the blue colour of the paper. This experiment, which has been repeated ſeveral times, and always attended with the ſame ſucceſs, proves that this humour is united with an acid, and not with an alkaline principle; we ſee, however, at the ſame time, that the quantity of acid it contains is very ſmall, and abſolutely incapable, as an acid principle, of occaſioning the ſmalleſt ſenſation on the tongue, or in the part pricked by the ſting of the animal.

A quantity of water impregnated with an equal bulk of fixed air, gives a red tinge to paper ſtained with the juice of radiſhes. This tinge, which is pretty ſtrong, continues a conſiderable time. A ſmall quantity of water impregnated with fixed air, ſcarcely contains a ſufficient degree of acid to be ſenſible

ſible to the taſte, and is likewiſe entirely innocent when applied to wounds.

We muſt therefore regard the hypotheſis of thoſe naturaliſts, who have advanced that this humour occaſions a ſwelling in the parts into which it is introduced, and that the volatile alkali, as ſaturating the acid principle, is a remedy againſt it, as falſe and erroneous.

Experiment ſeems to indicate that this humour acts by the medium of a bitter and cauſtick principle, which is neither acid nor alkaline. If it is put on the tongue, it has a hot bitter taſte, as I obſerved before, and not that of an acid or alkaline ſubſtance. There are many ſubſtances which, without being either acid or alkaline, are hot and acrid to the palate, and are productive of violent and diſagreeable ſenſations. Cantharides, and ſeveral aromatick plants, are of this claſs. In the preſent caſe it appears certain, that neither the pain, (which is frequently inſupportable, and greater than that which would be cauſed by oil of vitriol itſelf) nor the ſwelling nor inflammation of the parts, can be brought on by an acid principle introduced into the ſkin of the animals that have been ſtung; and therefore the theory laid down by certain authors to explain the effects of this humour, muſt be regarded as abſolutely falſe, and the conſequences they have deduced from it as no truer than the theory itſelf. A pretended concentrated acid, a *naked* acid, an uncombined acid, and a phoſphorick acid that produces ſuch wonderful effects, are hypotheſes that

are

are not capable of reſiſting the inveſtigation of reaſon and experiment, and are unworthy the enlightened age we live in. It is no longer the ſeaſon to imagine nature; we muſt conſult her. If chemiſtry has increaſed the number of our intelligences, the abuſe of chemiſtry has frequently retarded our progreſs in the ſciences. It has frequently led us into errour, and has ſubſtituted hypotheſes for facts and experiments.

Although bees, and the other inſects that are analogous to them as far as relates to the humour they throw out at their ſting, are not capable of killing, I think notwithſtanding, that if they are not conſidered as venomous animals in the moſt received ſenſe, they ſhould at leaſt be conſidered as animals that ſecrete in their bodies a ſmall quantity of a matter, which is not deſtructive ſimply becauſe it is in too ſmall a quantity. The moſt active poiſons and venoms, ſuch as arſenick, corroſive ſublimate, and the venom of the viper, when taken or applied in a very ſmall quantity, not only do not occaſion death, but do not even produce a ſenſible derangement, very far from their effects equalling thoſe that are produced by a large hornet, when it wounds with its ſting. Theſe quantities, however, although very ſmall, are capable of killing the ſmaller ſpecies of animals, whilſt more conſiderable ones are not ſufficient to kill thoſe of the larger ſpecies. Hence we ſee, that the difference entirely conſiſts in the quantity of the venom, and in the different degrees of ſtrength in the animal that receives

it, and not in the nature of the venom, which is always the ſame. The venom, for ſo I ſhall call it, of bees, is very active, conſidering the ſmallneſs of its quantity, and we may eaſily judge from the pain and inflammation it excites in an inſtant, that if the doſe of it were increaſed, it would produce the moſt violent derangements, and perhaps even a very ſpeedy death. Nay, I am almoſt inclined to think, that a grain in weight would kill a pigeon in a few ſeconds. The difference that is found betwixt the ſting occaſioned by a bee, and that of a hornet, notwithſtanding the difference in the reſpective quantities of their venom is but very inconſiderable, is very great. The ſame thing may be obſerved of the common ſcorpions of Italy, and thoſe of other countries, as well as of the bite of ſpiders. The larger produce in general the greateſt derangement, and thoſe of Africa, or of Aſia, even occaſion death: all of them, down to the ſmalleſt, poſſeſs a greater or leſs degree of activity.

There are other animals, particularly inſects, which when they bite or ſting, bring on a very violent pain and inflammation, ſo that they may reaſonably be ſuſpected of introducing a cauſtick and venomous humour into the wound. In this number we may reckon ants, which inſinuate into the ſmall wound they make in biting, a very ſharp and poignant humour, which they force from a veſicle ſeated in the hinder part of their body. I ſhall not make a digreſſion here to ſpeak particularly of this humour, becauſe I have treated of it in a very ample

ple way, in a paper of which the object was an examination of the *acids of animals*, &c. and particularly of *the nature of that of ants*, printed in the journal of the Abbé Roſier. I there demonſtrated that the humour of ants is a true acid, and that it is in reality the acid of fixed concentrated air, deprived of its elaſticity, and rendered liquid.

PART

PART III.

CHAPTER I.

Action of the Venom of the Viper on Parts of an Animal that have been previouſly bit.

THE ſubject of this part is the moſt intereſting one that the matter it treats of can preſent to a philoſophick obſerver.

All the queſtions that are here diſcuſſed become of conſequence, ſince they tend to throw great lights on the nature of venom. The animal economy itſelf is by their means better explained, and many hypotheſes that have been imagined, fall before experiment. It is the touchſtone that makes us ſoon diſtinguiſh all that does not belong to nature, all that is the effect of art, of prejudices, and of the imagination; in a word, of man.

Experiment alone may conduct us through the unknown paths of nature, and may lead us to new and unexpected truths. But at the very time that man, profiting by this torch, is making bold ſtrides towards the truth, and ſoars as if he meant to govern nature herſelf, ſhe ſtops him every moment,

ment, and by only diſcovering herſelf to him in part, ſeems afraid of being recollected; ſhe thus continually reminds him of his weakneſs, and ſhows him that his hopes are either vain, or confined within very narrow limits.

Man, who aſſigns to comets the courſe they are to keep, and who fixes the time that is employed by the light in its progreſs from the ſun to our hemiſphere, is not, with all this knowledge, acquainted with the air that ſurrounds him, or with the fire that warms him. Such is our condition, and ſuch is the ſtate of human ſcience.

The firſt queſtion that preſents itſelf, after what has hitherto been related, is to know whether the venom of the viper is a poiſon to all the animals with warm blood. It will be ſeen in a little time, that this large body of animals has not been ſeparated without deſign from the other, which comprehends thoſe that have the blood cold. When I ſay that a ſubſtance is venomous to an animal, I mean to expreſs, that it produces in it very violent diſorders, although it is only introduced into its body in a ſmall quantity.

To reply properly to the queſtion I have juſt propoſed, it is certain that all the animals with warm blood exiſting on the habitable globe, ſhould be bit by vipers. The ſubject is not ſufficiently intereſting to deſerve ſo long and difficult a labour. However, if the analogy betwixt the different animals with warm blood may be allowed, I am not afraid to advance, that the venom of the viper is a poiſon

to

to all of them. We have ſeen that it has proved ſo to all the ſeven ſpecies that have hitherto been examined; and I very well recollect that I could not find any animal in Italy, with warm blood, to which the venom of the viper did not prove a real poiſon. I tried it on all the birds I could meet with, and on all the quadrupeds I could procure, provided they were of a moderate ſize; as to the horſe, the camel, and the ox, ſetting aſide their bulk, I could not procure them eaſily for this purpoſe.

We may therefore, I think, conclude with a great deal of reaſon, that the venom of the viper is a poiſon to all the animals with warm blood; that is to ſay, that neither of them is beyond the reach of the effects it uſually produces, when it is introduced into the body in a ſufficient quantity.

The ſecond enquiry, which ſprings immediately from the firſt, is to know whether the venom of the viper is a poiſon to all the animals with cold blood.

It has already been ſeen in a former part of this work, that even the frog, a cold animal, and one very hard to kill, dies in a few hours, if it is bit by the viper. This, however, is not ſufficient to admit a certain concluſion, that all the other animals with cold blood would die in the ſame way. We frequently incur the riſk of being deceived by this method of employing analogies on too narrow and limited a ſcale.

A ſingle ſpecies of animals is not ſufficient to furniſh an analogical argument of any weight.

Had five or ſix hundred kinds of animals with cold blood been examined, and had certain ſymptoms of poiſon been obſerved in all of them after they had been bit, the analogy, in this caſe, would have formed an argument of probability, and we might have been enabled to draw concluſions on this ſubject, not only in regard to animals with warm blood, but likewiſe as to thoſe that have the blood cold.

We can ſcarcely do otherwiſe than ſuſpect that the venom of the viper is innocent to the viper itſelf. This animal, in all the diſeaſes or wounds of its mouth, would otherwiſe run a very great riſk of killing itſelf with its own venom. It is not very unuſual to find vipers with the bag or ſheath of their teeth inflamed and bloody. Small red ſpots are frequently obſerved in the mouth of this animal when it bites, and it is beſides eaſy to conceive, that if it ſhould be bit in the mouth by any other animal, its own venom would prove deſtructive to it, if its particular nature did not guard it againſt ſuch an accident.

The venom of the viper is conſtantly ſecreted and laid up, in the ſpongy gland. This gland has its canal continually open, through which the ſuperfluous venom that cannot be contained in the gland, is forced to ſhed itſelf into the viper's mouth.

However, it is eaſy to have recourſe to experiment. In the firſt part of this work a detail may be found of a great number of trials I made on this ſubject, and from which it reſults, that the venom

of

of the viper is not a poiſon to vipers, but that, on the other hand, it is altogether innocent to them. I was deſirous of repeating ſeveral of theſe experiments over again, and out of the great number, which brevity obliges me to omit, I think it ſufficient to relate a ſingle one.

After having enraged a viper very much, I forced it to bite itſelf ſeveral times in the part towards its tail; it, however ſuffered nothing from this, although it had certainly forced its teeth well into the part. I repeated this experiment on three other vipers, with the ſame ſucceſs. It is therefore very certain, that the venom, or bite, of the viper, is entirely innocent to this animal when it bites itſelf, and it likewiſe is when one viper bites another.

But this very ſingular exception is not confined to the viper. There are other animals to which this venom is innocent, and others again in which, although they were ſmall, one or two vipers are ſcarcely capable of producing any ſenſible change. I have mentioned ſome of theſe *cold animals (a)* in the firſt part of this treatiſe, but to come at the number of them, the experiments ſhould be extended to other ſpecies' I could not at that time procure, and on which I thought it ſuperfluous to make this trial.

If it is altogether extraordinary, that the ſame matter is entirely innocent to ſeveral ſpecies' of

(a) Animals with cold blood.

animals, and that it is mortal to an infinity of others, it is much more ſurpriſing, and at the ſame time more difficult to conceive, how, and by what principles, it happens, that an inſipid gum, as far as we can perceive, excites the moſt violent diſorders in ſo many very large animals, and that it does not bring about the ſmalleſt change in others that are incomparably ſmaller and weaker.

The known diſtinction of animals with cold, and animals with warm blood, which is only founded on a greater or leſs degree of heat, and on ſome other trivial difference in the circulation of humours, is of no uſe in the preſent caſe, ſince there are certain animals with cold blood that die of the venom, and others again that are not at all acted on by it.

If a compariſon is formed betwixt two cold animals, one that dies of the diſeaſe of the venom, and the other that ſurvives its action, they will be found to poſſeſs the ſame organs, the ſame circulation, an equal tenaciouſneſs of life, and, in a word, to the eyes of the obſerver, they will both of them appear perfectly alike.

What is it, then, that cauſes this matter which flows from the viper's tooth to be a poiſon to one, and not to the other? We are not only entirely ignorant of this, but it appears that we are likely always to remain ſo. To obtain ſuch a knowledge, it would be neceſſary to be acquainted with the moſt hidden nature of this extraordinary animal gum. It would be neceſſary to penetrate into the moſt internal and latent ſubſtance of the ſolids and fluids

fluids of animals with cold blood, to know the mechaniſm of their organization, and to comprehend perfectly the principle of life. We might then reply to all that could be aſked on this head. But how is it poſſible to acquire ſo extenſive an information, whilſt the activity and penetrability of our organs are ſo limited and confined?

But if we are not permitted to know what this very active principle of the venom of the viper is, which when it is introduced into a living animal cauſes its death; we are allowed, however, to enquire into the quantity of this venom, that is neceſſary to kill an animal of a certain ſize. This enquiry, very curious in itſelf, cannot but be of ſome uſe in the practice of medicine, particularly in cautioning us againſt thinking the danger greater than it really is, when any one of our own ſpecies has the misfortune to be bit by this animal.

To be able to ſpeak with ſome degree of preciſion, in this reſearch, it was proper to begin by determining very ſmall quantities of venom, and by introducing them without loſs into the ſubſtance of the body of a living animal. It was likewiſe expedient to operate on very ſmall animals, that would die ſoon and to a certainty, to the end that the conſequences might be leſs equivocal. It is true, that by an endleſs multiplication of experiments, the ſame conſequences might at length be obtained from large animals; but a longer time, and greater conveniences would be required, and one ought beſides

to be perſuaded of the importance of the undertaking.

In the following experiments, I made choice of ſparrows and young pigeons, knowing them by experience to be readily killed by the venom.

To determine ſmall known quantities of venom, I began by taking four grains in weight of the venom of the viper, and mixing with it eight grains of diſtilled water. I then, with a ſmall bruſh, ſpread it equally over a ſquare inch of thin paper. This may be done with a ſufficient degree of eaſe and preciſion to exclude any conſiderable errour, and indeed I found that the halves and quarters of the ſquare inch of paper were of the ſame weight when dried.

I cut this paper in two, and again divided one of the halves, continuing in this way till I had made ſix diviſions, reckoning the firſt. I then did the ſame with the other half, that I might have two pieces of the ſame ſize, and of each ſize, inſtead of one.

I ſtripped the muſcles of the leg, in ten ſparrows, of the ſkin, and bound upon them the ten bits of paper I have mentioned. The conſequences, beginning with the larger bits of paper, $\frac{1}{4}$, $\frac{1}{8}$, $\frac{1}{16}$, $\frac{1}{32}$, $\frac{1}{64}$, were as follows. Of the two ſparrows to which the papers marked $\frac{1}{4}$ were applied, one died at the end of fifteen minutes, the other not till the end of thirty-five. One of thoſe with the papers marked $\frac{1}{8}$ died at the end of an hour, the other ſurvived. One of thoſe with the papers marked $\frac{1}{16}$ died

died at the end of two hours, the other recovered. One of the two with the papers marked $\frac{1}{12}$ died at the end of two hours, the other at the end of five. And of the two with thoſe marked $\frac{1}{64}$, one died at the end of three hours, the other at the end of ſeven minutes.

On repeating this experiment, the conſequences were ſtill more irregular. I therefore abandoned this method, as altogether inſufficient and deluſory. This is probably owing to the paper, which, when put in contact with the humours of the animal, may not allow itſelf to be either equally or entirely deprived of the venom that adheres to it. This obliged me to have recourſe to another method, which is perhaps leſs exact in determining the preciſe quantity of venom, but which has afforded me conſequences as conſtant and as uniform as can be expected in ſo difficult an undertaking.

This is the method I employed:—I took a given quantity of venom, for example three grains, and ſpread it over a bit of glaſs, in ſuch a way that it occupied a determinate ſpace of a circular form.

The venom in the centre was not more than a quarter of a line in depth.

I procured a ſmall capillary glaſs tube, which terminated in a ſmall ſcoop of about half a line in diameter. I plunged this ſmall ſcoop vertically into the centre of the venom, and drew it out in the ſame direction.

To determine the quantity of venom that adhered to the ſmall ſcoop, and to know at the ſame time,

whether this quantity would be conſtantly the ſame, I put the three grains of venom, laid on the bit of glaſs, into a very nice ballance, and plunged the ſmall glaſs ſcoop ten times ſucceſſively into the liquor, taking care to wipe the ſcoop well every time. After the ten plunges, I found the equilibrium to be loſt, and that about the $\frac{1}{100}$ part of a grain of venom was deficient. I now plunged the ſmall glaſs ſcoop ten other times ſucceſſively into the venom, and the ballance having again loſt its equilibrium, found that the venom was diminiſhed about the $\frac{2}{100}$ part of a grain. With a little practice, one can make this experiment in leſs than two minutes, and in that ſpace the three grains of venom are not ſenſibly diminiſhed in weight by the natural evaporation, as I have aſſured myſelf by trial. I cannot take upon myſelf to ſay, that all the quantities are here rigorouſly the ſame; nay I agree that, in repeating this experiment ſeveral times, a ſenſible difference, which I have indeed met with myſelf, cannot fail to occur: but all theſe differences taken together can make but a very trifling variation in the quantity of venom that adheres each time to the ſmall ſcoop. On the whole, I can lay it down as an eſtabliſhed rule, that the ſmall glaſs ſcoop plunged perpendicularly into the venom, in the way I have deſcribed, carries away with it about the $\frac{1}{1000}$ part of a grain of the venom of the viper.

I laid bare a portion of the right leg of a ſparrow, and made a ſmall longitudinal inciſion into the muſcles with a lancet. Into this inciſion I introduced

at

at the very inſtant, the ſmall ſcoop armed with venom, and kept it in this ſituation for thirty ſeconds. The ſparrow died at the end of two hours, with a lividneſs of its leg.

I repeated this experiment on ſix other ſparrows, exactly obſerving the ſame circumſtances. They all died, one after the other, at the end of the following times, expreſſed in hours, 2, 2, 3, 4, 5, 7.

I again repeated it in the ſame way on twelve other ſparrows, and the conſequences were ſtill more irregular than thoſe of the preceding experiment. One of the ſparrows died at the end of four minutes, another at the end of three days, and another at the end of five. The ſymptoms of the diſeaſe were, notwithſtanding, indubitable in all the three. The nine others all died at the end of the times expreſſed by the following numbers, which denote ſo many hours; that is to ſay, 2, 3, 3, 5, 6, 9, 10, 12.

The firſt conſequences ſhow, that the quantity of venom I have mentioned is ſufficient to kill an animal of the ſize of a ſparrow, but that it produces in theſe animals very unequal effects, and a diſeaſe of greater or leſs violence. An animal that dies at the end of three minutes, and another perfectly ſimilar to the firſt, that does not die till the end of five days, prove that the diſeaſe in each of them has been very diſtinct. But ſuppoſing the quantities of venom which were introduced to have been equal, and that the inciſions were ſo too, a little more, or a little leſs, blood, oozing from the inciſed veſſels, might have cauſed all this difference, ſince it might have occa-

occaſioned a greater or leſs quantity of venom to enter into the circulation of the humours, or, to expreſs it ſtill better, into the animal itſelf.

I wiſhed to ſee whether I could bring on a more ſpeedy death by doubling the quantity of venom; and being at a loſs for a certain method of collecting this double quantity together, I made two inciſions inſtead of one, and introduced the ſcoop I have mentioned into each. Twelve ſparrows on which I made this experiment, all died, but at very different intervals. One died at the end of three minutes, another at the end of twenty-ſeven, and a third at the end of forty; the others at the end of the hours expreſſed by the following numbers, 1, 1, 2, 2, 2, 3, 3, 5, 6.

The experiments I made on ſparrows, and the method of introducing a given quantity of venom into the inciſion in their muſcles, have occaſioned me to make a very intereſting obſervation. I uſually kept the ſmall ſcoop in the inciſion for about twenty ſeconds, and obſerved that the lips of the wound became livid at the end of a certain time. I regarded this ſymptom as a ſure mark of the communication of the diſeaſe, and it will be ſeen by and by that I was not miſtaken in this conjecture.

I was deſirous of ſeeing what effect would reſult from a certain quantity of venom, applied to an animal larger than a ſparrow.

I ſtripped a part of a pigeon's leg of the ſkin, in ſuch a way that the muſcles beneath were entirely bare. Having made the uſual inciſion, I introduced

duced the ſmall ſcoop, which had been firſt plunged into the venom, and kept it there till I ſaw the lips of the inciſed part become livid; this happened, as in the ſparrows, in the ſpace of about twenty ſeconds. At the end of half an hour the leg became ſlightly livid, but neither ſeemed ſwelled nor hard. The pigeon neither died, nor ſuffered in any ſenſible degree.

I repeated this experiment on ſix other pigeons, exactly obſerving the ſame circumſtances. One of them had not the ſmalleſt ſymptom of diſeaſe, neither did the inciſion become livid, notwithſtanding I kept the ſmall ſcoop in it for upwards of a minute. Four others had ſymptoms of the diſeaſe of the venom, and two of theſe did not recover till after forty hours had elapſed. The ſixth, like the firſt, was not at all diſeaſed; the inciſion, however, I made in its leg, bled at the time I introduced the venom.

I repeated this experiment, with the ſame circumſtances, on ſix other pigeons. One of them died at the end of ſix hours. Three had all the ſymptoms of the diſeaſe of the venom, and did not recover till the third day. Two others had not any ſymptom of the diſeaſe. I think it proper to obſerve here, that the inciſions in the leg, in theſe two laſt, bled very ſenſibly when I introduced the venom. This ſhows that the blood which flows from the veſſels, may prevent the venom from entering them, or from remaining in them after it has entered,

I repeated this experiment over again on twelve pigeons, one of which died at the end of ten hours. Two others were exceedingly ill. The other nine had no ſenſible complaint.

Theſe new experiments ſhow, that the quantity of venom which uſually kills a ſparrow is not fatal to a pigeon. We ſee, however, at the ſame time, that a caſe may occur, in which ſuch a quantity of venom is introduced as is capable of killing a pigeon, notwithſtanding that the quantity employed in the experiment would ſcarcely have been ſufficient, generally ſpeaking, to kill a ſparrow.

I was deſirous of trying on pigeons, as I had done on ſparrows, what would be the effect of two inciſions, and a double quantity of venom.

Having laid bare the muſcles of a pigeon's leg, as uſual, I made two ſmall inciſions into them, introducing into each, the ſmall ſcoop armed with venom, in the accuſtomed way. The livid ſpot appeared at each inciſion, and almoſt the whole of the leg became livid, and remained in that ſtate for two days, at the end of which time the animal was perfectly recovered.

I repeated this experiment on twelve pigeons, and the conſequences were ſomewhat various. Two of the pigeons died at the end of three days. The others had all a lividneſs of their legs, and all recovered notwithſtanding.

On repeating this experiment on twelve other pigeons, four of them died; one at the end of ſix hours; another at the end of twenty; and the two

others

others not till the fifth day. All the others had the diſeaſe of the venom, but recovered.

Inſtead of making two inciſions only, I tried the experiment on twelve other pigeons, by making four inciſions at the ſide of each other. Nine of theſe pigeons died; one in ten minutes, two in an hour, two in two hours, and three in five hours. The other three had the diſeaſe of the venom, and their legs became livid, ſwelled, and hard.

What is the Quantity of Venom required to kill an Animal?

We may, I think, from the above experiments, determine with ſome probability, the quantity of venom it requires to kill an animal. This queſtion already begins to become important to us on our own accounts, ſince we may at length be enabled to flatter ourſelves, that the bite of the viper is not ſo dangerous as we have been hitherto taught to imagine it to be.

We have juſt ſeen that the $\frac{1}{1000}$ part of a grain of venom, introduced immediately into the muſcle by an inciſion, may be a ſufficient quantity to kill a ſparrow, although this animal does not always die in conſequence of the introduction of ſuch a quantity; and that it requires about four times as much to kill a pigeon. We may even ſuppoſe it to require about ſix times as much to kill the laſt animal to a certainty.

The

The ſparrows on which I made my experiments weighed ſomewhat leſs than an ounce each, and the pigeons ſomewhat more than ſix ounces each. Now let us ſuppoſe that ſparrows weigh exactly an ounce, and pigeons exactly ſix. The quantity of venom it will require to kill a large animal, an ox for inſtance, ſuppoſing it to weigh 750lb. will be about twelve grains; and it will require nearly two grains and a half to kill a man, ſuppoſing him to weigh the fifth part of what an ox weighs, that is to ſay 150 lb.

It is true that this calculation takes for granted ſome new hypotheſes more or leſs probable, but of which neither is unlikely. A ſufficient number of experiments are wanted, to render them either abſolute truths, or ſuſceptible of ſome reſtrictions.

The firſt hypotheſis ſuppoſed here is, that the venom of the viper acts on an animal in proportion to its quantity. There is nothing unreaſonable in believing this to be the caſe, ſince if a very ſmall portion of venom is capable of deranging the economy of an animal *to a certain point*, a greater doſe of it ought to produce a greater derangement, a more violent diſeaſe. Beſides we have ſeen, that animals bit ſeveral times by one viper, or by ſeveral, die ſooner than thoſe that are only bit once by a ſingle viper; and we know that a viper which bites ſeveral times, introduces freſh venom into the part at each bite.

The ſecond hypotheſis is, that the diſorder produced in the animal economy by the venom of the viper,

viper, is leſs in proportion, or rather that the power of the animal to reſiſt the action of the venom, is greater in proportion, as the animal is larger. This is generally ſo, although there may be exceptions to this law, that may prevent its being rigorouſly the caſe.

The third hypotheſis is, that from the effects produced in an animal of one ſpecies, we may argue as to the effects produced in an animal of another ſpecies; that is to ſay, from birds to quadrupeds. This argument is drawn from a ſimple analogy; but this analogy is at the ſame time formed betwixt animals with warm blood, and it may therefore be deemed of ſome weight.

Now granting that a viper of a middle ſize has in its veſicles two grains in weight of venom, it will require the venom of ſix vipers to kill an ox, and nearly of two to kill a man.

But if we reflect that a viper which bites, does not leave itſelf without venom; that at each bite, at leaſt for the firſt three or four, it may bring about the death of an animal with almoſt the ſame facility; it will not appear altogether unlikely, that it may perhaps require twenty vipers, each biting only once, to kill an ox, and five or ſix, with the ſame reſtriction, to kill a man.

CHAPTER

CHAPTER II.

Of the Time it requires for the Effects of the Venom of the Viper to become ſenſible.

A QUANTITY of the venom of the viper which ſcarcely weighs the $\frac{1}{100}$ part of a grain, produces, on being introduced into the body of a ſmall animal, ſo violent a diſeaſe, that death follows in a few minutes. It is therefore very clear, that it muſt poſſeſs a great degree of activity, and that its effects muſt be both ſudden and powerful. I have advanced in ſeveral parts of this work, that the venom of the viper renders the parts that have been bit, in animals, and that almoſt in an inſtant, incapable of exerciſing their uſual functions. I am at leaſt certain, that I have obſerved this effect in ſeveral that I have had bit. It has been ſeen that the venomed part becomes livid after it has received the bite, but that this does not happen till within a certain ſpace. The wounded parts ſoon become ſwelled and painful, and the adipoſe membrane is ſhortly after filled with a black and diſſolved humour; whilſt the blood that remains in the veſſels is black and coagulated.

One would naturally ſuppoſe, that the action of this venom on the organs of an animal is momenta-

neous, and that it is not different from that which takes place when two ſubſtances of different natures are blended together, and of which chemiſtry furniſhes us a thouſand examples.

Deſirous of purſuing this idea, and flattered with the hope of diſcovering ſome effect, or ſome particular, that might be ſerviceable to my preſent reſearches, I formed a new plan of experiments.

My firſt trials had for their principal aim, the obſerving of the changes the venom of the viper would produce, when introduced into a part cut from an animal, but ſtill warm and palpitating.

Experiments on the Limbs of an Animal, recently ſeparated from the Body.

At the very inſtant the part was cut off, I had it bit by a viper, ſo that when the experiment ſucceeded well, as it frequently did, there could ſcarcely paſs a ſecond betwixt the amputation, and the bite.

I made choice of young pigeons for this experiment, becauſe I had obſerved in theſe animals, that the venom of the viper very ſpeedily produces a livid ſpot, in the part of the muſcles through which it has introduced itſelf.

To make this experiment, a perſon holds the animal in one hand, and in the other a pair of open ſciſſars, betwixt which is the leg of the pigeon to be cut off. Another perſon holds this leg in one hand,

and in the other the head of a viper with the teeth laid bare, and forces theſe teeth deeply into the muſcles of the leg, the inſtant it is ſeparated from the body. The head of the viper has been ſeparated from the body ſome minutes before, and, to make the experiment more commodiouſly, has been deprived of the lower jaw. This head is ſtill alive, and the ſmalleſt compreſſion that can be made, is ſufficient to make it of itſelf draw its teeth from out of their bag or ſheath, and force them into the parts that are made to approach it.

It is certain that there never paſſed, in any one of the dozen experiments I firſt made, more than three ſeconds betwixt the amputation and the bite; ſeveral of theſe experiments were made in a ſingle ſecond, or preciſely at the very inſtant of cutting off the limb.

In ſome of the legs that were cut off, the venom was ſeen ſurrounding the holes made by the teeth; in others it was ſeen flowing out of the holes; and in others no venom was to be perceived exteriourly. On examining the muſcles bit in this way by the viper, I could diſcover no ſign of a communicated diſeaſe, neither could I obſerve any ſupervening lividity about the holes. The blood continued in a fluid ſtate in the veins and arteries.

Theſe legs, which were yet warm and palpitating, and which bled, being kept for minutes, and even for hours, afforded me nothing further that was worthy of obſervation.

I re-

I repeated this experiment on the bared, and almoſt pale and tranſparent, muſcles of twelve frogs. The event was exactly the ſame; there was not the ſmalleſt apparent ſymptom of communicated diſeaſe.

I repeated theſe experiments afreſh, as well on pigeons as frogs, having the amputated legs bit by freſh vipers, previouſly well irritated. The reſult was the ſame in all.

I got ready the legs of pigeons and frogs, and as ſoon as they were cut off, wounded them with teeth taken from the head of a dried viper. The ſymptoms that reſulted from theſe ſimple mechanical wounds, were not ſenſibly different from thoſe of the wounds into which the venom had been introduced, although made within the ſame time.

It ſeems then to be an eſtabliſhed truth, that the venom of the viper produces no ſenſible change in parts ſeparated from an animal, notwithſtanding they are yet in a ſtate of palpitation. This truth appears to me of the higheſt importance in eſtabliſhing the theory of the venom, and deſerving of the utmoſt attention.

In the firſt place, it is certain, as I have particularly aſſured myſelf, that there ſtill ſubſiſts in the amputated leg, for upwards of twenty ſeconds, the degree of heat it had before it was cut off. A perfect irritability is ſtill retained in the muſcles, which continue to move, even for whole minutes.

The arterial and venous fluids ſtill remain in the part, at leaſt in a great meaſure, and they ſtill keep in motion there during ſome time.

Thoſe who have examined the circulation of the blood in cold adimals, know that this fluid ſtill continues to circulate for a long time, in the parts of theſe animals that have been cut off.

Notwithſtanding this, the venom ſeems to be entirely inactive and innocent, in all the caſes I have related above, although every thing ſubſiſts in the part bitten; that is to ſay, humours, arteries, veins, nerves, irritability, and motion.

This circumſtance appeared to me ſo new, and at the ſame time ſo paradoxical, that I was deſirous of trying a new kind of experiments, in which the amputated part of the animal ſhould approach ſtill nearer to its natural ſtate, at the moment of being bit by the viper.

I divided the muſcles, nerves, and blood-veſſels, of a pigeon's leg with a ſharp knife, and left the bone untouched. The inciſion was made at the beginning of the *tibia*, immediately below the *femur*. At the ſame inſtant, I had the fleſh beneath the inciſion bit by a viper.

But in ſpite of all this precaution, I could not perceive, either that the muſcles were rendered livid, or that the diſeaſe of the venom had been communicated to them.

I repeated this experiment on eleven other pigeons, and, although I kept them alive a great while

while, could never obſerve any ſymptom which indicated in the ſmalleſt degree, that they would have died in conſequence of the operation.

We may therefore regard this circumſtance, however paradoxical it may appear, as beyond all kind of doubt. On obſerving it, I began to flatter myſelf that ſome truth in phyſicks, relating to the mechaniſm of the venom of the viper, might be drawn from it; and that we might likewiſe gather from it ſome principle that would be uſeful to the comprehending of animal motions. In the firſt place it is certain, that the venom, as far as can be obſerved, does not act by a ſimple mechanical motion, or by a ſimple mixture of fluids; ſince, if that were the caſe, as the muſcles were provided with both the accuſtomed humours and motions, it ought to have produced its ordinary effects in the inſtances related above. Neither does its action ſeem to depend on an effect in chemiſtry, ſuch as is brought about, for example, by the contact of an acid with an alkali; and preciſely for this reaſon, that no effect is produced, although the venom is in contact with the humours of the leg of the animal.

Experiments to determine the Time the Venom of the Viper requires to produce its Effects, after it is introduced into a Wound.

To have excluded an hypotheſis of any kind on the mode of action of the venom of the viper, may without doubt be a ſtep towards the truth; this is not, however, ſufficient to inſtruct us how, and on what part of an animal, it acts. My curioſity was therefore rather excited than ſatisfied, and I immediately began to conſider how I ought to purſue my experiments.

I reflected that, if the venom of the viper produces no effect on a detached part of an animal, however near it may be to its natural ſtate, it is certain that it produces very violent, and very ſudden effects, on parts that have not been yet ſeparated.

The firſt enquiry that naturally preſented itſelf, was to know whether this venom produces its uſual effects, or rather, whether it communicates its diſeaſe to the part bit, at the inſtant, or not till the end of a certain time.

With this view, I enraged a large viper, and made it bite the leg of a pigeon twice, the ſecond bite inſtantly ſucceeding the firſt. I immediately cut off the leg, and examined it with attention. It was very eaſy to diſtinguiſh in it the holes made by the teeth; but although I kept it a great while, I could never diſcover any mark of diſeaſe or lividneſs.

I had

I had ſix other pigeons bit in the ſame way, each repeatedly by a ſingle viper, and almoſt immediately after cut off the leg that had been bit. There was very little difference in the time of my doing this, to all the ſix. As there appeared no ſymptom of diſeaſe in the part, it follows as an inconteſtible truth, that the venom of the viper does not act inſtantly on the part that has received the bite, but that it requires a certain time for this purpoſe; ſince it is well known, that the parts wounded by this animal, ultimately become livid and ſwelled.

The ſpace of time it requires to act, was to be determined by experiment.

For this purpoſe I had a dozen pigeons bit, each once by a diſtinct viper. I meaſured with a watch the ſeconds that paſſed betwixt the bite of the viper, and the ſucceeding amputatiop, and managed in ſuch a way, that the intervals of time increaſed in a ratio of ten ſeconds; ſo that the legs were cut off at the end of 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, ſeconds after they had been bit. I had previouſly ſtripped the ſkin from the muſcles, without cutting or lacerating them; and wiped away the blood that flowed from them after the inciſion, with a wet ſponge. In the leg of ten ſeconds, I could perceive no change, nor any livid ſpot; but in that of twenty there were ſymptoms of diſeaſe. I conceived, at leaſt, that I ſaw an incipient lividneſs about the holes made by the teeth of the viper. In all the others, the diſeaſe of the

venom was ſo decidedly apparent, that not the ſmalleſt doubt could remain on the occaſion.

I repeated this experiment on twelve other pigeons; but inſtead of making the intervals of time betwixt the cutting off of the legs in an increaſed ratio of ten ſeconds, I made them in an increaſed ratio of ſeven.

The leg cut off after ſeven ſeconds had no appearance of diſeaſe. That of fourteen was in the ſame found ſtate; but all the others, beginning at that of twenty-one, had marks of lividneſs about them. The livid ſpots were in general greater in proportion to the delay that was obſerved in the amputation. This rule was, however, not without ſome exceptions, occaſioned by a great variety in the circumſtances, which, as any one may readily conceive, are never exactly the ſame.

To obtain a more preciſe information of the time in which the diſeaſe is communicated, I had twelve other pigeons bit, and in cutting off their legs obſerved a ratio of from five to ſix ſeconds, beginning with five.

There was ſome doubt in that of twenty ſeconds, but the diſeaſe was certain in that of twenty-five. Thoſe of five, ten, and fifteen, were without any marks of diſeaſe, or the ſmalleſt livid appearance.

A certain concluſion may, I think, be drawn from theſe repeated experiments, that the action of the venom of the viper on the part bitten is not inſtantaneous, but that it requires a certain time for its effects to become ſenſible in that part.

The ſpace of time that elapſes before the venom gives manifeſt tokens of the diſeaſe it produces, is from fifteen to twenty ſeconds, or thereabouts.

We muſt naturally conceive that this time varies in different animals, and that the diſeaſe diſcovers itſelf ſooner in ſome, and later in others. The peculiar conſtitution of the animal, and its ſize, ought to make a ſenſible variation, and to modify in a greater or leſs degree, the action of the venom of the viper.

But it is ſufficient for us to know, that this venom does not operate inſtantaneouſly, and to be in ſome meaſure acquainted with the time it requires in acting on certain ſpecies of animals. Theſe *data* open the way to further reſearches.

Is it by the ſimple local Diſeaſe, or by a Diſorder excited in ſome of the moſt eſſential Principles of Life, that the Death of the Animals bit by the Viper is occaſioned?

The firſt enquiry that preſents itſelf, and which is very important, is to know whether the venom of the viper produces a diſeaſe, independant of that which diſcovers itſelf in the part of the animal that has been bit; that is to ſay, whether it deranges the animal economy in ſuch a way, after a bite has been received in any particular part, that the animal may die in conſequence of ſuch a derangement alone.

I have ſeen animals, even pretty large ones, ſuch as dogs, fall proſtrate on being bit by a viper, without

out being able to ſtir for ſome time, and with a ſcarcely ſenſible reſpiration. I have ſeen others void their urine and excrements at the very inſtant, as if their ſphincters had become paralytick at the moment of their being bit. It is not a rare caſe to obſerve men fall into a ſwoon almoſt immediately after they have received a bite from a viper. But the agitation of certain animals, and the fear of others, may contribute a good deal to the producing of theſe effects; and ſince it is invariably the caſe that there is ſtill a communication of organs, and a continuation of humours, betwixt the animal and the part that has been bit, we may miſtake for a communication of diſeaſe, what is no more than a ſimple correſpondence betwixt the part bit, and the other parts of the animal. After all, it muſt be left to experiment to decide on this, as well as on every other point.

I had a pigeon's leg bit repeatedly by a viper, and cut off the part ſoon after at one blow, at the articulation of the femur with the tibia.

The leg, when cut off, had all the ſymptoms of diſeaſe; the holes made by the teeth of the viper were livid, and the uſual ſmall ſpots were diſtinguiſhed. The pigeon died at the end of four minutes.

I had remarked, in making the experiments related above, that the amputation of the leg is not mortal to pigeons; at leaſt, I found ſeveral that were deprived of that part, ſtill living at the end of ſeveral hours.

To

To prevent the following experiments from being in the leaſt equivocal, I cut off in the firſt place the legs of ſix pigeons, a leg from each, to ſerve by way of compariſon to the others.

I had twelve pigeons bit ſucceſſively, ſome once, others ſeveral times. Betwixt the bite and amputation, there did not elapſe in any one of them, leſs than one minute, and more than two. All the pigeons died, and the times of their death are expreſſed by the following numbers, denoting ſo many minutes, 2, 2, 3, 4, 4, 4, 7, 7, 10, 12, 12, 14.

Of the ſix pigeons mentioned above, the legs of which I had cut off without having them bit, neither died, nor did either of them appear to have ſuffered in any ſenſible degree. I let them live eight days, during which time they fed as uſual, and they then ſerved me for other purpoſes.

Theſe firſt experiments ſhow, and that in an unqueſtionable way, that a mortal diſeaſe is communicated to the animal in a very little time; and that it dies, independently of the local diſeaſe, by an interiour derangement, which the venom has already communicated to its whole ſyſtem.

This new diſcovery was of too much importance not to require ſtill further experiments.

I had twenty-four pigeons bit by as many vipers, and at the end of a minute, or with very little variation, if any, from that time, cut from each of them the leg that had been bit. They all died, at the times expreſſed by the following numbers, denoting

ing ſo many minutes, 3, 3, 3, 4, 4, 5, 5, 6, 7, 7, 7, 9, 9, 10, 10, 10, 10, 10, 12, 12, 13, 13, 14, 20.

It is certain, as I have ſince aſſured myſelf by freſh experiments, that the amputation of the leg is not only not mortal to pigeons, but that it does not ſeem to be productive in them of any kind of complaint. It is equally certain, as we ſee by the experiments related above, that the pigeons bit in the leg by the viper die, notwithſtanding the part is removed, provided the amputation is delayed till the end of a certain time. It is therefore a demonſtrated truth, that the venom of the viper excites in an animal that has been bit, a diſeaſe independent of the part bitten; and that the animal dies of this ſecond diſeaſe, and not of the local diſeaſe of the leg; ſince the latter ſubſiſts no longer when the part is cut off, which does not however prevent the death of the animal. This at leaſt has unqueſtionably been the caſe in the pigeons on which the above experiments have been made. But what is ſtill more extraordinary, is that theſe animals die much ſooner when the venomed leg has been removed, than when it has not. We have already ſeen, that in pigeons the ſimple amputation of the leg is of no conſequence, and it is therefore very ſurprizing that the local diſeaſe, which becomes extremely violent, being removed, this circumſtance, inſtead of retarding the death of the animal, rather accelerates it. This would lead one to ſuſpect that the part bitten ſerves to divert the vitiated humours in the animal, and that it is, if I may ſo expreſs myſelf, a diſeaſe excited by the animal

animal itſelf, or rather by that principle which exiſts in a living animal, and which, agreeably to the opinions of Hippocrates and Sydenham, ſeems to preſide over its life, and to be the moderator of it.

Is the internal Derangement which the Venom of the Viper cauſes in Animals that are bit, produced at the Inſtant of the Bite, or ſome Time after?

What is now of the greateſt conſequence to be known, is whether the diſeaſe of the venom is communicated to the animal inſtantly, or not, on the introduction of the venom itſelf.

We have already ſeen what the local malady is, and what are the ſymptoms of it; the time has likewiſe been determined that the venom requires to produce any ſenſible effect on the part bitten. The internal diſeaſe is that which becomes univerſal in the animal, and which even occaſion its death, independently of the external and local diſeaſe juſt mentioned.

To determine whether the internal diſeaſe is inſtantaneous, or not, I made the following experiments.

I had a dozen pigeons bit in the leg by as many vipers, and cut off the part immediately after, in each of them, at a ſingle blow. There was not more than three or four ſeconds betwixt the bite,

and

and the amputation. Neither of the pigeons died, nor had any ſymptom of diſeaſe.

I repeated this experiment on twelve other pigeons, which were likewiſe bit and mutilated within the ſpace of three or four ſeconds. Neither of them died, nor had the leaſt apparent illneſs.

It is therefore certain that the venom of the viper does not produce the internal diſeaſe inſtantaneouſly, but that it requires a certain time to communicate itſelf to the animal. We are now to enquire what that time is. Is it the ſame as that which it requires to produce the external diſeaſe?—If this is the caſe, by what common principle do theſe two effects go hand in hand together? And why may not the external diſeaſe be anteriour to the internal one? The venom begins by touching the local part, and previouſly mixes with the humours of that part.

But let us proceed to experiment.——I had a dozen pigeons bit, each once in the leg by a diſtinct viper, and cut the leg from each, with an interval of five ſeconds betwixt the reſpective amputations. The firſt leg was taken off at the end of five ſeconds. The others at the times expreſſed in ſeconds by the following numbers, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60.

That of ſixty ſeconds died at the end of ſeven minutes; that of fifty-five at the end of ſix; that of fifty at the end of ſeven; that of forty-five at the end of ſix; that of forty at the end of twenty; that of thirty-five at the end of an hour; that of thirty at the end of three hours; and that

of

of twenty-five at the end of ten hours. Thoſe of twenty, fifteen, ten, and five ſeconds, neither died, nor ſeemed to ſuffer in any ſenſible degree.

However irregular the time of death in theſe animals may appear, we, notwithſtanding, remark, in one particular ſenſe, a degree of regularity. Neither of the pigeons died on which the amputation had been made before twenty-five ſeconds; and neither of thoſe recovered, the legs of which had been cut off, on or after twenty-five ſeconds.

We likewiſe obſerve, that in general the pigeons on which the amputation was made the lateſt, were thoſe that the ſooneſt fell victims to the diſeaſe of the venom.

I was deſirous of repeating this experiment of twelve other pigeons, obſerving the ſame intervals of time. The conſequences were it is true ſomewhat different; but there was ſtill a great regularity betwixt the time of the amputations, and that of the deaths.

The pigeons on which the operation was made at the end of 5, 10, 15, ſeconds, recovered. That of twenty died at the end of ſeven minutes, and that of twenty-five ſurvived. Thoſe of thirty, thirty-five, forty, forty-five, fifty, fifty-five, and ſixty, all died; and the times of their death, beginning with that of ſixty, and going back, are 5, 10, 7, 7, 6, 40, minutes, and eight hours.

Here again we obſerve, that neither of the pigeons the leg of which was amputated before twenty ſeconds, died; and that only one of thoſe

lived

lived in which the operation was performed at twenty ſeconds, or afterwards. They in general died the ſooner, in proportion to the delay obſerved in the cutting off of the leg.

The pigeon that died, notwithſtanding it was mutilated as ſoon as twenty ſeconds; made me ſuſpect (as in the former caſes, neither of thoſe that had been ſubmitted to the operation at this period died) that the ſize of the viper, and ſtill more the circumſtance of its having been irritated, might, partly, however, have occaſioned this difference.

To be certain of this, I had two pigeons, perfectly alike in ſize, bit, one of them by a large well-enraged viper, the other by a ſmall one, that was not irritated. I cut off a leg from each pigeon at the end of twenty ſeconds. The firſt died at the end of five minutes; the ſecond had not the ſmalleſt ſymptom of diſeaſe.

This experiment convinced me, that the time in which the internal diſeaſe is communicated, may be greater or leſs, according to the different circumſtances, that the vipers, and the pigeons or other animals, may be in at the time, and according to the manner of biting.

To aſſure myſelf ſtill more fully of this circumſtance, I had two other pigeons bit, one by a very large viper, the other by a very ſmall one. The firſt was enraged, and hiſſed at the time of biting. The other was made to bite, without being provoked in the leaſt. The amputation of the leg was made in both of them at the end of fifteen ſeconds.

ſeconds. The firſt pigeon died at the end of nine minutes; the other had not the ſmalleſt complaint.

It follows from all that has been obſerved, that it requires a certain time for the venom of the viper to be communicated to an animal, and that this time is ſomewhere betwixt fifteen and twenty ſeconds.

It has been ſeen above, that it requires pretty much the ſame time for the external diſeaſe to be communicated to the part bitten; and hence it appears, that theſe two diſeaſes accompany each other, and that the venom produces both within the ſame ſpace of time.

This agreement of diſeaſes and effects, which has thus far appeared ſo very regular and conſtant, fully deſerved to be confirmed by a continuation of experiments, ſtill more preciſe and ſimple than the preceding ones.

Of the Symptoms which characterize the Diſeaſe.

The difficulty conſiſts in determining the death or diſeaſe of the animal by the ſymptoms that appear in the part bitten; and *vice verſa*, in determining the ſymptoms of the part bitten by the death of the animal. On one hand, theſe ſymptoms, as has been remarked before, are neither equivocal nor difficult to obſerve; and on the other, the death of

the animal, in conſequence of the introduction of the venom, is a truth eſtabliſhed by experiment.

It would be long and tireſome to enter into a detail here, of the diſtinct conſequences of the experiments, more than eighty in number, that I made with this view. It will be ſufficient for me to ſay in general, that neither of the animals on which they were made (except one indeed, the caſe of which was doubtful) died without manifeſt ſymptoms of the diſeaſe of the venom in the part bitten; and that (except in five inſtances only) I obſerved in all the others, that when the animal recovered, there was no ſymptom of the local diſeaſe of the venom. The few exceptions that occurred, which might have depended on a thouſand accidental cauſes, do not render the law that theſe two diſeaſes obſerve, nor the conſtancy with which they are at the ſame preciſe point of time excited in the animal, leſs certain.

This agreement, ſo conſtantly obſerved, made me ſuſpect ſtill more the exiſtence of a certain principle in the animal machine, which preſides and watches over life.

Scarcely has an animal encountered any thing that troubles and deranges the functions of its life, than a new force ſeems at the ſame time to be excited, and to be, as it were, awakened, which endeavours ſtrenuouſly to keep the cauſe of death from the organs that are the moſt eſſential to life, and to carry the morbifick matter to the part that is the moſt diſpoſed to receive it, whether on account

count of wounds that have previouſly been made in it, or of humours that are extravaſated by the rupture and laceration of veſſels.

The venom of the viper occupies but a very ſmall ſpace in the leg of an animal, and may, if one wiſhes it, be driven into ſo narrow a compaſs, as ſcarcely to occupy the hundredth part of a line in ſuperficies, without any phyſical or ſenſible ſolidity.

Now granting the ſuppoſition that this ſmall quantity of venom is entirely abſorbed, and carried into the torrent of the circulation, it ought to be equally diſtributed in the maſs of humours of the animal, to the ſize of which, or to its veſſels, the diſtribution of it ought to be proportioned.

But it is quite the contrary; the humours and the blood are carried tumultuouſly, and in haſte, to the part that has been bit, and the blood not only collects about the ſimple mechanical wound made by the tooth, but ſpreads to a great diſtance, and changing its colour, pours in torrents into the adipoſe membrane, whilſt another part of this fluid penetrates in a diſſolved ſtate through the coats of the veſſels.

It therefore appears, that all the efforts made by an animal which has been bit by a viper, are directed to the diſcharging of the blood and humours that are affected by the obnoxious principle the latter conveys by its venom, and to the throwing of them, as much as it can; on the part that has been bit. If it ſucceeds in this way, in ſupporting

the highly neceſſary functions in the vital parts, it ſurmounts the very ſudden and dangerous internal diſeaſe that would otherwiſe have been deſtructive to it.

As to the external diſeaſe, the circumſtances are altogether different. It becomes ſimilar to many other diſeaſes cauſed by an obſtruction of humours in the veſſels, of fluids extravaſated in the adipoſe membrane, and of blood which threatens gangrene and ſphacelus. If the animal is very ſtrong, however great the local diſeaſe may be, it at length recovers; and I have obſerved monſtrous ſwellings, enormous extravaſations, and an entire lividneſs and gangrenouſneſs of the parts, and, notwithſtanding all this, the animal has got about again. This is frequently obſerved in the larger ſpecies of animals, ſuch as reſiſt for ſeveral days the action of the venom.

I wounded the crural muſcles of three pigeons with venomous teeth, and, almoſt at the very inſtant, cut off the leg in each of them. The muſcles of the firſt pigeon's leg had no apparent ſymptom of diſeaſe. Thoſe of that of the ſecond had a ſmall red ſpot, which penetrated through the fibres without changing its colour. In thoſe of the third pigeon's there was a ſmall red ſpot, ſimilar to the former one, which penetrated to the tibia itſelf, where it appeared ſomewhat darker than uſual.

I wounded the crural muſcles of two other pigeons, with teeth which had been dried a long time, and which I had previouſly well waſhed, and a moment

a moment after cut off the leg in each of them that had been thus punctured. In one of theſe legs there was no ſymptom of diſeaſe or wound; in the other there were two red ſpots, which penetrated into the muſcles, inſenſibly loſing their redneſs.

I wounded the crural muſcles of three other pigeons with venomous teeth, and bound and cut off the legs at the very inſtant. In one of theſe legs there was an appearance of black and extravaſated blood. The ſymptoms of diſeaſe in the other two were perfectly viſible and certain; that is to ſay, a livid colour, and black and extravaſated blood for the whole depth of the muſcle.

I wounded the crural muſcles of two other pigeons with dried teeth, and at the ſame time bound and cut off the legs. The blood was extravaſated in both, and was become of a dark colour.

Experiments to determine whether at the Moment of Amputation, a ſubtile Principle of ſome Kind does not eſcape from the Blood.

The little conſtancy I met with in theſe experiments, and the ſuſpicion that a volatile fluid of ſome kind might have eſcaped from the blood on the moment of its being diſcharged from the veſſels, and expoſed to the open air, induced me to engage in ſome other trials, which I conducted in the following manner:—I held the pigeons in ſuch a way, that although their legs were perfectly dry, their

thighs were entirely plunged in water. The amputation was made beneath the water; in the thigh, ſo that the inciſed part could have no communication with the air; and the muſcles were wounded under water with venomous teeth. This being done, I kept the foot under water for three or four minutes, drew it out again, and examined it.

I repeated this experiment on the ſame number of pigeons, and ſimply wounded their muſcles with dried teeth. There were marks of the ſimple mechanical wounds, not only in the venomed muſcles of the pigeons in the former experiment, but likewiſe in thoſe that had not been venomed of this one. As I found no difference betwixt them, I cannot take upon me, with any degree of probability, to eſtabliſh a fact of any importance on theſe appearances.

I examined ſeveral times into the parts adjacent to the one that had been bit, either in animals which were already recovered, or in thoſe in which there were no longer any certain ſymptoms of diſeaſe, and of which the parts had almoſt regained their uſual motions. I obſerved with ſurpriſe in ſeveral of theſe animals that had been bit in the leg, that there was ſtill a great extravaſation of humours in the adipoſe membrane, at a very great diſtance from the part that had been bit; and likewiſe that all the abdominal muſcles were ſtill red and inflamed. Every thing in ſhort concurred to perſuade me of the exiſtence of that principle, which

which has been either ſuſpected or admitted by others; and to convince me that the local diſeaſe is not a mechanical effect of the introduction of the venom into the part, but is rather the means this vital principle employs to drive towards the exterior parts the morbifick matter that circulates in the humours, and to relieve from it the organs that are the moſt eſſential to the preſervation of the animal. I ſhall point out at the concluſion of this work, the purpoſe that may be drawn, and the utility that may be derived, from this diſtinction of the two diſeaſes that the viper occaſions in an animal by its bite. The want of attention to theſe two ſtates of the animal, ſo different from each other, has thrown the greateſt perplexity on this ſubject, and has enveloped it in errour and obſcurity. That which belonged to the one has been aſcribed to the other, and thus has every thing been confounded.

CHAPTER III.

On the Action of the Venom of the Viper upon the Blood of Animals.

IF the matter of the preceding chapter has been of ſome importance, which cannot be denied; if it has preſented new and altogether unexpected phenomena; if it has been a guide to us in eſtabliſhing principles and vital powers in the living machine; the ſubject of the following chapters will certainly not be leſs important, whether we regard the novelty of their contents, or the uſe and applications that may be made of them, in obtaining a knowledge of the venoms that are analogous to that of the viper, and in explaining the animal mechaniſm, as well in a ſtate of diſeaſe as in perfect health.

Mead, to determine whether the venom of the viper had any degree of action on the blood of an animal that had been bit, mixed five or ſix drops of it with half an ounce of blood, in the colour and conſiſtence of which he could obſerve no change, as the conſequence. There was in ſhort no difference betwixt this blood and another quantity drawn at the ſame time, which he had put into a veſſel ſimilar to that which contained the firſt, by way of comparing the two together. This experiment

ment I repeated, and received the blood which flowed from the divided veſſels of an animal, immediately into a concave glaſs, which I had previouſly warmed, and into which I had put five grains in weight of the venom of the viper. The paſſage of the blood from the veſſels to the glaſs was ſo quick, that it is not poſſible to have it out of the veſſels, in a condition approaching nearer to its natural one. On the moment of the union of the venom with the blood, I obſerved the latter, the quantity of which was about an ounce, or ſomewhat more, with a very ſtrong lens. I could never perceive any kind of motion in it, neither could I diſtinguiſh in it any diſſolution, nor the ſmalleſt appearance of coagulum; in a word, it was entirely in its natural ſtate. Its globules were of their uſual ſhape, and its colour was equally preſerved. This particular ought not to ſurprize us, after the experiments that have been made on the legs of pigeons bit by the viper at the very juncture of their being cut off; and likewiſe after thoſe in which they were cut off ſometime after they had been bit. The blood in theſe caſes certainly approaches much nearer to its natural ſtate than when it is drawn from the veſſels. There is here both the natural heat and ordinary motion of the humours, and in ſhort the life of the organs themſelves.

Nothing appears more natural than the deducing from theſe particulars, that the venom of the viper has no action on the blood of the animal that receives the bite. This is indeed the inference that

Mead

Mead has drawn from the above recited experiments on the blood of animals taken warm from the veſſels.

However perſuaſive this experiment on the blood might have been, and however reſpectable the authority of Mead, I determined to try a new kind of experiments, partly analogous to thoſe related above, but more direct and more ſimple. Theſe experiments conſiſt in introducing the venom of the viper in an immediate way, without touching any of the parts that are previouſly cut, into the blood. They are indeed ſomewhat difficult, but are ſtill poſſible, and are made by injecting the venom of the viper, by means of a ſmall glaſs ſyringe, into a vein that has been opened with a lancet. I foreſee an objection that will be made, that experiments of this kind are altogether uſeleſs after thoſe that have been related, to which they are beſides perfectly analogous; and that ſeeing there has been no change obſerved in the venomed blood, it ought, from a parity of reaſoning, to be concluded, that there will be no greater change in it, in theſe experiments. Such is the riſk of being miſtaken, that thoſe incur who love rather to reaſon than to experiment; and this is the mode of arguing of thoſe philoſophers, who, perſuaded that they are arrived at the fountain-head of natural ſciences, flatter themſelves that they know every thing, and are capable of explaining every thing.

Injection

Injection of Venom into the Blood Veſſels; and its Effects.

The experiments I am about to relate, were made on large rabbits. The jugular vein was the veſſel on which I operated.

When a great portion of hair has been removed from the inferiour part of the ſide of a rabbit's neck, and a large inciſion made in the ſkin, the jugular vein is diſcovered dividing itſelf into two ſmaller branches. I ſtrip in the experiment, the two branches and a part of the trunk of the jugular vein, of about ten or twelve lines at leaſt in length, of the adipoſe membrane, and the other neighbouring parts. I tie one of theſe branches with a thread, at the diſtance of ten lines from the trunk, and tie another thread to the ſame branch, about ſeven lines below the firſt, ſo that this ſecond thread is only three lines from the trunk. This laſt thread has a knot, ready to be drawn tight at a proper time. But before I go any further, I think it neceſſary to explain the manner of making uſe of the ſmall ſyringe, intended to convey the venom into the veſſels.

This is a ſmall common glaſs ſyringe, terminating in a capillary tube of ten lines in length, and crooked. I put into this ſyringe the venom that I mean to introduce into the vein. I uſually cut off two vipers' heads, and receive all the venom from

from their veſicles in a ſmall cryſtal ſpoon. I add to this venom the ſame quantity of water, and when the liquors are well blended together, draw them up by ſuction into the ſyringe. There uſually enters into the ſyringe at the ſame time, a ſmall air-bubble, which is eaſily diſperſed, by puſhing the piſton forward a little towards the tube. The ſmall quantity of the liquor that flows with the air out of the point of the tube, is received in the ſmall ſpoon, and is ſucked up again into the ſyringe, by once more withdrawing the piſton a little.

The ſyringe being thus freed of the external air, I withdraw the piſton in an almoſt inſenſible degree. The venom retreats a little, and leaves the point of the capillary tube, which remains full of air, for the length of four lines. The quantity of this air is very trifling, on account of the ſmallneſs of the diameter of the tube in that part. I now wipe the crooked part or extremity of the ſyringe with a piece of very fine moiſtened linen, and introduce a very fine and dry linen thread, for the length of two lines, to cleanſe the venom, and likewiſe the ſmall ſpace in the capillary tube, that is occupied by the air.

The ſyringe being thus in readineſs, I raiſe a little, by the uppermoſt thread, the branch of the jugular vein to which the two threads are faſtened, betwixt which I open it with a lancet, and introduce the capillary extremity of the ſmall ſyringe at the orifice, continuing this till it has entered four or five lines into the principal trunk. I now draw the

end

ends of the threads together, the lower one of which binds the coats of the veſſel very ſtrongly to the capillary tube of the ſyringe. Things being in this ſtate, I puſh forward the piſton of the ſyringe by degrees, and force out of it all the venom, which paſſes entirely to the trunk of the jugular vein, to be carried an inſtant after to the heart.

This experiment requires two perſons at leaſt, and ſucceeds ſtill better when there are three. If the ſyringe has been previouſly got in readineſs, it does not continue at the moſt for more than two minutes altogether; and when the parts of the animal are known, and it has been made a few times, is not ſubject to any inconvenience.

Before the ſyringe is drawn out of the veſſel, I have been accuſtomed to withdraw the piſton a little, that a ſmall quantity of blood may enter the capillary tube, and that none of the venom may remain at the orifice of it. At the moment of my drawing out the ſyringe, I again tighten the lower thread, ſo that the veſſel remains perfectly cloſed. I raiſe with pincers the portion of the jugular vein betwixt the threads.

It was not without reaſon that I made choice of a veſſel which branches out into two others; neither was it at hazard that I introduced the capillary part of the ſyringe into the principal trunk itſelf.

I wiſhed that the venom ſhould be carried immediately to the heart, and I could not think of a better expedient than that of procuring a very large lateral veſſel, where the blood continuing to run in a

full ſtream towards the heart, muſt neceſſarily carry with it the venom that it meets with in the trunk.

Theſe experiments are too important not to be related with ſome degree of detail. They at leaſt require me to deſcribe the principal circumſtances by which they were accompanied. I ſhall give them here in the order in which they were made.

I injected into the outer jugular vein of a large rabbit that weighed ſeven pounds, the venom of two viper's heads, got ready in the manner deſcribed above, and with a nice obſervance of all the precautions I have juſt laid down. The venom ſcarcely began to enter the vein, when the animal gave ſeveral horrible cries, diſengaged itſelf, writhed itſelf about, and died a moment after.

The novelty of this ſtrange and unexpected event, prevented me from calculating the exact time the animal lived after the injection of the venom; neither could I aſcertain the time I employed in propelling the whole of the venom from the ſyringe. It is certain, however, that the animal did not live more than two minutes, and that the injection was made within the ſpace of eight or ten ſeconds.

As I was deſirous of ſeeing whether this experiment was a certain one, or whether the animal died in conſequence of ſome circumſtance I was ignorant of, I examined the ſtate of the viſcera in the dead animal, and likewiſe that of the blood in its veſſels. I was likewiſe induced to vary ſome of the circumſtances in making the ſucceeding ones.

I got

I got ready another rabbit in the above manner, and began by injecting a quantity of water into a branch of the jugular vein, equal to that of the mixture of venom and water in the preceding experiment. The rabbit did not ſuffer in the leaſt. I kept it in this ſtate during five or ſix minutes, and perceiving that it did not become at all diſordered, ſat about injecting into the ſame vein, the quantity of venom mentioned above.

The animal, however, neither cried out nor was agitated. At the end of a few minutes I perceived that it had ſickened, and it died at the end of twelve hours. All the parts I had ſtripped of the ſkin, to lay bare the jugular vein, were violently inflamed, and very livid. The adipoſe membrane was filled with black extravaſated blood. All the pectoral muſcles at the ſide on which I had injected the venom, and a part of the abdominal ones, were already become livid. The very inteſtines were inflamed. The inner part of the thorax was inflamed likewiſe, and was bloody; and the heart had formed adheſions. The blood, both in the large veſſels and heart was coagulated and black; and the lungs were marked here and there with ſomewhat livid ſpots.

This ſecond experiment convinced me of the very great importance of thoroughly examining the ſtate of an animal after its death. It is principally by this ſtate, that we ought to judge of the action of the venom on the blood.

But how came it about that the firſt rabbit died as it were inſtantly, and the ſecond not till the end of

of twelve hours? To what is this difference to be aſcribed?

I inſtantly proceeded to a third experiment, hoping to draw ſome further information from it.

I got ready a rabbit, and injected the venom of two vipers as before, into the branch of the jugular vein. The rabbit did not ſeem to ſuffer in the leaſt from this operation, and recovered of the external diſeaſe in a few days, as readily as if it had only undergone the preparation neceſſary to the injection of the venom. An hour after this injection had been made, I found it eating, as if in perfect health.

This third experiment completed my perplexity, and I began to miſtruſt altogether. In the firſt place I ſaw an animal die, as it were at the moment of injection, and diſtinguiſhed a real diſeaſe in that which lived twelve hours. It was therefore certain that the venom, when united with the blood, was capable of producing ſuch a derangement in the animal machine, as to excite in the animal a very violent diſeaſe, terminating in death. All this was real; but how could theſe two caſes be reconciled with the third?

Some doubts occurred to me as to the method I had purſued in making theſe experiments, which had not been altogether conducted with the exactneſs and preciſion I deſcribed a little time ago. I did not make the ſecond ligature in the vein; I did not examine whether the capillary tube reached into the principal trunk; and I did not withdraw the piſton of the ſyringe, before I drew the capillary tube

tube of the latter out of the veſſel. The neglect of theſe precautions made me look upon the three experiments I have juſt related as ſuſpicious, and I ſat about experimenting afreſh, with a greater degree of attention and preciſion than before.

For this purpoſe I got ready a large rabbit, healthy, and in good plight. I made the two ligatures in the external branch of the jugular vein, and introduced the capillary tube into the common trunk, tightening the thread on the tube, and injecting the whole of the liquor at once. I took care to withdraw the piſton a little, before I drew the ſyringe out of the vein, and to tighten the thread once more. In a word, I did not neglect any one of the precautions I had previouſly determined to take. The conſequences were as follows.

The whole of the venom had ſcarcely paſſed from the ſyringe into the jugular vein, when the rabbit gave ſeveral horrid ſhrieks, and was ſeized with very violent convulſions. It died in leſs than a minute and an half. There were not more than ſeven ſeconds ſpent in the injection.

The blood in all the large veſſels was black and coagulated. It was likewiſe ſo in the heart and auricles. The coronary veins were ſwelled and livid; and an extravaſated black blood, in large ſpots, was ſeen about them in the muſcular ſubſtance of the heart. The pericardium was entirely filled with a liquor, ſo as to be diſtended like a bladder; and this liquor was tranſparent, with a ſlight red tinge.

The lungs were full of the uſual ſpots, through which, when they were touched in the ſlighteſt degree, the air ruſhed out of the water that covered this viſcus. The inteſtines, ſtomach, and meſentery, were covered with ſmall, livid, and red ſpots.

This experiment ſucceeded too well to leave me in any kind of doubt as to the nature of its conſequences. The animal died almoſt inſtantly, and cried out the moment the venom had entered the veſſel.

The two viſcera that are the principal organs of life, were inſtantly attacked by a violent and mortal diſeaſe. The blood was immediately coagulated in the large veſſels, in the lungs, and in the heart. In a word, every thing concurred to the ſudden ſtoppage of the circulation, and to the death of the animal.

The extravaſation of the blood of the coronary veins is ſurprizing, and the livid ſpots of the lungs, and dilacerations of this viſcus ſtill more ſo. But what ſuprized me moſt of all, was the blood collecting in ſuch an abundance, in ſo many veſſels, and in ſo many cavities. In this diſeaſe an extreme diſſolution of a part of this humour, exuding every where through the veſſels, takes place; and, at the ſame time, a coagulation of another part, which fixes and condenſes in a few moments.

Every advance I made, in this new career of experiments, preſented me either with ſomething paradoxical, or with a novel and unexpected circumſtance. I paſſed on to the fifth experiment, which

I made

I made exactly as I had done the fourth. Although the reſult of it was ſomewhat different from that of the laſt, it agreed very well as to the nature of the diſeaſe, and as to the opinion that ought to be held of the introduction of the venom of the viper into the blood. On the injection being made, the rabbit did not cry out, neither did it ſeem to ſuffer in any ſenſible degree. At the end of an hour, however, it appeared to be ſick, refuſed its food, and died at the end of twenty-four.

On opening its body, I did not find the abdominal viſcera to be much inflamed; but to atone for this, the uſual livid ſpots, and the air guſhing freely out of them, were ſeen on the lungs. All the muſcles of the breaſt were conſiderably inflamed, and the whole of the adipoſe membrane, from the neck to the lower part of the belly, was filled with black, extravaſated, and fluid blood. The blood in the heart, in the lungs, and in the large venous veſſels, was coagulated; but much leſs ſo than in the caſes related above in which the rabbit died almoſt inſtantaneouſly.

I immediately proceeded to the ſixth experiment, to ſee whether there would be any degree of uniformity betwixt the injection of the venom, and the death of the animal. I neglected to remark, in relating the preceding experiments, that I had found ſometimes a greater, ſometimes a leſs, quantity of venom in the viper's heads, and that in ſome of them I had even obſerved a white and ſomewhat glutinous matter flow from the tooth.

I had likewiſe found the palate of ſome of the vipers I employed inflamed to a certain degree, and the two ſheaths or bags of the teeth likewiſe inflamed and red.

But I could not ſay poſitively, whether theſe circumſtances could have been capable of influencing the effects of the venom on the animal. I therefore reſolved to take it in future from no other heads of vipers, but ſuch as were perfectly ſound, and the beſt ſupplied; and to procure it in a greater quantity.

I got ready a large and ſtrong rabbit in the uſual way, and introduced into the ſyringe the venom of two very large vipers, the heads of which were in a ſound ſtate.

The venom was not yet completely injected, when the rabbit began to ſhriek, and died in leſs than two minutes in very violent convulſions. Having opened the breaſt, I found the auricles and ventricles filled with grumous blood. That of the large venous veſſels was in the ſame ſtate. There was a great deal of lymph in the pericardium, in which there was likewiſe extravaſated and concreted blood. All the inteſtines were in a very inflammatory ſtate, as were alſo the ſtomach and meſentery. The arteries were in general empty. The lungs were but little ſpotted; but in inflating them beneath the water, the air was ſeen ruſhing out in ſeveral parts, and the ſmall ſpots were then apparent. The blood in the lungs was likewiſe concreted.

I got

I got ready another rabbit, and injected in the uſual way into the jugular vein, the cuſtomary quantity of venom.

It ſcarcely began to enter the vein, when the rabbit cried out, and in leſs than two minutes died, with the moſt terrible ſhrieks and convulſions.

I opened it, and found the lungs ſpotted as uſual, and the blood coagulated in the two ventricles. It was much more ſo in the right ventricle than in the left, as I had alſo found it in all the preceding caſes. It was likewiſe in the ſame ſtate in the auricles and veins. The pericardium was filled with water mixed with blood. The coronary veins had, for their whole circumference, two large, longitudinal, and livid ſpots. The blood in the lungs was black and grumous, and the air guſhed out as uſual. The inteſtines were inflamed; as were alſo all the abdominal muſcles; and there was a great deal of extravaſated and diſſolved blood in the adipoſe membrane.

Theſe two laſt caſes are very uniform, and agree too well with the preceding ones, to admit of any doubt as to the immediate action of the venom of the viper on the blood.

Further Experiments on the Jugular Vein of Rabbits.

Notwithſtanding the uncertainty and obſtacles that are met with in experimenting on the blood-veſſels, I was deſirous of making ſome further trials

on them, conducted with all poſſible care and attention, as the very great importance of the ſubject ſeemed to require. I choſe for this purpoſe, two of the largeſt rabbits I could procure, each of them weighing ten pounds. I took the venom from two ſound vipers, which I had previouſly examined with great attention for that purpoſe. I had not yet finiſhed the injection in either of the two rabbits, when they gave ſeveral loud ſhrieks, and died in the moſt violent convulſions in leſs than two minutes. Having opened the thorax of each rabbit, I found the lungs ſpotted as uſual, and the blood veſſels and auricles filled with black and grumous blood. The pericardium, as in the former caſes, contained a humour, and the inteſtines and muſcles were, as uſual, inflamed.

The immediate action of the venom of the viper on the circulation of animals with warm blood, is therefore both indubitable and conſtant. It is a fact, however, that would not have gained any degree of credit, had it not been for theſe laſt experiments, ſince it ſeemed in ſome meaſure to be contradicted by the former ones, which, although they were leſs direct and leſs ſimple, were nevertheleſs made on the blood. This ſhows us how cautious we ought to be in the inferences we draw from experiments; and at the ſame time proves to us, that we know little or nothing, at leaſt with any certainty, and without incurring the riſk of being deceived, beyond that which is demonſtrated to us by experiment alone.

But

But in what way are we now to reconcile the immediate action of the venom of the viper on the blood, when it is injected by the veins; and the inactivity of this ſame venom, not only on the parts of an animal recently cut off, but likewiſe on thoſe that have remained in an entire ſtate, and united with the animal, during a period of 15 or 20 ſeconds, after it has been introduced?

I muſt acknowledge that this is a very great difficulty, and that it would be no eaſy taſk to explain it in a ſatisfactory way. It appears that there can be nothing deficient in the parts that are ſtill connected with the animal when they are bit by the viper. It even ſeems probable that theſe caſes have an advantage over thoſe in which the trial is made on the blood, ſince both the muſcular fibres and nerves are wounded by the viper's teeth, inſtead of which the venom, when injected into the veſſels, certainly touches neither one nor the other of theſe parts. What is then the cauſe that retards the diſeaſe of the venom for ſeveral ſeconds in the part of an animal that has been bit; and how it is that the venom does not produce any diſeaſe in the parts that are either cut off and bit immediately after, or are cut off immediately after they are bit.

It is probable that there may be an unknown principle in the blood circulating in the veſſels, which ceaſes to exiſt the moment this fluid is drawn out of them, and which is likewiſe no longer to be found in the parts that are recently ſeparated from the body of an animal. This principle, granting

the ſuppoſition, therefore, poſſeſſes ſuch very active and ſubtile qualities, that it is diſſipated at the very moment the part is ſeparated from the animal.

We have ſeen that the venom ſcarcely comes in contact with the blood in a veſſel, when the moſt violent derangements are produced. The animal ſuffers extremely, and the blood is condenſed in an inſtant. If this ſame venom is mixed with the blood as it flows warm from a veſſel that has been opened; or if it is introduced into any part of a muſcle that has been ſeparated an inſtant before; it produces no effect, and no appearance of diſeaſe is obſerved, nor any condenſation of humours. Here, however, every thing is the ſame, unleſs it be that in the caſe in which the venom is introduced into the veſſel, there is a blood circulating with the reſt of the humours, and always covered by the coats of the veſſels; inſtead of which, the blood that is drawn from a vein is out of the torrent of the circulation; and that of the parts which have been recently cut off has already been in contact with the air, and the veſſels which contain it are open.—However it may be, ſince the effects are very different from each other, the circumſtances muſt be ſo likewiſe; and the only concluſion we can draw, as to the humour contained in a veſſel, and the humour drawn out of a veſſel, is, that there exiſts in the firſt caſe, ſomething that is not to be met with in the ſecond.

Agreeably

Agreeably to this hypotheſis, this new principle which exiſts or reſides in the blood, in the veſſels of a living animal, does not produce the ſame effects every where equally, and in the ſame time. The venom has no ſooner united itſelf with the blood in the jugular vein, than the animal is attacked by a very violent diſeaſe, and the blood is coagulated a very few inſtants after. Inſtead of which, in the parts that are more diſtant from the heart, where the veſſels are ſmaller, it requires a certain time for the diſeaſe to diſcover itſelf, and for any ſenſible change to take place in the part into which the venom has been introduced.

It thererefore appears, that this principle obſerves certain laws in governing the animal economy, and that it is itſelf ſubject to certain regulations.

In the caſes in which the diſeaſe is the moſt remote from the heart, and the leaſt dangerous, the blood coagulates by degrees, is driven back to the parts bitten, and affords time and opportunity for the efforts of nature to overcome the diſeaſe, and to preſerve the circulation in the organs the moſt neceſſary to life.

But at length what is this new principle, and what are the organs that ſecrete it, and convey it to the veins?

In this very difficult enquiry, it appeared to me that experiment could alone furniſh me with ſome light, and conduct me to ſome new truth. But where are the experiments to begin?

CHAPTER IV.

Experiments on the Nerves.

IN the long courſe of my experiments on the venom of the viper, and in collecting together the circumſtances and ideas that preſented themſelves, I never loſt ſight of the principle of ſenſation in an animal, which appeared to me to be acted on by the venom of the viper. I have in conſequence of this, judged it neceſſary to examine the nerves in which it reſides, or which are the organ and inſtrument of it.

Mead ſays, in the introduction to his work on Poiſons, that having better conſidered the nature and quality of the ſymptoms of the bite of the viper in animals, he is certain that this diſeaſe is altogether nervous, and that it is communicated by the medium of the nerves, and not of the veſſels. In conſequence of this theory, he has recourſe to the animal ſpirits, againſt which he believes that the immediate action of the venom of the viper is exerciſed. Indeed if we examine the ſymptoms that this venom produces in animals, we are eaſily led to believe that a diſeaſe of ſuch a nature belongs to the claſs of diſeaſes which the phyſicians ſtile

nervous. In the courſe of my experiments, I have ſeen a pretty large dog fall down motionleſs, the moment after it had been bit by two vipers. I at firſt thought it dead, but at length perceived ſome little remains of reſpiration, which was, however, ſo ſlight and feeble, that it could ſcarcely be diſtinguiſhed. The dog continued in this lethargick ſtate for more than half an hour. I have ſeen ſeveral others thrown by the venom into very violent convulſions. Vomiting, anxiety, and rage, occur very frequently; the motion of the heart is irregular and convulſive, and the arterial ſyſtem hard and contracted. In ſhort, they die in the midſt of the moſt unequivocal ſymptoms of ſpaſms and contractions, and, in a word, with the affections that are by the faculty termed *nervous*.

Another idea occurred to me, that perhaps an active principle, a ſubtile fluid, is ſecreted by the nerves themſelves, which, when it mixes with the blood, contrives in ſome way to animate it, to render it vital, and to maintain its fluidity. In this caſe, the action of the venom of the viper may perhaps have been directed againſt this principle itſelf; and thus we may explain, why it is that the blood, when drawn from the veſſels, and in the open air, is no longer ſuſceptible to the action of this venom.

Expe-

Experiments on the Nerves, Spinal Marrow, and Brain, of Frogs.

I opened the belly of a frog, and laid bare the crural nerves. I poured a ſmall quantity of venom on theſe nerves, taking care that it did not ſpread to the ſurrounding parts. At the end of two hours I pricked the nerves with the point of a needle, and the muſcles of the foot contracted. At the end of four hours, however, no part of the animal was ſenſible to ſtimulation. A frog intended for a compariſon, lived twelve hours, notwithſtanding I had opened the abdomen, lacerated the inteſtines, and punctured the lungs.

I repeated this experiment twice, and the event was each time pretty much the ſame. I began, however, on a little conſideration, to think the method I had adopted fallacious. It is almoſt impoſſible to prevent the venom, when applied to the nerves, from communicating to the adjacent parts. In this caſe, the diſeaſe, or death, of the frog, may be the effect of the communication of venom to the other parts of the animal, and not the conſequence of its contact with the nerve itſelf.

I changed my mode of experimenting, but ſtill employed the ſame animals.

I cut off the heads of two frogs, alike in ſize, and touched the ſpinal marrow of one of them, but not of the other, repeatedly with the venom. At the end

end of three hours the venomed frog appeared to be dead, whilſt the other continued to live, and to leap about.

I introduced a pin into the ſpinal marrow of the frog to which the venom had been applied. Its fore-legs remained motionleſs, but there was a ſlight tremulus in the feet. The heart and auricles likewiſe had a ſmall degree of motion. In an hour more every part was at reſt. The ſecond frog leaped about the chamber at the end of twenty-four hours.

I cut off the head of another frog, and introduced a drop of venom into the ſpinal marrow. At the end of an hour the frog ſcarcely gave any ſigns of life. On opening the breaſt, the heart and auricles ſeemed ſtill to preſerve ſome degree of motion, which was however perceived with difficulty. A pin introduced into the ſpinal marrow, occaſioned an almoſt imperceptible motion of the fore-legs and feet. The heart, however, on being ſtimulated, performed its oſcillations for a long time.

I cut off the head of a frog, and removed a ſmall portion of the ſpinal marrow. I introduced by the great opening of the vertebræ, a drop of venom. At the end of two hours the frog was to appearance dead. The heart ſcarcely preſerved ſome little remains of motion, which were not encreaſed by ſtimulation. A pin introduced into the ſpinal marrow was barely capable of exciting a feeble motion in ſome of the muſcles.

I cut

I cut off the head of another frog, and having removed a ſmall portion of the ſpinal marrow, introduced a drop of venom into the great foramen. At the end of three hours the frog appeared to be dead. Having opened the thorax, I remarked that the heart was ſtill irritable; a pin, however, that I introduced into the ſpinal marrow ſcarcely occaſioned a ſenſible contraction of the feet.

I repeated this experiment on two other frogs, and the reſult was pretty much the ſame as in the above experiments. The death of the frogs ſucceeded the operation and introduction of the venom in a ſpace of betwixt two and three hours. The heart was ſomewhat irritable, but the muſcles were little ſo, or not at all, notwithſtanding my ſtimulating the ſpinal marrow with a needle.

I now thought it proper to make a little variation in my experiments.

I removed a portion of the cranium of a frog, and applied a ſmall quantity of venom to the brain. At the end of four hours the frog was dead, and the heart inſenſible to every ſtimulation. On pricking the ſpinal marrow with a needle, not the ſmalleſt motion was reſtored in it.

I opened the cranium of another frog, and applied a drop of venom to the brain. The frog ſurvived this operation two hours, at the end of which time the heart had ſtill retained a ſlight degree of motion; it was ſhrivelled, black, and contracted. On ſtimulating the ſpinal marrow, there was an almoſt inſenſible contraction of the muſcles.

I re-

I repeated this experiment on the brain of four other frogs, and the conſequences were very analogous to thoſe of the two preceding ones. However having removed the cranium of the two frogs without applying the venom to the brain, by way of a comparative experiment, they both died in the ſpace of ten hours.

The conſequence of theſe experiments appearing neither ſufficiently clear nor uniform, I had once more recourſe to the cutting off of the heads, thinking that by dint of multiplying my experiments in that way, I might aſſure myſelf of the action of the venom on the nerves.

I cut off the heads of two frogs, and applied venom to the ſpinal marrow of one of them, but did not venom that of the other. At the end of three hours the venomed frog was to appearance dead; the other was ſtill living, and had a free motion in all its parts. I introduced a pin, which had been dipped in venom, into the vertebral opening of the firſt frog, and it excited a very feeble motion of the feet, but had no ſuch effect on the fore-legs. I ſcarcely ſtimulated the ſpinal marrow of the other frog with a needle, when the frog leaped about briſkly. At the end of the fourth hour there was not the ſmalleſt perceptible degree of motion in the venomed frog, and neither the heart nor auricles were any longer ſenſible to ſtimuli. The other frog was ſtill leaping about at the end of thirty hours.

I cut

I cut off the head of another frog, and introduced the venom into the ſpinal marrow. At the end of two hours the frog was to appearance dead. Having opened the thorax, the heart was motionleſs, and even inſenſible to ſtimulations. The ſpinal marrow, when likewiſe ſtimulated, ſcarcely excited any degree of motion in the feet.

I repeated this experiment, obſerving the ſame circumſtances, on another frog. At the end of three hours I found it dead. The heart and muſcles were perfectly motionleſs. On treating another frog in the ſame way, the conſequences were altogether ſimilar.

I cut off the head of another frog, and applied venom to the ſpinal marrow. At the end of five hours the frog ſtill retained ſome feeble ſigns of life. On opening the thorax, I found the heart motionleſs; it however renewed its oſcillations on being touched.

The conſequences of all theſe experiments together may reaſonably induce us to ſuſpect, that the venom of the viper acts on the nerves, and that when it is applied to theſe parts in frogs, it produces a mortal diſeaſe. But this mode of experimenting is not entirely irreproachable. The ſpinal marrow and brain are too ſmall to enable us to be certain that the venom does not communicate to the adjacent parts. No precaution whatever can in my opinion prevent this. When the venom is applied, it is too near to the veſſels and other parts; and how indeed can it be kept from the blood-veſſels of both brain and ſpinal marrow?

This

This enquiry is too important to be confined to the limits of ſimple probability. I ſtill flattered myſelf, that the purſuit of it would tend very much to the obtaining of a true knowledge of the venom of the viper and its qualities, and of the animal economy itſelf.

With this view I formed a plan of experiments, to be made on the nerves of the largeſt rabbits I could procure. This animal is hard to kill, and as it is gentle in its nature, may be managed agreeably to one's wiſh; it is likewiſe not ſo ſmall, but that its nerves may ſerve for the moſt deciſive experiments.

Experiments on the ſciatick Nerve of Rabbits.

I made choice of the ſciatick nerve as the ſubject of my principal experiments. I removed the hair with ſciſſars from the ſkin that covers the great gluteus muſcle, and made an inciſion, beginning on the great trochanter, and deſcending in a direction with the thigh. I detached the anteriour part of the gluteus muſcle from the os innominatum and trochanter, and gradually raiſed the muſcle with my fingers, freeing it from the adipoſe membrane. A little cuſtom in theſe experiments, enables one to make them in leſs than two minutes, and in ſuch a way, that after removing the ſmall quantity of blood which flows from the integuments, it is not ſucceeded by any freſh hemorrhage, that is capable

of retarding or diſturbing the operation. Now, holding the great gluteus muſcle with one of my hands, I paſſed, with the aſſiſtance of ſmall pincers, a piece of fine linen in ſeveral folds under the ſciatick nerve, upon which, being in this ſtate, I began my experiments.

Having got ready one of the ſciatick nerves of a large rabbit, in the way I have juſt deſcribed, I wounded it in ſeveral places with a venomous tooth. The rabbit ſhook itſelf a little at the time of my doing this. At the end of twenty hours it ate, and ſeemed in full vigour. It died however at the end of ſeven days, with a large wound in the part that had been cut. This experiment was not made ſo well as it ſhould have been; more than half the gluteus muſcle was cut, and there was a great hemorrhage.

I laid one of the ſciatick nerves of another rabbit perfectly bare, paſſing beneath it ſeveral folds of linen. I then wounded it, in upwards of twenty places, with the venomous teeth of two vipers. The rabbit ſcarcely gave any ſigns of feeling pain, and at the end of ten hours, ate and appeared lively. It was in this ſtate at the end of twenty-four hours, but died at the end of forty-eight. The nerve was marked here and there with dark red ſpots; the parts about it were violently inflamed; and the blood in the auricles and heart black and coagulated.

In wounding the part with the venomous teeth, I took the greateſt care imaginable to prevent the venom from communicating to the adjacent parts;

and conſtantly covered the nerve after I had wounded it.

Having got ready one of the ſciatick nerves of another rabbit, I paſſed under it the uſual folded linen, and wounded the nerve in ſeveral places with the venomous teeth of two vipers. I covered the nerve well with linen, and ſtitched up the ſkin. I had obſerved this latter precaution in the preceding experiments.

The nerve was prepared for the introduction of the venom in leſs than two minutes, and the hemorrhage from the integuments was very ſlight indeed. Notwithſtanding this the rabbit died at the end of eighteen hours. The nerve was to appearance in its natural ſtate. The blood in the heart and auricles was black and grumous. The muſcles in the vicinity of the nerve were a little inflamed, and there was a degree of lividneſs on their ſuperficies.

Theſe experiments, although few in number, and not diſtinguiſhed by any great uniformity, began however to make me ſuſpect, that the bite of the viper is leſs dangerous to the nerve, than to many other parts of an animal. The rabbits lived much longer than one would naturally have conceived, and notwithſtanding it appears that they all died ſooner or later, I conjectured that here, as in the caſes of the tendons, the venom might have communicated to the neighbouring parts, and that the animal might probably have died rather from this, than from any other cauſe.

As a ſtill further precaution in purſuing theſe experiments, I had recourſe to the piece of lead I had before made uſe of, putting it betwixt the folds of the linen. In this way the nerve was very well ſecured from the other parts, and it did not ſeem poſſible for the venom to ſpread beyond it.

I wounded a ſciatick nerve of a rabbit in ſeveral places, after having got it ready in this way, with the venomous teeth of two vipers; covering it with linen, and binding it up ſecurely afterwards. During the time employed in forcing the teeth into the nerve, the rabbit cried out ſeveral times, and was ſeized with violent convulſions. It died at the end of twenty hours. All the muſcles about the nerve were livid and ſphacelated for their whole ſubſtance; and the ſphacelus extended for the whole length of the leg. The lungs were ſpotted; and the nerve itſelf was likewiſe covered with red and livid ſpots. The blood in the auricles and great venous veſſels was black and coagulated.

The circumſtances that accompany this experiment are ſufficient to induce one to believe, that the venom of the viper has in effect a ſtrong action on the nerves. The ſphacelus of ſo many muſcles, even of thoſe that were diſtant from the wounded part, made a great impreſſion on me. I, however, did not on this account terminate my experiments.

Having laid one of the ſciatick nerves of another rabbit perfectly bare, I wrapped it carefully in the linen, in which I had not, however, encloſed the bit of lead. I now wounded it in ſeveral places with the

the teeth of two vipers, and covered it with linen as uſual. The rabbit died at the end of thirty-two hours. The nerve was but little redder than it naturally is, and was not ſpotted. The blood in the auricles and large veſſels was but ſlightly coagulated. When I opened the rabbit, I found it ſtill to poſſeſs a degree of warmth.

This experiment is very different from the preceding one, and ſhows how little confidence we ought to place in experiments themſelves, however nicely they may have been made, unleſs they are in a great number, and agree with each other.

I laid bare a ſciatick nerve of another rabbit, and wrapped it carefully in the folded linen, in which I had previouſly encloſed the bit of lead. I wounded it in ſeveral places with the venomous teeth of two vipers, and afterwards covered it ſecurely. The rabbit died at the end of thirty-two hours. Several parts of the nerve were red, and there were livid ſpots in it. The muſcles adjacent to it were in their natural ſtate; but the lungs were livid and ſpotted. The heart, auricles, and principal veſſels, were filled with black grumous blood.

I repeated in four other rabbits, the application of venom to one of the ſciatick nerves, but with ſome little change in the circumſtances. I apprehended, that probably the linen which encloſed the nerve on all ſides, and remained on the wound, might occaſion the death of the animal, and the derangements we have obſerved. It therefore became neceſſary to ſeparate theſe two circumſtances,

and to remove the linen after wounding, the nerve with the venomous teeth. Before the linen was removed, I wiped the venom carefully from the ſurface of the nerve with ſmall bruſhes, which I repeatedly changed. After this, I dipped bits of linen in water, and holding them with pincers, employed them in waſhing the whole extent of the nerve. The linen I had paſſed under the nerve, folded upwards of ten times, prevented the water from communicating to the adjacent parts. I now removed this linen, and threw on the nerve a conſiderable quantity of water, which waſhed at once, nerve, muſcles, &c. ſo that it was not poſſible for any particle of venom, however ſmall, to continue lodged in the parts that ſurrounded the nerve.

Theſe four rabbits all died in leſs than thirty-ſeven hours. In three of them no ſenſible change was to be obſerved in the parts adjacent to the venomed nerve. The muſcles, except that they were a little redder than uſual, were in their natural ſtate.

I confeſs that on one hand it did not appear poſſible to me, that the venom could, notwithſtanding all the precautions I had taken, have been communicated to the ſurrounding parts; and on the other hand, I could not find any ſymptom of diſeaſe, any effect of the venom, in the muſcles adjacent to the venomed nerve. The death of the animal was the moſt conſtant reſult in theſe experiments; and this did not however take place till very late, and was not attended with either ſpaſms or convulſions. If

the

the bite of the viper is really venomous to the nerves of animals, it is certain that it acts on these parts with less force and activity than on many others.

As this enquiry appeared to me a very important one, I thought it proper to persevere, making some little change in my experiments.

Experiments on the Sciatick Nerve, divided in its upper Part.

I laid bare the sciatick nerve of a rabbit on one side, in the usual way, and with a pair of sciſſars divided it in its upper part, as near to the vertebræ as possible. The cleared part of the sciatick nerve, towards the extremity, was about an inch and an half in length. I wrapped it in linen, which was as usual in several folds, wounded it in several places with venomous teeth, and covered it securely, to prevent the venom from communicating to the surrounding parts. The rabbit died at the end of thirty-six hours.

I opened it whilst it was yet warm. The blood in the heart and auricles was black, but not grumous. The muscles adjacent to the nerve were somewhat inflamed.

I laid bare one of the sciatick nerves of another rabbit, and divided it in the manner described above. I wrapped it in linen, wounded it with venomous teeth, and covered it. The rabbit died at the end of eighteen hours. The nerve in some parts was

dark and livid; the adjacent muſcles were in a very ſlight degree inflamed; and the blood in the heart ſtill fluid.

The principal aim of this method of experimenting, was to ſee what effect the venom of the viper would produce, when applied immediately to a nerve, leading it is true to an organized part, and one endued with ſenſation, but which had no longer any immediate communication with the life of the animal. The action of the venom in the two above caſes could not in any poſſible way be communicated from the nerve to the animal, and could not awaken in it any immediate pain or ſenſation. This however did not prevent the nerve from communicating the diſeaſe of the venom to the inferiour part in which it terminated. It muſt be obſerved, that in this part the humours continue in motion as before; that the muſcles are in their entire natural ſtate; that the fibres ſtill preſerve their irritability; and that the part retains its ſenſibility, in conſequence of the other nerves that terminate there. With all this, no diſeaſe is obſerved in it. There is no tumour, no gangrene nor ſphacelus, and no extravaſation of black and grumous blood.

Thinking, however, that two experiments alone, would not be ſufficient to render this particular, which is of ſo much importance, certain, I was deſirous of repeating them in the ſame way.

I deſtined ſix rabbits to this purpoſe, and in each of them laid bare and divided one of the ſciatick nerves, wounding it as uſual with the venomous teeth,

teeth, and covering it carefully with linen. Theſe rabbits all died, two in eighteen hours, the other four before the expiration of thirty-ſix. The adjacent muſcles were in their natural ſtate, and the nerves, in a greater or leſs degree, dark and ſpotted. It is therefore certain, that the venom of the viper is not communicated by the nerve to the parts into which that nerve enters and ramifies; notwithſtanding it is true that the animal dies on which this experiment is made.

Experiments on the Sciatick Nerve, divided in its inferiour Part.

But if the diſeaſe of the venom is not communicated to the parts beneath that in which the nerve has been divided, it may nevertheleſs be communicated to the upper parts, with which the nerve ſtill preſerves its former union and correſpondence entire. The animal continues to be ſenſible to the ſmalleſt violence offered to the nerve, which conſequently never ceaſes to be an organ and inſtrument of ſenſation, and in which that principle, whatever it may be, ſtill exiſts, that produces ſenſation in the machine.

Having laid bare the ſciatick nerve in the uſual way, inſtead of dividing it in the upper part towards the vertebræ, I divided it in the inferiour part towards the feet. The cleared part of the nerve was, as uſual, about an inch and an half in length. I wrapped

wrapped it in linen as in the preceding experiments, and wounded it with the venomous teeth, taking care to cover the whole carefully, to prevent the communication of the venom to the adjacent parts.

The following are the experiments I made in this way:

The ſciatick nerve of a rabbit being laid bare, I cut it in the inferiour part towards the feet, and wrapped it in linen which had been folded ſeven times. I now wounded it repeatedly with the venomous teeth of two vipers, and, in the midſt of this operation, the rabbit exhibited ſtrong marks of pain. It died at the end of twenty hours. The nerve was ſpotted and livid, as were likewiſe the lungs. The blood in the heart was black and grumous; but the muſcles about the nerve ſcarcely ſeemed to have undergone the ſmalleſt change.

This experiment ſeems to confirm us ſtill more in the opinion, that the venom is not communicated to the muſcles adjacent to the nerve, and that there is no local diſeaſe in theſe parts.

I laid bare the ſciatick nerve of another rabbit, cut it in its inferiour part, and wounded it as uſual with the venomous teeth of two vipers. The rabbit cried out and writhed itſelf during the act of wounding the nerve, and died at the end of ſixteen hours. The nerve was livid and inflamed in ſeveral parts. The lungs were covered with large black ſpots. The heart, auricles, and large venous veſſels, contained black grumous blood. All the adipoſe membrane covering the muſcles of the abdomen, was inflamed,

inflamed, and inner part of the ſkin was ſo likewiſe. The ſkin, adipoſe membrane, and muſcles towards the breaſt, were all gangrened. The muſcles adjacent to the nerve were livid for the depth of a line.

This experiment is very different from the preceding one, and may induce one ſtrongly to ſuſpect, that the venom of the viper is alſo venomous to the nerves, and that in theſe caſes, the diſeaſe of the venom is communicated to all the parts of the animal, above that where the nerve has been cut. In ſuch an uncertainty there is no way of coming at the truth, than that of purſuing the experiments. It is almoſt impoſſible not to obtain, in a long continuance of them, ſome agreement and conſtancy in the effects.

I divided one of the ſciatick nerves of a rabbit in the uſual way, and having wrapped it in linen, wounded it with the venomous teeth of two vipers. The rabbit ſhrieked violently at the moment of its being wounded, and died at the end of thirty-ſeven hours. The nerve was full of black and livid ſpots, and the parts adjacent ſomewhat inflamed. The heart was very hard, and very much ſhrivelled. I did not open the rabbit till upwards of an hour after its death. The venæ cavæ, however, ſtill oſcillated with force. Their motion began at the part where they open into the auricle, and they continued to move for upwards of five hours longer, notwithſtanding the cavity of the thorax was expoſed to the open air.

Having

Having divided the ſciatick nerve on one ſide, of another rabbit, and wrapped it carefully in linen, I wounded it in ſeveral places with the venomous teeth of two vipers. The rabbit died at the end of ſixteen hours. The nerve had ſeveral black ſpots on its ſurface, and the adjacent muſcles were livid throughout their whole ſubſtance. The blood in the heart, auricles, and large venous veſſels, was fluid, and a little darker than uſual.

I repeated this experiment with the ſame circumſtances on ſix other rabbits, and the conſequences were perfectly analogous to thoſe I have related above. The rabbits all died ſooner or later, but neither of them in leſs than ſixteen hours, nor after thirty-ſeven. In ſome of them the muſcles circumjacent to the nerve were inflamed and livid throughout their whole ſubſtance; and in others, on the contrary, they were ſimply a little redder than uſual. The blood in the heart, in ſome of the caſes, was fluid, and in the others coagulated. The muſcles, adipoſe membrane, and ſkin, of the breaſt, were inflamed in only one inſtance. The only thing conſtant in theſe experiments was the death of the animal.

We may, in my opinion, deduce in general, from the experiments I have thus far related on the nerves, that the changes obſerved in the muſcles adjacent to the ſciatick nerve, or thoſe in other parts of the animal, are entirely accidental; ſince they ſometimes exiſt, and ſometimes do not.

Experiments

Experiments on the Sciatick Nerve, on which a Ligature was made.

A new ſpecies of experiments remained to be made on the nerves, which might probably decide the queſtion. I reflected that the nerve could only communicate the diſeaſe of the venom to the animal, in conſequence of there being a free communication betwixt the nerve and the animal itſelf, and thought of putting an entire ſtop to this communication, without even cutting the nerve. We know that a thread which makes a ſmall preſſure on a nerve, entirely prevents this communication; that the muſcle no longer obeys the will of the animal; and that the nerve is no longer the inſtrument or organ, either of motion or ſenſation.

In conſequence of this hypotheſis, I laid bare the ſciatick nerve on one ſide, of a rabbit, and tied it ſtrongly in two parts with a thread. There was a portion of nerve betwixt the two ligatures of more than ten lines. I covered it with a piece of linen in ſeveral folds, and wounded it repeatedly with the venomous teeth of two vipers, taking care to cover all the parts about it effectually, to prevent a communication of the venom. The rabbit died at the end of ſixteen hours. The part of the nerve betwixt the ligatures was white; the muſcles adjacent to the nerve were but very little redder than uſual; the heart, auricles, and great venous veſſels,

were

were filled with a fluid blood, ſcarcely darker than it is in its natural ſtate.

I laid bare the ſciatick nerve on one ſide, of another rabbit, and tied it in the way I have juſt deſcribed. I then wounded it betwixt the two ligatures with venomous teeth, and covered it with linen. The rabbit died at the end of eighteen hours. The nerve was in its natural ſtate. The adjacent muſcles were red and livid, for the depth of four lines and more.

Having laid bare one of the ſciatick nerves of another rabbit, I wounded it as above. The rabbit died at the end of ſeventeen hours. The nerve was in its natural ſtate, and the muſcles about it ſcarcely inflamed.

Theſe three experiments ſhow, that the greater or leſs degree of inflammation and lividity in the muſcles adjacent to the ſciatick nerve, is not owing to the venom; and that even the death of the animal may ariſe from ſome other cauſe. It is very certain, that in the caſes in which the nerve is tied, we do not ſee any livid ſpots on this part; and, conſequently, that they are occaſioned by the free communication of the nerve with the animal.

I repeated this experiment, with the ſame circumſtances, on four other rabbits, all of which died in leſs than nineteen hours. The nerve in each of them was white, and in its natural ſtate. The adjacent muſcles in two of them were ſcarcely inflamed; in the other two they were livid for a certain

tain depth. In one of theſe two laſt, a part of the pectoral muſcles was inflamed.

I confeſs, that in combining all theſe experiments together, I find nothing that can give me the ſmalleſt ſuſpicion, of the nerve being a means of communicating the venom of the viper to an animal, and of exciting in it the diſeaſe this venom occaſions. It is true, that there are livid ſpots on the venomed nerve, which are not obſerved when it is tied; but may not theſe be purely mechanical, and the effect of the wounds made by the teeth? And even though they ſhould be occaſioned by the venom itſelf, does it on that account follow, that the venom acts on the nerve, as a venom and not otherwiſe? Is it therefore demonſtrated, that the nerve ought to communicate it to the other parts of the animal?

We are now acquainted with the conſequences that enſue from the application of the venom to the ſciatick nerve; when this nerve is entire; when it is cut, as well above as below; and laſtly, when there are two ligatures made in it. It remains to compare all the effects already known, with thoſe that will be obſerved, in inflicting on the nerve ſimple mechanical wounds. After what we have ſeen, theſe comparative experiments can leave no future doubt.

As the experiments thus far related on the ſciatick nerve, were made in three different ways, ſo I ſhall divide the comparative experiments into three correſponding claſſes.

Expe-

Experiments on the Sciatick Nerve, in which mechanical Wounds were made.

Having laid bare the ſciatick nerve on one ſide, of a rabbit, and wrapped it, as uſual, in linen, to the end that all the circumſtances might agree with thoſe of the preceding experiments, I wounded it in ſeveral places with a viper's tooth, that had been dried for upwards of a month, and had been carefully waſhed in water, to remove all ſuſpicion of its concealing any venom. The rabbit appeared to ſuffer violently when the tooth pierced the nerve. It died at the end of twenty-four hours. The nerve was in ſeveral parts red and livid; the muſcles adjacent to it were inflamed and dark, and theſe appearances extended to the lower part of the leg. The abdominal muſcles and integuments were likewiſe inflamed. The right ventricle contained grumous blood.

I laid bare one of the ſciatick nerves of another rabbit, and having wrapped it in linen as uſual, I pierced it in ſeveral places with the point of a fine needle. The animal ſhrieked terribly, and died at the end of thirty-ſix hours. There were ſeveral dark ſpots in the nerve, and the parts adjacent to it were ſomewhat inflamed. The blood in the heart was black and coagulated.

Having laid bare one of the ſciatick nerves of another rabbit, and wrapped it in linen, I pricked

it ſeveral times with a needle. The rabbit exhibited marks of pain, and died at the end of twenty-ſeven hours. The muſcles about the nerve were ſomewhat livid and inflamed, and the nerve itſelf covered all over with red and black ſpots. The blood in the heart was black and coagulated.

Several important truths are demonſtrated by theſe experiments.

I. That the livid and red ſpots of the nerve are the effect of ſimple mechanical wounds.

II. That the death of the rabbits is owing to the ſimple wound of the nerve, and not to the venom.

III. That the venom of the viper, communicated to the nerves, neither occaſions in any degree the diſeaſe of the venom, nor haſtens the death of the animal.

IV. And laſtly, that the venom of the viper is altogether innocent to the nerves, having no greater action on them than pure water, or the ſimple ſolution of gum arabick in diſtilled water. I have aſſured myſelf by other experiments, that it is not at all offenſive to theſe organs.

The experiments I have juſt related were not yet ſufficient to ſatisfy and convince me perfectly. I knew by experience how eaſy it is to be miſled by facts, when they are but few in number, and was therefore deſirous of repeating the ſame proceſs over again on four other rabbits. The event was perfectly ſimilar to that of the three caſes related above. The rabbits all died; the ſciatick nerve,

in each of them, was more or leſs covered with livid and red ſpots; the adjacent muſcles were, in a greater or leſs degree, inflamed and livid; and the blood in the heart was in general black and coagulated.

Experiments on the Sciatick Nerve.

Having laid bare the ſciatick nerve on one ſide of a rabbit, I tied it in two places with a thread, and pricked it ſeveral times with a needle betwixt the two ligatures. The rabbit died at the end of thirty-three hours. The lungs had ſeveral dark ſpots; the nerve was white, and in its natural ſtate; the blood in the heart was dark, but fluid. The rabbit, when I opened it, was ſtill warm.

Having laid bare the ſciatick nerve on one ſide of a ſecond rabbit, and tied it in two places, I pricked it betwixt the two ligatures with a needle. The rabbit died at the end of eighteen hours. The nerve was white, and in its natural ſtate; the blood in the heart black and coagulated; and the muſcles that ſurrounded the nerve red and livid.

I repeated this experiment on two other rabbits, tying the nerve, and pricking it with a needle, as uſual. Both rabbits died; one at the end of thirty hours, the other at the end of thirty-five. The nerves were in a natural ſtate, but the muſcles were inflamed, and in one of the rabbits, livid for a conſiderable

ſiderable depth. The blood in the heart was black and grumous.

Experiments on the Sciatick Nerve divided above and below.

Having laid bare the ſciatick nerve of a rabbit, I divided it in its inferiour part, and wrapped it in linen, in the way I had done in all the caſes related above. I pricked it ſeveral times with a needle. The rabbit gave repeated ſhrieks, and died at the end of thirty-ſeven hours. The nerve was covered with black and livid ſpots; the adjacent parts were ſomewhat inflamed, and the heart ſhrunk, and very hard. The venæ cavæ continued to move for five hours after I had opened the thorax, their motion beginning at where they ariſe from the auricles.

I divided the ſciatick nerve of another rabbit, and having wrapped it in linen, pricked it ſeveral times with the point of a needle. The rabbit died at the end of fifty-four hours. There were black ſpots in ſeveral parts of the nerve; the muſcles about it were ſcarcely inflamed; the blood in the heart was in a fluid ſtate.

I made the ſame experiment on another rabbit, the ſciatick nerve of which, when divided, I pricked ſeveral times with a needle. The rabbit died at the end of thirty hours. The nerve was in ſeveral places red and livid; the muſcles were livid

and inflamed; and the blood in the heart black and grumous.

I was desirous of repeating the same experiment, with precisely the same circumstances, on four other rabbits, all of which died in less than forty hours, and one of them before the expiration of eighteen. The muscles in all of them were in a greater or less degree inflamed, and the nerve more or less red and livid. The blood in the heart, in some of them only, was black and coagulated.

Seeing that all these experiments correspond in a certain degree, both with each other, and with the relative ones of the venomed nerves, I did not think it necessary to make a great number of trials on the sciatick nerve cut in the upper part. I therefore made only two, and these agreed in their consequences with those of a similar kind in which I employed the venom.

I do not conceive that any doubt can remain after these experiments, as to the entire innocence of the venom of the viper, applied to the sciatick nerve, and as to the impossibility of the bite of this animal producing the disease of the venom, when confined to a nerve alone.

This new truth in animal physicks is of the greatest importance in understanding the nature of the venom of the viper, and its action on the animal body. I must acknowledge, that I had need of all the experiments on the nerves thus far related, and which are in so great a number, and varied so many different ways, to be fully and clearly persuaded of

this circumſtance. Every thing concurred to a belief of the contrary. The rapidity of the diſeaſe, the ſuddenneſs of the death, the momentaneous loſs of ſtrength, the very violent convulſions, the very acute pain, and, in a word, every ſymptom that characterizes the diſeaſes of the nerves, ſeemed to exiſt in the animal when the nerve was bit. It is however certain, that the venom of the viper does not communicate itſelf to the other parts by the medium of the nerves, and that the ſubſtance of them, which cauſes the ſenſation of the animal, and on which life itſelf ſeems to depend, is not changed by the action of this venom. The experiments are direct, they are in a great number, and varied exceedingly; the fact is certain, and the errour was on our ſide, the offspring of prejudice and opinion, and not of nature and experience. On the other hand, we have ſeen that the venom of the viper, introduced into the blood, without touching any veſſel or any ſolid part, kills animals inſtantly, bringing on very acute pains, and very violent convulſions. I have ſeen the relaxed ſphincters give paſſage both to the urine and fœces.

Here an occaſion preſents itſelf of examining the principles and grounds on which this doctrine of theoretical and practical medicine, that aſcribes diſeaſes to the nerves, and ſubmits ſo many motions and functions to a nervous principle, is ſupported. The field is ſo very extenſive, that, although this diſcuſſion might be very uſeful to the practice of medicine, I cannot allow myſelf a moment's ſtay

in it. It will be ſufficient for the preſent to draw this general concluſion—that the uſual ſymptoms of nervous diſeaſes are equivocal and deceptious; that they may exiſt without there being any diſeaſe of the nerves; and that a ſimple change in the blood may be capable of producing all this derangement, and that inſtantly.

Had the celebrated Engliſh phyſician, Mead, known that a ſmall quantity of the venom of the viper introduced into the blood, kills a large and ſtrong animal almoſt inſtantly; and that this venom is entirely innocent to the nerves; he certainly would not have had recourſe to the animal ſpirits, and to the nerves, to explain the action of the venom on animals that have been bit. But he was entirely ignorant of theſe two important truths, which were likewiſe unknown to all the other phyſicians of his time.

Mead employed the ſame principle, that is to ſay, the nerves and animal ſpirits, to explain the nature and effects of the other poiſons. The nerves are affected every where; the animal ſpirits are deranged and in motion every where; and tumult and nervous agitations are obſerved every where. He will have this principle applied, not only to the effects of the venom of the viper and the other poiſons, but likewiſe to ſeveral other very violent diſeaſes, and amongſt others, the plague. This theory is abſolutely falſe as to the venom of the viper, which it ſeemed to favour the moſt. I do not believe it any truer as to ſeveral other poiſons, par-

particularly thoſe of the animal kingdom; and after the experiments I have made, I do not find it demonſtrated as to the plague, and other diſeaſes.

When we examine the reaſons that have determined naturaliſts and phyſicians to recur to the nerves in explaining theſe diſeaſes, (whether natural or the effect of poiſon) we find them to be founded on two principles—the rapidity of the diſeaſe itſelf; and the convulſions and very ſudden proſtration of ſtrength in the animal.

The firſt of theſe two reaſons is of no weight, ſince I have ſhown, that a very ſmall quantity of the venom of the viper introduced into the blood, kills an animal in a few inſtants; and the ſecond is neither evident nor certain, ſince experiment itſelf has demonſtrated to us, that a little of this venom, in the ſame circumſtances, produces very violent convulſions, and cauſes a proſtration of ſtrength in the animal in a few moments, notwithſtanding it only touches the fluid parts of the blood. I do not beſides think it difficult to explain the convulſions, without recurring either to the animal ſpirits, or to the nervous ſyſtem. In the firſt part of this work, I have mentioned the circumſtance of convulſions ariſing ſimply from a want of equilibrium in the parts, occaſioned by the unequal diſtribution of blood in the organs, and by the unequal loſs of irritability in the muſcles. I did not know at that time, either that the nerve was not obnoxious to the venom of the viper, or that this venom was mortal when ſimply introduced into the blood. This ſub-

ject appears to me of ſufficient importance to deſerve a work apart. In that work the other poiſons might be examined, as I have done the venom of the viper; their effects might be analized; and all the circumſtances attending them might be noted. What lights might not ſuch an undertaking throw on animal phyſicks, on the theory of poiſons, and on medicine itſelf! The road is open, and nothing is needed but a patient and induſtrious obſerver. I can venture to promiſe him, in the courſe of a few years, the moſt brilliant, and perhaps too, the moſt uſeful, diſcoveries.

But let us return to our experiments.

Although I had aſſured myſelf of the innocence of the venom of the viper, when applied immediately to the nerves, I had a conſtant ſuſpicion however, that the nerves might at leaſt be neceſſary to the action of this venom on the blood. They might perhaps depoſit ſome unknown principle, ſome ſubtile fluid, in the veſſels of the animal, and this principle, or fluid, might be eſſential to life, and likewiſe to the ſound ſtate of the blood. This was a new mode of conſidering the action of the venom on the living body, differing eſſentially from all the hypotheſes that phyſicians had hitherto invented; and it remained to know whether the venom of the viper would produce a diſeaſe of greater or leſs violence, on the nervous communication betwixt the part bitten and the animal being ſtopped.

Bite of the Viper on Frogs that had been deprived of the Head.

To obtain ſome information on theſe reſearches, I made the following experiments.

I cut off the head of a frog, the leg of which I had bit, twice by a viper. This was followed by no ſymptom of diſeaſe in the part.

I cut off the head of another frog, and, after ſtripping off the ſkin, had one of its legs bit repeatedly by two vipers. The frog has no ſymptom of diſeaſe.

I cut off the head of a third frog, which I had bit repeatedly in the leg, the ſkin ſtill remaining on. There appeared ſome ſlight ſymptoms of diſeaſe in the leg. At the end of two hours I introduced a pin into the ſpinal marrow, and there was ſome ſmall degree of motion in the muſcles.

I repeated this experiment on four other frogs that had been deprived of the head. Three of them had no ſign of diſeaſe; in the fourth there were ſome little indications of it.

Theſe experiments appearing to me neither ſufficiently clear nor uniform, I was deſirous of repeating them on twenty-four other frogs, in which I cut off the head, as uſual. Twelve were bit in the leg repeatedly by ſeveral vipers, and the twelve others had punctures made in the leg, either with fine needles, or with vipers' teeth dried, and free

from

from venom. The conſequences were very vague. Of the twelve that were bit, three only had the diſeaſe of the venom; and of the twelve that were not bit, but pricked with needles, or wounded with venomous teeth, one had ſymptoms of inflammation and lividity in the leg, that might have been confounded with the ſymptoms that accompany the diſeaſe of the venom.

We may, I think, ſay in general terms, that a frog deprived of its head, contracts the diſeaſe of the venom with greater difficulty on that account; and that in theſe circumſtances the part bitten is leſs changed by the venom than otherwiſe. Theſe experiments, however, did not throw a ſufficient light on the queſtions I propoſed to myſelf to clear up, and I therefore determined to fall on a new mode of experimenting.

Experiments on Frogs, in which the Spinal Marrow was divided.

I divided the ſpinal marrow of a frog, at two lines above the part where the nerves that are ſent to the legs and feet, riſe out of the vertebræ. I then had one of the legs of the animal bit repeatedly by two vipers. There was no appearance of a communication of diſeaſe.

I repeated this experiment on four other frogs, and the event was the ſame; I could not obſerve any ſign of diſeaſe in the legs that been bit.

After

After having ſeparated the head from four other frogs, I entirely deſtroyed the ſpinal marrow with a bit of wood. I then had the legs of the animals bit, but could obſerve no ſubſequent ſymptom of diſeaſe.

On repeating the experiment of the divided ſpinal marrow on ſix other frogs, I found in four of them not the ſmalleſt appearance of diſeaſe. There was ſome doubt as to the fifth, but the ſixth appeared to have a real attack of the diſeaſe of the venom.

This laſt caſe made me ſuſpect the experiments I have related above on the nerves of frogs, and I therefore thought it expedient to proceed to ſome experiment that would be more luminous and leſs equivocal. I procured large rabbits for this purpoſe.

Bite of the Viper on the Parts, the Nerves of which had been divided.

I divided the ſciatick and crural nerves of the right leg of a rabbit. I ſtitched up the inciſed ſkin, and had the leg bit by three vipers, each viper biting thrice. The rabbit returned to its food a little time after it had been bit, and recovered. At the end of twenty days I employed it in another experiment. I muſt remark, that a degree of motion ſtill continued in the leg, and that I had

my

my doubts as to the crural nerve having been effectually divided.

I cut the ſciatick and crural nerves of the right leg of another rabbit, aſſuring myſelf that they were completely divided. Having ſtitched up the ſkin, I had the leg bit by three vipers, each of which bit thrice. The rabbit died at the end of eighteen hours; the muſcles of the leg that had been bit were black, livid, ſwelled, and ſphacelated; the abdominal muſcles were inflamed; as was alſo the whole of the internal part of the ſkin.

Theſe two experiments could not have been more contradictory to each other; it is however certain, that in the ſecond caſe there was a true diſeaſe of the venom. The firſt caſe proves nothing more, than that an animal may, on ſome particular occaſion, be bit repeatedly, even by ſeveral vipers, and not have an attack of the diſeaſe. Caſes analogous to this have been related in the preceding parts of this work.

I cut the ſciatick and crural nerves of the leg of another rabbit; the nerves were effectually divided, and the leg had loſt all motion. I then had the part bit by three vipers, repeatedly by each. The rabbit died at the end of ſixteen hours. The muſcles of the leg were livid and gangrened throughout their whole ſubſtance.

I repeated this experiment with the ſame circumſtances on two other rabbits, one of which died at the end of twenty hours, the other at the end of twenty-four. They had both the moſt determined

ed ſymptoms of the diſeaſe of the venom, in the leg that had been bit.

Theſe experiments are poſitive and uniform, and prove indubitably, that it is of no conſequence as to the diſeaſe of the venom, whether the nerves of the parts that are bit, are, or are not, cut; or whether they communicate, or have no longer any communication, with the animal.

But in theſe experiments, there ſtill ſubſiſts a nervous communication betwixt the part bitten and the animal. This communication is formed by the ſkin of the animal that covers the part where the venom is received. We muſt therefore put a ſtop likewiſe to this communication, by removing the ſkin.

After having divided the ſciatick and crural nerves on one ſide, of a rabbit, and ſtitched the inciſed ſkin, I made a circular inciſion in the leg, at four fingers diſtance above the part where I had propoſed to have it bit by vipers. As ſoon as the inciſion was made, I ſtitched it all round. I then had the leg bit repeatedly by three vipers, the teeth penetrating into the ſkin. At the end of two hours there was no ſymptom of diſeaſe; at the end of ſix the part that had been bit was viſibly inflamed; at the end of ten, there was an oozing of blood from the part of the ſkin that had been punctured by the teeth of the viper; at the end of twenty-two, the blood flowed in ſtill greater abundance; at the end of twenty-four, the part was very much ſwelled, but no lividneſs was to be diſcerned; at the end of

thirty

thirty the ſkin opened, and a wound formed. The rabbit after living eight days, was employed on another occaſion.

In this laſt experiment, there can be no doubt but that the diſeaſe of the venom was communicated to the part bitten, although it was not very violent.

I now thought of making a comparative experiment.

For this purpoſe I had a rabbit bit repeatedly in the leg by three vipers, without cutting either the nerves or the ſkin. At the end of eight hours the leg was ſwelled, but not livid. At the end of twenty-two, a cyſt or bladder, filled with a dark-coloured humour, formed betwixt the legs, near the part that had been bit. The rabbit died at the end of forty hours. The ſkin was broken, and corroded at the part bitten; the muſcles of the leg were livid and gangrened; the heart, auricles, and large veſſels, were filled with black and grumous blood; and there were concretions of blood even in the aorta, which is uſually found empty.

I repeated the preceding experiment on three other rabbits, and, after dividing the nerves, making the circular inciſion in the ſkin, and ſtitching it, had each of them bit repeatedly by three vipers. They all had ſymptoms of the diſeaſe of the venom in the part bitten.

I had another ſuſpicion, that there might ſtill ſubſiſt, after all, ſome nervous communication betwixt the animal and the leg, after the ſciatick nerve was divided. I conjectured that the viper

might have ſtruck its teeth on ſome fibre of the great glutous muſcle, which deſcends very low in the leg. This was a ſufficient inducement to me to make the following experiments.

I divided the ſciatick and crural nerves on one ſide, of a rabbit, and made a circular inciſion in the ſkin, which I afterwards ſtitched. I had the leg bit by three vipers, by each repeatedly, but in ſo low a part as to avoid the above-mentioned muſcle. At the end of two hours the part that had been bit began to inflame; at the end of twenty-two the ſkin was broken, but not puffed up; at the end of forty-two the animal was apparently recovered; and at the end of eight days, was employed in other purpoſes.

I divided the ſciatick and crural nerves on one ſide, of another rabbit, and made an inciſion, which I afterwards ſtitched, for the whole circumference of the leg. I had the leg bit in its inferiour part by three vipers, each viper biting thrice. At the end of eight hours, the ſkin burſt, and diſcharged a humour; at the end of twenty-two this broken ſkin was ſwelled and livid; and at the end of ſixty, the rabbit was in a dying ſtate. I opened it, and found all the muſcles of the leg gangrened, and almoſt the whole of the adipoſe membrane that covers the abdominal muſcles full of extravaſated blood. The blood in the heart was in a diſſolved ſtate.

I divided the ſciatick and crural nerves of another rabbit, and made an inciſion in the ſkin, which I ſtitched, all round the leg. I had the inferiour part

of the leg bit repeatedly by three vipers. At the end of two hours the part that had been bit was become ſwelled; at the end of eight it was conſiderably ſo; at the end of twenty-two, the ſkin was broken, but without ſwelling; at the end of forty-two hours, there was ſimply a wound at the part bitten. The rabbit lived ten days, and was employed in other experiments.

Theſe experiments ſhow, that the venom of the viper produces its uſual effects on the parts bitten, notwithſtanding all nervous communication has been ſtopped, betwixt theſe and the other parts of the animal. But it is not yet decided, whether, ſuppoſing ſome active principle which mixes with the blood to be ſeparated from the nerve, this principle does, or does not, ceaſe to exiſt the inſtant the nerve is divided; particularly as the nerves ſtill remain in the part bitten, although they are no longer the inſtruments of ſenſation, and voluntary motion. This reflection made me fall upon the following experiments.

I divided the ſciatick and crural nerves on one ſide, of a rabbit, and likewiſe made the circular inciſion in the ſkin, which I ſtitched. I left the part in this ſtate during ſixteen hours, and then had it bit below the inciſion by three vipers, each of which bit repeatedly. The rabbit died at the end of twenty-two hours. All the muſcles of the leg were livid, gangrened, and putrid; the pericardium was filled with a tranſparent fluid; the right ventricle and right auricle were filled with black grumous blood;

blood; and the blood in the large veſſels was in the ſame ſtate.

Effects of the Bite of the Viper on Rabbits, the ſpinal Marrow of which was divided.

I ſhall conclude my experiments on the parts deprived of their nerves, and bit by the viper, by relating three experiments made on rabbits, the ſpinal marrow of which had been entirely cut through. I divided it beneath the loins, and in ſo effectual a way on all ſides, that there could be no ſuſpicion of any communication of nerves, betwixt the legs and the other parts of the animal.

The ſpinal marrow of a rabbit having been divided in the way I have juſt mentioned, and a circular inciſion round the leg made in the ſkin, which I had ſtitched, I had the inferiour part of the leg bit by three vipers, by each repeatedly. At the end of an hour, a ſmall tumour formed in the part bitten; at the end of two this part was very much ſwelled, and very livid; and the rabbit died at the end of ſeven. The part that had been bit was gangrened all over, and the gangrene penetrated into the whole ſubſtance of the muſcles that had been wounded by the teeth. The blood in the heart was black and grumous.

I divided the ſpinal marrow of another rabbit, and by the help of ſciſſars detached a great portion

of the ſkin that covers the crural muſcles. When the latter were in this manner laid bare, I had them bit repeatedly by three vipers. A few minutes after, there were ſymptoms of the diſeaſe of the venom in the part, and the rabbit died at the end of ſeven hours. The muſcles that had been bit were livid and inflamed; the blood all about them was extravaſated into the cellular membrane; there were livid ſpots in the lungs; and the heart was filled with blood, which was almoſt in an entire ſtate of diſſolution.

I repeated this experiment on another rabbit, with the ſame circumſtances, and the reſult was likewiſe the ſame. The rabbit died at the end of ſix hours, with the muſcles of its leg affected by the diſeaſe of the venom.

Thus are we aſſured, that the nerves which are ſent to the parts bitten, in no way contribute to the diſeaſe of the venom of the viper, and that this venom is altogether innocent to the nerves; important truths we were before ignorant of. What is ſtill hidden from our ſight, is the occaſion of the blood, united with the venom, coagulating in an inſtant when it is incloſed in the veſſels of the animal, and not coagulating in the open air.

Effects of the Venom on the Parts of an Animal, the Circulation of which was interrupted.

I hoped to draw ſome information from the experiments which follow. They conſiſted in examining the effects of the bite of the viper, on the parts of animals in which the arteries and veins had been previouſly tied by ligature. This was altogether a novel enquiry, and it was not at all amiſs to know the effects that would be produced in ſimilar caſes.

I made a ligature in the belly, on the aorta deſcendens and vena cava of a rabbit. Having ſtitched the ſkin, I had the leg of the rabbit bit by three vipers, by each repeatedly. The animal died at the end of nine hours. The leg was gangrened in the part where the bites had been received, but in no other.

I divided and removed in the belly of a rabbit, the arteries and veins that go to the right leg, and likewiſe removed a large portion of the ſkin of the leg, which I had bit at the part where the muſcles were bare, by three vipers, each of which bit thrice. At the end of an hour there were certain ſymptoms of local diſeaſe. At the end of two hours the leg was livid at the part bitten, but not elſewhere. The heart was filled with black and grumous blood.

I made a ligature, as in the firſt experiment, on the arteries and veins within the belly of two rabbits. Each of them was bit repeatedly by three vi-

pers. In one of them the ſkin was in an entire ſtate; in the other it was cut circularly, as in an amputation, and ſtitched. They were both dead at the expiration of twenty hours. There were ſymptoms of diſeaſe in the parts bitten. However, the diſeaſe was ſlight, and neither deep nor extenſive. The blood in the heart was black and coagulated.

I divided in the belly, the arteries and veins of another rabbit, but neglected to have it bit by vipers. It died at the end of ſixteen hours. The lungs were livid; and the heart, auricles, and great veſſels, filled with black and grumous blood. This experiment is a ſtill further demonſtration to us, that the grumous blood in the heart and neighbouring veſſels is an equivocal ſign, when it is taken alone, and without being accompanied by others.

I repeated this experiment of making a ligature in the belly, on the veins and arteries, on three other rabbits, having each of them bit in the leg by three vipers. They all three died in leſs than ſeventeen hours. The muſcles that had been bit were attacked by the diſeaſe of the venom, but not the adjacent ones. This local diſeaſe was but of little conſequence.

We may deduce with certainty from theſe experiments, that the venom of the viper produces its uſual effects, even when the parts bitten no longer participate of the circulation of the blood in the animal machine. In theſe caſes we ſee in general, that the diſeaſe is leſs extenſive and leſs violent, than when the blood circulates in the part; and this

particular agrees very well with the experiments in which the venom was injected into the jugular vein.

Effects of the Venom on Parts, the Veſſels of which were cut.

I was deſirous of ſeeing what would happen to a rabbit, the crural arteries and veins of which were tied, and cut beneath the ligature, ſeveral hours prior to the leg being bit. In theſe caſes the blood has not only ceaſed to circulate in the leg, but has been a long time ſtagnant: it may already be changed in a great meaſure, may have ſuſtained a conſiderable loſs in its quantity, and may be deprived of a ſubtile principle of ſome kind or other. The rabbit that I got ready in this way, remained in this ſtate during upwards of eight hours. At the end of that time, I had it bit in the leg by three vipers, each of which bit repeatedly, the ſkin having been previouſly removed from the part. The rabbit died three hours after. The muſcle at the part where the vipers had bit, was ſomewhat darker than in the adjacent parts; but this was ſcarcely ſenſible.

I cut the crural artery and vein of a rabbit, in the ſame way beneath the ligature, and waited ten hours before I had it bit. At the end of twenty hours it was very lively, and I had it again bit repeatedly by three vipers, the leg being previouſly bared of the ſkin, in the part bitten. It died ſix hours after. The muſcles that had been bit were livid throughout

their whole ſubſtance, but the diſeaſe was confined to the part on which I operated.

I repeated this experiment on two other rabbits, having them bit each in a leg, without removing the ſkin, eight hours after the ligature had been made, and the crural artery and vein cut. I took the precaution to make repeated compreſſions on the leg, that the arterial and venous blood might flow out at the opening in the veſſels. Both rabbits died in leſs than eleven hours. The fleſh where the teeth had entered appeared of a deeper colour than uſual, and this diſcoloration penetrated to the depth the teeth had extended to. The other parts were in a natural ſtate.

I got ready two other rabbits in the ſame way, as a comparative experiment; they were therefore not bit by vipers. They were both dead at the end of ſeventy-two hours.

It now remained to examine the effects of the venom of the viper, after having tied the arterial and venous veſſels ſeparately.

For this purpoſe I made a ligature on the vena cava in the belly of a rabbit, and afterwards made the circular inciſion in the ſkin of the leg, and ſtitched it. I then had the leg bit repeatedly by three vipers. At the end of twenty-four hours, there were ſymptoms of the diſeaſe of the venom in the part bitten. I killed the rabbit in this ſtate, and found that the diſeaſe was circumſcribed to the inciſion made in the ſkin. The muſcles were livid, and the

the adipoſe membrane filled with dark extravaſated blood.

I tied in the belly, the vena cava of another rabbit, and had its leg bit repeatedly by three vipers. At the end of two hours there was an extenſion of the ſkin at the part that had been bit, but ſcarcely any ſenſible ſwelling; at the end of four hours, a moiſture exuded from it; at the end of ten hours the ſwelling had encreaſed; and the rabbit died at the end of fifteen. The part bitten was livid and gangrened throughout its whole ſubſtance; the diſeaſe was, however, entirely confined to the leg.

The conſequences of experiments on two other rabbits treated in the ſame way, were pretty ſimilar.

I tied the aorta in the belly of a rabbit, and had its leg, covered by the ſkin, bit repeatedly by three vipers. At the end of ſix hours, the ſymptoms of diſeaſe were perceptible, and the rabbit died at the end of fifteen. The leg was ſwelled and livid, and the diſcoloration penetrated ſome depth into the muſcles. The blood was black in the part that had been bit, and was coagulated in the large veſſels.

This experiment repeated on two other rabbits was attended with pretty much the ſame ſucceſs.

I ſhall conclude this chapter, by relating in a few words, two experiments made on rabbits, in the belly of which I had divided all the lymphatick veſſels I could find, as far as the ductus thoracicus. An hour after this operation, I had both rabbits bit in the legs, covered by the ſkin, repeatedly by three

vipers. At the end of ſix hours the leg in each, exhibited the moſt certain marks of the diſeaſe of the venom. It was livid and ſwelled, and a good deal of humour oozed from it. Both rabbits died at the end of eighteen hours, and the muſcles of the leg in each, were livid throughout their whole ſubſtance.

Expecting nothing from the continuation of theſe experiments, and ſeeing that the ſtoppage of the circulation of the lymph and chyle has no influence on the uſual effects of the venom of the viper, I did not think it neceſſary to make any further progreſs in this ſubject.

CHAPTER V.

Effects of the Venom of the Viper on Blood expoſed to the open Air.

ALTHOUGH the experiments hitherto related afford us very important information, we are ſtill in the dark as to the circumſtance of the blood coagulating when united in the veſſels with the venom of the viper, and not coagulating when blended with the latter in the open air. I have at leaſt conceived that I conſtantly diſtinguiſhed a very ſenſible difference in this fluid, when I had the leg of an

animal

animal bit after its being ſeparated from the animal itſelf, and when I had it bit, either ſtill adhering in an entire ſtate to the animal, or faſtened to it mechanically.

In this uncertainty, I judged it expedient to make an analyſis, followed by the experiment of Mead, that relates to the effects of the venom of the viper on the blood drawn from an animal; and as Mead made his experiment on a ſmall quantity of venom and a large quantity of blood, I determined that my trials ſhould be on much ſmaller quantities of the latter, that the effects might be more ſenſible.

I let fall into a ſmall conical glaſs, three drops of the venom of the viper, and twenty drops of the blood that flowed from the neck of a fowl, into which I had made an inciſion. I reclined the glaſs, and ſhook it circularly for ten ſeconds, that the venom and blood might mix well together.

At the ſame time I let fall into a ſimilar glaſs, twenty drops of the blood of the ſame fowl, in the ſame ſtate as the laſt. I ſhook the glaſs as I had done on the preceding occaſion, that, the venom excepted, all circumſtances might be alike. At the end of two minutes, the blood unmixed with the venom coagulated, and was of a fine vermillion colour. On the contrary, the blood united to the venom was black and fluid, notwithſtanding it was a little viſcous and compact.

I repeated this experiment, and the event was the ſame. The venomed blood did not coagulate, and, as before, was of a dark colour; whilſt, on the contrary,

trary, the blood that was not venomed coagulated very ſoon, and ſtill preſerved its bright red colour.

I repeated this experiment on the blood of a guineapig, one of the legs of which I had cut off. The venomed blood at the end of twenty-four hours was ſtill black and diſſolved, whilſt the other coagulated in leſs than two minutes, and continued to preſerve its red colour. The venomed blood did not harden till it dried by degrees and ſplit into ſcales, and preſerved its black colour till the laſt; inſtead of which the blood in a pure ſtate, ſtill continued red, even after it had dried and ſplit.

The black colour of the blood that was mixed with the venom, agrees very well with the moſt uſual effects of the bite of the viper on animals, and with the effects of the venom introduced into the jugular vein of rabbits. But the other circumſtance in theſe latter experiments is altogether ſingular and unexpected. Inſtead of the venom coagulating the blood, as one would ſuppoſe it ought to do, it even keeps it from that coagulation which is natural to it in the open air, and preſerves it in a conſtant ſtate of diſſolution. Here then the venom not only does not produce its ordinary effect on the blood, namely, that of coagulating it, but produces an altogether contrary effect, namely, that of keeping it in a diſſolved ſtate, and ſo preventing that coagulation which would otherwiſe take place.

This ſingular effect of the venom on blood expoſed to the air, ſeemed to promiſe ſome important diſcovery, relative to the action of it that ſucceeds

the

the bite of the viper in animals. I reflected, that this bite is entirely innocent to the viper itſelf, and likewiſe to many other animals with cold blood, and that it is not mortal to certain animals, ſuch as frogs, in which it does not produce the diſeaſe of the venom till very late, and with difficulty. In conſequence of this, I perſuaded myſelf that the effects of the venom on the blood of vipers and frogs muſt be very different from thoſe it produces on the blood of warm animals, and that from this difference, that of the diſeaſe and death of theſe animals muſt preciſely depend. Theſe were my reaſonings, and I drew no ſmall expectations from them.

I accordingly put into a ſmall conical glaſs, three drops of venom, and thirty drops of the blood that flowed from the neck of a viper, after I had cut off its head. I ſhook the glaſs, as uſual. The blood, which became ſomewhat darker, did not coagulate. At the end of two hours there was a ſeparation of ſerum, which floated on the top, the red part of the blood being beneath. This was dark, and viſcid like glue, but was not coagulated.

I had at the ſame time got ready a comparative experiment. I had put into a glaſs of the ſame kind, thirty drops of the blood of the ſame viper, but without having introduced the venom. I ſhook the glaſs, as uſual. The blood did not coagulate, and became covered with a great deal of ſerum, through which the ſanguineous fibres, which were very red, were diſtinguiſhed. At the end of two hours

hours the ſerum was in a much larger quantity than in the preceding experiment; at the end of twenty-four the red fibres were in their uſual ſtate; but notwithſtanding this, the blood was thinner than that which had been venomed. At the end of thirty-five it was ſtill fluid, with a great deal of ſerum ſwimming on the ſurface; at the end of fifty it was become thicker and more tenacious; and at the end of ſixty was dry and red.

I mixed in a glaſs, three drops of venom with fifty of the freſh blood of a viper, and received in another glaſs fifty drops of the ſame blood, without making any addition to it. I ſhook both glaſſes a little, and equally. The blood unmixed with venom, remained till the laſt redder than the other, and with a greater proportion of ſerum. At the end of thirty hours the venomed blood coagulated, but the other did not.

We ſee by theſe experiments, that the colour of the blood of vipers which has been united with the venom of the animal, agrees very well with that of the blood of warm animals blended with the venom in the ſame way, notwithſtanding there is a great difference in all the other circumſtances. Theſe experiments, however, are as yet too little varied to admit any certain concluſions to be drawn from them.

I put three drops of venom into a glaſs, and added thirty drops of the blood of a frog, the head of which I had juſt cut off. I likewiſe put thirty drops of this blood into a glaſs, without adding any venom.

venom. I ſhook the two glaſſes, as uſual, and at the end of thirty minutes examined the blood in each. I found the venomed blood black, but not coagulated. The blood unmixed with venom had leſs ſerum than the other, was redder and more fibrous, and was likewiſe in a fluid ſtate. At the end of three hours the blood was black and diſſolved, but viſcous, and without any apparent ſerum. The other blood had a great deal of ſerum on its ſurface. At the bottom it was red and coagulated, but the coagulum was moveable, fibrous, and viſcous.

Not ſatisfied with this laſt experiment, which I repeated twice with pretty much the ſame ſucceſs, I determined to make at the ſame time, experiments on the blood of vipers, on that of frogs, and on that of guineapigs, and to follow minutely all the changes that I ſhould obſerve.

I took ſix conical glaſſes, ſimilar to thoſe I had employed before, and put into each of the three firſt, four drops of venom, and thirty drops of blood. In the firſt glaſs was the blood of a viper, in the ſecond that of a frog, and in the third that of a guineapig. In each of the other three glaſſes I ſimply put fifty drops of blood, taken from one of the ſame animals, and in the ſame order. I ſhook the ſix glaſſes a little, and equally, and then left them in an undiſturbed ſtate for ſome time. At the end of a few minutes the blood in the three venomed glaſſes was black, and much more diſcoloured than that in the three others, and was already

ready coagulated; the blood of the viper unmixed with venom, was however much leſs ſo than the others, and was perhaps rather viſcous than effectually coagulated. The blood of the viper is beſides of a darker red than either that of the frog or guineapig. At the end of ſome time, I obſerved that the venomed blood of the viper, and that of the frog, had a good deal of ſerum on their ſurface, but that there was none on that of the venomed blood of the guineapig. There was likewiſe no appearance of ſerum on the ſurface of the blood in the three glaſſes, in which it was unmixed with venom. At the end of eight hours the blood of the frog contained as much ſerum as the venomed blood, and was in the ſame diſſolved ſtate as the laſt, but redder. The pure blood of the viper never had any ſerum on the top, and continued coagulated as uſual; but the venomed blood of the viper was darker than the other, diſſolved, and extremely viſcous. At the end of three days the venomed blood of the viper ſtill preſerved the large quantity of ſerum it had at the beginning; it was, however, black and viſcous. The blood of the viper, unmixed with venom, had a little ſerum, was red, fibrous, and almoſt wholly coagulated. The venomed blood of the frog was entirely diſſolved, of a greeniſh hue, and contained a little ſerum; but that which had not been venomed, had a great deal of ſerum on its ſurface, was coagulated, and redder than the other. The venomed

blood

blood of the guineapig was black, viſcous, and without ſerum.

At the end of eight hours I examined the red globules of the blood in the three venomed glaſſes, and found them but very little changed in ſhape, and ſcarcely different from the globules of the pure blood in the three other glaſſes. At the end of eight days, however, I found that the globules of the venomed blood of the viper were conſiderably altered in ſhape; ſeveral of them were broken, and they were in general much more changed than thoſe of the viper's blood unmixed with venom. The globules of the venomed blood of the frog were almoſt all diſſolved; thoſe that were not were diſfigured, and very much broken. The venomed blood of the guineapig had, on the contrary, its globules enlarged; they had in ſome meaſure changed their ſhape, and were in a greater or leſs degree diſſolved; they did not differ a great deal from thoſe of the pure blood of the ſame animal.

Theſe laſt obſervations on the globules of blood can be of no uſe in explaining the immediate effects of the venom of the viper on its being introduced into the veins; and the changes we have ſeen in theſe globules are not obſerved till a long time after the action of the venom on the animal. If the animal is ſmall, it is dead a long time before any of theſe changes are obſerved.

I repeated the experiment on the blood of the viper, frog, and guineapig, twice, and the conſequences, although they were not in every reſpect

perfectly ſimilar, were very uniform. I have therefore deemed it unneceſſary to enter into a detail of them.

We ſee in a general way, that the venom of the viper changes the blood both of warm and cold animals black; that of the animals on which it acts as a venom, and that of thoſe on which it has no ſuch action. This very uniformity of change of colour ſhows, however, that the venom of the viper does not kill animals, in conſequence of the principle that changes the blood with which it is united, black. It would otherwiſe be a venom to the viper itſelf, which it is not.

But it is not the ſame as to the coagulation of the blood. The venom acts little, or not at all, on the blood of the viper, and the trifling variations we have obſerved in this reſpect are not at all to be attended to. It is otherwiſe as to the blood of the frog, and ſtill more ſo as to that of the guineapig. Scarcely is the latter in the glaſs, when it coagulates; inſtead of which if it is blended with a few drops of venom, it coagulates no more, and remains black, viſcous, and without ſerum. This effect of the venom is the more ſingular, as one would ſuppoſe that it ought to produce a very contrary one. Does the venom, when united with the blood, becauſe it is a venom, or from ſome other principle, deprive it of the power of coagulating?

It has been ſeen that the venom of the viper produces a ſenſible change on the blood drawn from the veſſels of an animal. In theſe caſes the blood

blood becomes black, and remains fluid, inſtead of coagulating, as it conſtantly does, when it is not mixed with the venom. On the contrary, this venom, when it is introduced into the circulation of an animal, ſuddenly impedes it, by cauſing a coagulation of the blood. The effects of the venom on the blood of animals are certain, but we do not on that account know, on what they depend, nor by what mechaniſm all theſe changes are wrought. Does the venom of the viper act on the blood ſimply as a venom? that is to ſay, by the very principle that renders it mortal?—We have ſeen that this venom is a true gummy ſubſtance, and that it has all the properties which characterize gums. We have likewiſe ſeen, that gums are entirely innocent to animals; and I have obſerved, that when they are injected in a very ſmall quantity into the blood, the death of the animal is by no means ſubſequent. But why may not the black colour of the venomed blood, and the fluidity it preſerves out of the veſſels, depend on the gummy principle of the venom? We know that gums abound with phlogiſton, and that phlogiſton gives a black tinge to the blood. It is true, that as a gummy ſubſtance, it ſeems that it ought rather to coagulate the blood, than to keep it in a ſtate of diſſolution; but experiment alone can reply to all theſe doubts.

Comparative Experiments made with Gum Arabick.

I diſſolved a few grains of gum arabick in a ſmall quantity of warm, diſtilled, water. The mucilage was tranſparent, and almoſt fluid. I put three drops of this mucilage into a glaſs, and added ſixty drops of the blood of a pigeon, at the juncture of its flowing from the divided veſſels.

At the ſame time I put three drops of the venom of the viper into another glaſs, and added ſixty drops of the ſame blood, perfectly in the ſame ſtate.

I ſhook both glaſſes for a minute, that their contents might be well blended together. At the end of two minutes the blood united with the gum and coagulated, its colour continuing red as in its natural ſtate, and no ſeparation of ſerum followed, although I kept it two days in the glaſs. The blood in the other glaſs ſuddenly became black, and continued fluid as uſual.

We ſee by this experiment, that gummy ſubſtances do not give a dark tinge to the blood, and that they have not the property of keeping it in a diſſolved ſtate, and preventing its natural coagulation. Thus, then, the changes that are wrought in the blood by the venom of the viper, are not the effect of a gummy principle, but of ſome other principle yet unknown to us, probably the very one that conſtitutes it a venom; and indeed we have hitherto

hitherto been able to diſtinguiſh nothing more in this humour, than a gummy principle, and a venomous principle deſtructive of animal life.

I was afterwards deſirous of trying whether the venom of the viper would be no longer a venom when mixed with the blood. For this purpoſe, I put into a glaſs thirty drops of the blood of a pigeon warm from the veſſels, and three drops o venom. I blended the liquors well together, and after twenty-four hours were elapſed, applied ſeveral drops of the mixture to the muſcles of a pigeon. The pigeon ſurvived, and at the end of thirty hours ſeemed ſcarcely to have any ſymptom of diſeaſe.

I prepared a mixture of venom and blood in the ſame way, in another glaſs, but made it with an equal quantity of each fluid. Two minutes after, I covered with it the wounded muſcles of a pigeon, which, although it ſurvived, had certain ſymptoms of the diſeaſe of the venom.

I repeated this experiment on four other pigeons. Three of them died in leſs than eighteen minutes; and the fourth had ſo violent an attack of the diſeaſe, as not to recover till the end of the ſixth day. I got ready two other pigeons in the ſame way, and did not employ the venom till half an hour after its union with the blood in the glaſs. Both pigeons died.

It appears from all theſe experiments, that the venom does not loſe its deadly qualities in conſequence of being united with the blood.

We have ſeen that the venom of the viper is a true gum, and that it has all the eſſential properties of a gum. Why may not this venom prevent the coagulation of the blood in warm animals, and likewiſe in ſeveral of the cold ones, as a ſimple gum, and not as a venom? And why likewiſe may not the blood of the viper be different from that of other animals, ſince we ſee that the venom is innocent to the viper, and not ſo to the others?

In this caſe again it belonged to experiment to decide.

As it did not appear to me that the experiments I have hitherto related were ſufficient to explain this difficult phenomenon, of blood which coagulates in the encloſed veſſels of an animal, and which does not coagulate in glaſſes in the open air, I thought it neceſſary to examine more attentively than before, the effects of the venom on the legs of animals cut off, and bit by the viper; and likewiſe on thoſe on which a ligature had been made previouſly to their being bit. I was apprehenſive that I might have made ſome miſtake in my prior experiments, and that ſome neceſſary attention might have eſcaped me. It was natural to conceive, that after all I had ſeen in my laſt experiments, I was better prepared for nice obſervance than before.

With this preſumption, I made the following experiments.

I had a pigeon bit in the leg repeatedly by a viper, and a few ſeconds after cut off the limb. There was a degree of lividity at the preciſe ſpot

wher

where the teeth had penetrated; but this was ſcarcely perceptible.

On repeating this experiment with the ſame circumſtances, the conſequences were perfectly ſimilar.

I had the leg of another pigeon bit by a viper, a moment after I had cut it off. There was no ſymptom of diſeaſe, nor any livid appearance.

I wounded the leg of a pigeon with a venomous tooth, and cut it off immediately after. There was ſome appearance of grumous blood in the muſcle the tooth had pierced.

I wounded the leg of a pigeon with a viper's tooth that had been dry a long time, and at the ſame time wounded the pigeon's other leg with a venomous tooth. The wounds made with the venomous tooth became livid, and this lividity penetrated into the whole ſubſtance of the muſcle. There was no appearance of any change in the part of the other leg that had been pierced by the dried tooth.

I wounded the leg of a pigeon with venomous teeth, and cut it off immediately after. There was a dark ſpot, which however I could ſcarcely diſtinguiſh, at the part the teeth had penetrated.

I forced the venomous tooth into the leg of a pigeon, and cut it off immediately after. There was no ſymptom of the diſeaſe of the venom.

I cut off the leg of another pigeon, and wounded it immediately after with a venomous tooth. There was a ſlight appearance of dark extravaſated blood.

I again forced a venomous tooth into the leg of a pigeon, and cut it off immediately after. This was not ſucceeded by any ſymptom of the diſeaſe of the venom.

I pricked the leg of a pigeon repeatedly with the point of a needle, and cut it off immediately after. I obſerved dark and extravaſated blood at the part that had been pricked.

Although the greater part of theſe experiments ſhow, that the venom of the viper has no action on the parts of animals that have been cut off, there are ſome of them, however, in which we find a ſlight appearance of dark and extravaſated blood.

The experiment made with the needle rendered what I was deſirous of deducing from the others ſtill more uncertain. One would conceive that every time the large veſſels are ruptured, and there is a ſenſible hemorrhage from them, the ſpots and dark colour may appear, without the intervention of venom.

It is in general true, that there exiſts a notable difference betwixt the effects of the venom of the viper introduced into a leg that has been cut off, and the effects of the ſame venom communicated to a leg that continues to make a part of the animal. This difference may occur, either becauſe the quantity of blood in the amputated leg is leſſened, becauſe the blood receives ſomething from the air, or becauſe it, on the contrary, loſes ſomething when in contact with the air. To diſcover which

of

of theſe hypotheſes is the moſt probable one, I made the following experiments.

Experiments on the Venom of the Viper on Limbs that were ſheltered from the Air.

I placed a pigeon in water in ſuch a way, that I could cut off one of its legs without there being any communication betwixt the divided part and the air. A moment before I cut it off I had wounded the leg with a venomous tooth. At the end of four minutes I drew it out of the water. At the part where the tooth had pierced the muſcle, there was a ſmall livid ſpot, on which I immediately made an inciſion, and found it to have penetrated juſt as far as the tooth and venom had reached.

I repeated this experiment twice, and the conſequence was the ſame. The livid ſpot had extended in the ſame way into the ſubſtance of the muſcle.

The blood of the leg amputated in the water, flows from the veſſels in the ſame way as if the part had been cut off in the open air. The ſymptoms, therefore, of the diſeaſe of the venom in the leg ſtill adhering to the animal, and the abſence of theſe ſymptoms when it is detached, do not depend on the different quantity of blood that is found in the two different ſtates of the leg.

This experiment ſeems likewiſe to determine that the blood meets with no eſſential loſs when it

is expoſed to the air, ſince it does not ſeem probable that the water which ſuffers the blood to flow from the leg, prevents this ſuppoſed principle from eſcaping with it.

It remains probable, then, that the contact of the air cauſes ſuch a change in the blood of the leg, and that the air itſelf unites in ſuch a way with it, as to produce the diverſity of effects we have obſerved, although it is certain that we cannot explain what this change conſiſts in, and how, in theſe caſes, the air unites itſelf with the blood.

New Experiments on Parts that were cut, after the Circulation had been interrupted in them by a Ligature.

I had now an important experiment to make, to determine the effects of the venom of the viper on parts of animals tied, and afterwards cut off.

I had the leg of a pigeon bit by a viper at the ſame inſtant that I had it tied and cut off. The whole operation was made in three ſeconds, and required the aſſiſtance of three perſons. The amputation was made over the ligature, which was ſo very tight as to prevent the ſmalleſt hemorrhage. The leg when cut off had the moſt decided ſymptoms of the diſeaſe of the venom. There were livid ſpots on it, the veſſels were black and ſwelled, and the blood black, and partly condenſed. Having opened the muſcles, I found that the livid colour

extended

extended for the whole depth of thoſe that were wounded.

I immediately made a ſecond experiment, ſimilar in every reſpect to the foregoing, except that the leg was not bit by a viper. It had not the ſmalleſt ſymptom of diſeaſe.

I had the leg of another pigeon bit once by a viper, and, after four ſeconds were elapſed, made the ligature and the amputation at one and the ſame inſtant. In leſs than a minute the ſymptoms of the diſeaſe were apparent, the whole ſubſtance of the wounded muſcles being in a livid ſtate.

I tied and cut off the leg of a pigeon, and immediately after had it bit once by a viper. The ſymptoms of the diſeaſe of the venom in the part were very violent, and the muſcles were livid throughout their whole ſubſtance.

I tied and cut off the leg of another pigeon, and had it afterwards bit by a viper. The whole ſubſtance of the muſcles was livid.

Theſe experiments ſeemed to me ſufficiently uniform to diſpenſe with my multiplying them any further. They demonſtrate, that the venom of the viper acts as a venom on theſe parts, although they are detached from the animals, provided there is no hemorrhage from them.

We likewiſe ſee, that it is not neceſſary that the uſual circulation of the blood and other humours ſtill continue in the part, ſince I have ſince obſerved that the venom acts on the legs that have been

been tied, even when they have been bit a pretty conſiderable time after the ligature has been made.

Experiments on Animals with warm Blood, deprived of the Head.

The experiments made on the frogs that had been deprived of the head, in which it appeared to me that the diſeaſe of the venom was communicated with difficulty, gave me the idea of ſeeing whether it would be the ſame with warm animals. Theſe experiments have ſome relation to thoſe that were made on the legs of animals cut off and afterwards bit, and only differ from them in this particular—that here the greater part of the body continues in conjunction with the leg, notwithſtanding the blood flows in a great quantity from the neck, where the inciſion has been made.

I divided the trachea arteria of a fowl, and having fixed in it the nozzle of a ſmall pair of bellows, inſtantly cut off the head. I now began to work the bellows, and at the ſame time had the leg of the fowl bit repeatedly by two vipers. The fowl lived for upwards of fifteen minutes. There were deep livid ſpots in the leg, at the part the teeth had penetrated.

I repeated this experiment on two rabbits, and on a guineapig. They lived beyond all compariſon longer than the fowl, and their life was not equivocal, as was eaſy to be diſtinguiſhed by the volun-

voluntary motions of the parts. It is true, that in theſe laſt I prevented the loſs of blood, in a great meaſure at leaſt, by tying the veſſels; and it is certain that they might have lived much longer, if the total effuſion of blood could have been prevented. The ſymptoms of the diſeaſe of the venom were manifeſt in all the three, and in all of them the muſcles that had been bit were livid.

This experiment ſhows, that the head of warm and perfect animals is not neceſſary to life, although it is ſo to a continuation of it. In a word, an animal may live very well, although deprived of its head, and may be ſenſible to external objects. The pulmonary reſpiration, and the circulation of humours in the parts, are ſufficient for this effect. This principle of life is ſtill ſuſtained in the animal, which may reaſonably be ſaid to be not altogether dead, but only ſo in part.

CHAPTER

CHAPTER VI.

On the Cauſe of the Death of Animals bit by the Viper.

My experiments on animals in which the nerves were bit by vipers, ſhow that the venom is a ſubſtance perfectly innocent to theſe organs, that it does not occaſion in them any ſenſible change, and that they are not even a means or vehicle of conveying it to the animal. In a word, it appears that the nervous ſyſtem does not concur more to the production of the diſeaſe of the venom, than does the tendon, or any other inſenſible part of the animal. On the other hand, all the experiments on the blood, the injection of venom into the veſſels, and ſo on, conſtantly evince that the action of the venom of the viper is on the blood itſelf. This fluid is alone changed by the venom, and this fluid conveys the venom to the animal, and diſtributes it to its whole body. The action of the venom, and its effects on the blood, are almoſt inſtantaneous. The colour of the latter is ſuddenly changed, and the bright red colour that is natural to it, becomes livid and black. This firſt effect is ſucceeded by a ſecond. The blood coagulates very ſuddenly in the lungs, heart, auricles, liver, and in the large venous veſſels. Sometimes the

the heart ſtill continues its oſcillatory motions, notwithſtanding the blood it contains is, at leaſt in part, coagulated. At other times, the heart beats with greater force, as if it wiſhed to ſtop the principle of coagulation that exiſts in the blood.

The coagulation of the blood of animals is certainly the moſt remarkable effect of the venom of the viper, and it is this which ought principally to occaſion the derangements in the viſcera, and in their functions. But the whole maſs of blood is not coagulated in the animal, ſince a part of it appears in a diſſolved ſtate. The red and lymphatick parts alone form the coagulum, the ſerous part is more fluid and diſſolved than before. It is certain at leaſt, that the latter is thrown in great abundance on the venomed parts, and ſheds itſelf in great plenty on the adipoſe membrane.

If the coagulated part of the blood is left for ſome time in water, it loſes the black colour it had contracted, depoſits the red part, which unites with the water, and leaves a tenacious, white, fibrous, ſubſtance, ſimilar to the polypus.

The blood, partly coagulated, and partly diſſolved, produces a very violent derangement in the organs of the animal. The part bit by the viper ſwells inſtantly, and becomes livid by ſucceſſive degrees. The blood in the large veins ſtops and coagulates. The ſerous part tranſudes into the adipoſe membrane, which it entirely fills. The circulation is deranged in the viſcera, diminiſhes by degrees, and at length ceaſes. The lungs are the

viſcus

viſcus in which the circulation ceaſes ſooner than in the other parts. In a moment after the injection of venom into the jugular vein, the blood coagulates in the lungs, the veſſels of which are filled and diſtended with this humour, in a black condenſed ſtate. In a word, the circulation is totally impeded and ſtopped, and the animal dies. It is a known fact, that as ſoon as the circulation is ſtopped in an animal with warm blood, death enſues in a few minutes, whatever the principle may be that binds and unites together the circulation and the life, the motion of the fluids and the ſenſitive faculty.

It will not be foreign to the purpoſe to ſpeak here of the animal irritability, or of that property of the muſcular fibres, by which a muſcle contracts on the ſlighteſt touch. We muſt conceive this property of the muſcular fibres, as ſomething that differs from the nerve, or from ſenſation; notwithſtanding it is true, that the nerve is the organ of the voluntary motions of the animal, and that when it is touched, it excites irritability in the muſcle. The nerve, in whatever way it is ſtimulated, is always motionleſs, and the muſcle continues to contract after it is ſeparated from the animal; whence it follows, that the nerve is rather the occaſion than the cauſe of the contraction of the muſcles.

In my work entitled *De Legibus Irritabilitatis nunc primum Sancitis*, printed at Lucca in 1767, I demonſtrated that the nervous fluid cannot be the *efficient cauſe* of muſcular motion. The arguments I adduced in that work are drawn from the hypo-

theſis, that the nervous fluid acts agreeably to the laws of fluids in general. If the nervous fluid was different from fluids in general, if it had laws altogether different to theirs, or if it was analogous to electricity, my reaſons would be no longer applicable to the preſent caſe.

However this may be, it is certain that the motion of a muſcle ſeparated from the animal, does in no way depend on the animal, or on the ſenſitive principle that reſides in it, and that the irritability in the fibres ſubſiſts from itſelf alone. The irritability of the fibres is therefore diſtinct from the ſenſibility of the animal, and two things which appear ſo different, and which ſeem to have been ſeparated by nature, ought no longer to be confounded.

But if this ſenſitive principle, which conſtitutes the life of the animal, is different from the irritability of the fibres, why, in a part ſeparated from the animal, may there not ſubſiſt an obſcure ſenſation, an imperfect life, relative to the ſize and to the nature of the part ſeparated from the animal, and to the nerves that are found in that part?

In this ſuppoſition, there is no agreement, no harmony, betwixt the life of the entire animal, and the obſcure ſenſation of the part that has been ſeparated; but I do not ſee why, in this caſe, the irritability may not likewiſe depend on the ſenſation of the part. The irritability would then depend on the partial ſenſibility, or would be the ſame as the latter, that is to ſay, it would depend on the ſenſibility

lity of the part cut, and not on the ſenſibility of the animal.

But the opinion that an obſcure ſenſation of life ſubſiſts in the parts ſeparated from animals, is founded on an immenſe number of obſervations and experiments, which I have promiſed to give in the ſecond volume of my Philoſophical Enquiries on Animal Phyſicks, *(Recherches Philoſophiques ſur la Phyſique Animale)* the firſt volume of which, in quarto, was printed in Italian at Florence, in 1775. In the mean time I can venture to aſſert, that I know a very great number of animals, even amongſt thoſe that are called perfect, that is to ſay, that have humours, heart, and viſcera, in which the hypotheſis of the continuance of animal ſenſation, in parts that have been divided, is verified.

But whatever opinion may be adopted on irritability, it is ſtill certain that this property exiſts in the muſcular fibres, that it is the principle of all the motions of the animal, and that without it, all would be ſtill, the organs would become uſeleſs, and the functions would be ſuſpended.

When I wrote the firſt part of the preſent work, I was of opinion that the venom of the viper attacked the irritability in an immediate way, and that the animal died from the loſs of irritability in the fibres. But I did not then know, that the venom of the viper has no action on the nerves, and that when it is introduced into the blood, it kills an animal in a few inſtants. This hypotheſis ought now to be partly modified. It is not that in effect the

irritability is not diminiſhed, in the animal that has been bit, and that it is not even deſtroyed in a little time; but this is rather an effect than a cauſe, and is a conſequence of the change cauſed in the blood by the venom, rather than an effect of the venom on the muſcular fibres. It ſometimes occurs that we ſee an animal, at the moment of its being bit, loſe all its voluntary motions, and ſcarcely diſcover any of the lateſt ſymptoms of life.

The debility of an animal, after it has been bit, is in general very great; but this ſhows equally, that the ſenſibility is affected: and as the venom does not act on the nerves, but on the blood, this diminution of ſtrength and ſenſation, and likewiſe the diminution of the irritability itſelf, may depend on the blood.

I have had frogs bit in the leg by the viper, and have found upon pricking the crural nerves a little time after the bite, or upon drawing electrical ſparks from them, that they had loſt but little, if any, of their irritability. It is very true, that this irritability diminiſhes with time, and that, on the death of the animal, it is frequently loſt altogether; but in theſe caſes, the ſenſibility is likewiſe diminiſhed and loſt. It is beſides certain, that if the crural nerves of the leg that has not been bit are ſtimulated, they contract with greater force than thoſe of the other; and that they frequently contract ſtill, when thoſe of the venomed leg have no longer in any degree that property.

The irritability of the fibres, in animals bit by the viper, diminiſhes in proportion as the diſeaſe is more conſiderable, and as it continues a longer time. An animal that dies in a few minutes preſerves in its muſcles more irritability, than one that dies at the end of ſeveral hours, or of ſeveral days. The irritability ceaſes much later in the heart, ſtomach, and inteſtines, than in the other parts. It particularly ceaſes very late in the inteſtines, which continue to move, notwithſtanding the animal has been dead ſome time. The irritability of the diaphragm, or the motion of the thorax, ceaſes much later than that of the other muſcles that depend on the will.

I made all theſe obſervations on animals with warm blood, in which it appeared to me, that the electrical ſparks were drawn with greater difficulty from the parts bitten, than from the other parts of the animal. This experiment ſucceeds particularly in fowls, in which there is no difficulty in laying the muſcles of the leg bare, and in having them bit.

The diminution of irritability in the muſcular fibres, is occaſioned by the changes the venom cauſes in the blood. The latter in this ſtate, in which it is partly diſſolved and partly coagulated, is diſpoſed to a ſpeedy putrefaction, and being pent up in the veſſels, diſſolves the texture of them, paſſes through their coats, and ſheds itſelf in the adipoſe membrane, corrupting and decompoſing whatever it meets with. In animals, the parts that have been bit by the viper, paſs in a ſhort time to

the

the ſtrongeſt putrefaction, and preſent gangrenes and ſphacelations. The ſkin is ſpeedily corroded and deſtroyed; the muſcles black and fœtid; and the adipoſe membrane diſſolved.

I have known a rabbit die in leſs than three hours, with the muſcles of the leg already gangrened throughout their whole ſubſtance; they were black and offenſive, and were divided by a knife without any reſiſtance. In a word, this putrefactive tendency of the muſcles, in animals that have been bit by the viper, cannot be denied, and is occaſioned by the change wrought in the blood by the venom.

It is very true, that when the animal dies in a few minutes, there is as yet no actual putrefaction in the ſolid parts, although the humours have a true tendency to this ſtate. The diſeaſe reſides ſolely in the humours, and the ſtoppage of theſe humours in their natural courſe, occaſions the death of the animal. Whatever tends to impede the motions in the animal machine, neceſſarily tends likewiſe to deſtroy in it the ſenſitive principle and life; and we cannot conceive life there, where every thing is in a perfect repoſe.

Senſation is an active principle, and neceſſarily expreſſes an action, and we cannot conceive action without motion. We ſay in effect, that an animal is dead, when it is no longer ſenſible; and we ſay, that it is no longer ſenſible, when there are no longer in its organs, the ſigns, the external motions, that indicate ſenſation. The moment theſe motions ceaſe, we ſay that an animal is dead. This manner

of judging is founded on obſervation itſelf. We have ſeen that when an animal is reduced to this ſtate of repoſe, it does not return again to life; and think, we may, on the other hand, reaſonably conclude, that an animal, when it is dead, can no longer revive in any manner whatever. This ſecond opinion, if we pay attention to it, actually appears to be derived from the firſt, ſince, after all, we do not know the principle that conſtitutes life and ſenſation in animals; it is, however, contradicted by obſervations and experiments of more modern date.

The obſervation that an animal deprived of motion does not return to life, appears to be combatted, as I have ſaid, by modern obſervations of a quite contrary nature. We have heard of ſtrong aſphyxies, in which there was no longer any ſign of motion. We are likewiſe told of drowned perſons, who have preſented the ſame phenomenon, although death in them was nothing more than apparent. I therefore do not ſee, why a certain obſcure motion may not ſubſiſt in the organs of an animal, which may not come within the reach of the evidence of our ſenſes. A motion to be inſenſible is not the leſs real; and when a motion ſubſiſts in an animal, there may ſtill ſubſiſt in it a principle of ſenſation.

I cannot deny but that, when there no longer ſubſiſts any principle of ſenſation, the animal is in all phyſical rigour dead; ſince we cannot poſſibly have any conception of life, in an animal without ſenſation. In the ſame way it ſeems equally clear, that

a total repoſe in the organs of an animal, ought to cauſe this ſenſation to ceaſe, and conſequently to occaſion the death of the animal. But is there any method by which we can aſſure ourſelves of the total immobility of the organs of an animal, in which the humours are ſtill in a fluid ſtate? I cannot conceive any one. A very ſmall motion is entirely imperceptible to us, and we ſee only the greater ones. Every thing in nature is in motion, and it is not poſſible that a body, or any of its parts whatever, can be found for a ſingle inſtant in a total and perfect repoſe. Perfect repoſe is beſides repugnant to the general laws of gravity, and to the nature of fluids, which are in a greater or leſs degree penetrated by heat. Hence ariſes the difficulty of pronouncing on the death of animals, ſince in ſhort there may ſtill ſubſiſt in them a motion which may be inſenſible to us, but which may yet be ſufficient to maintain in them an obſcure ſenſation, to prevent their being altogether dead, and to enable them to return to life.

The motion of the heart being ſuſpended, and the reſpiration and circulation ſtopped in an animal, it is ſoon reduced to that ſtate in which we ſay of it that it is dead; notwithſtanding that this may probably not always be the caſe, when we believe it to be ſo. I know of only two ſtates of an animal, that can make us certain of its being really dead. One of theſe is the total putrefaction of its organs; the other, the abſolute deſiccation of its humours.

The firſt renders all animal function impoſſible; the ſecond deſtroys all principle of motion.

The total deſiccation of the fluids and ſolids of an animal, not only forbids the uſe of the organs, but even conveys an abſolute immobility into all the parts. An animal, in this ſtate of a total deſiccation of parts, and of an immobility of its organs, is in my opinion certainly dead, and ought to be ſo in the opinion of every body; elſe we ſhould be expoſed to a capricious and unreaſonable pyrrhoniſm. A fiſh, for example, dried in the ſun, or by artificial heat, during twenty years, ſo as to become hard as wood, might ſtill paſs for being alive. I muſt confeſs that I cannot conceive life without action, nor action without motion, nor organical motion when the organs are dry; and this ſtate is therefore to me a ſtate of death. The naturaliſt ought not, however, to confound with each other theſe two different ſtates of death, that is to ſay, the putrefaction of the parts, and the deſiccation of the organs. In the firſt the animal is dead for ever; in the ſecond, it may yet again return to life. We do not know any power, nature herſelf does not diſcloſe any, that can recompoſe an organ that is deſtroyed, and entirely decompoſed by putrefaction, or by the concuſſions of external bodies. This is what has never yet either been accompliſhed or ſeen. We have therefore every poſſible reaſon, not only to believe an animal that is reduced to this ſtate dead, but likewiſe to believe it dead for ever. But if the animal is ſimply dry, if there is no phyſical diſeaſe

in its organs, if the component particles of the different parts ſtill preſerve their reſpective ſituations, the animal may in this caſe very well return to life, to which effect it is only neceſſary, that the organs are reſtored to the ſtate they were in when the animal was alive. And why ought not an animal to revive in theſe caſes, provided it has every thing that concurred to make it live before? Whoever had reaſoned in this way a century ago, would have advanced matters both probable and reaſonable, but would not have been liſtened to, even by philoſophers, and would have riſked the paſſing, at leaſt for an extravagant perſon, or for a viſionary.

But let us return to the animals that die by the bite of the viper.

The blood coagulates in the veſſels of an animal bit by the viper, and the animal itſelf is in a ſtate of death. The blood, changed by the venom, corrupts and deſtroys the organs of animals, and renders the leaſt ſuſpicion of life altogether improbable.

It is true that, in proportion as the circulation of the blood ſtops in the veſſels, and as the death of the animal approaches, we likewiſe ſee a perceptible diminution of the ſenſibility; but this does not yet demonſtrate to us, that the nerves are either changed, or have received an injury.

There may perhaps be ſuch an harmony or agreement betwixt the circulation of the blood, the air of the lungs, the principle of ſenſation, and the

nerves, that the one being removed, the other may diminiſh, although one may not act on the other.

My experiments have demonſtrated, that an animal may loſe its ſenſibility from quite another cauſe than from that of an injury to the nerves; and I therefore think that any one would reaſon ill, who ſhould ſay that the death of an animal depends on the nervous principle alone, becauſe in proportion as the animal draws towards its death, its ſenſibility is found to diminiſh. The diminution of ſenſibility in the nerves may be a ſecondary effect of the cauſe that kills an animal; and indeed, if the repoſe, if whatever puts a ſtop to motion in the animal, produces death, it muſt likewiſe be productive of a loſs of ſenſation, which cannot ſubſiſt without motion.

Such is the death of animals with warm blood bit by the viper; but in cold animals it is not exactly the ſame. Animals with cold blood, ſuch for example as frogs, may live a certain time without the circulation of the blood, and without reſpiration. It is preciſely on this account, that the venom of the viper operates on them with leſs activity than on warm animals, and that they ſurvive much longer than theſe laſt, in proportion to the ſize of their bodies. The action of the venom of the viper is inſenſibly communicated to the whole animal; the muſcles diſpoſe to putrefaction, and the part bitten becomes in a little time livid and gangrenous. The death of the animal then follows, but happens much later than in animals with warm blood, be-

cauſe

cauſe the principle of life is not ſo intimately connected with the circulation of the humours.

Why the circulation is thus cloſely connected with life in animals with warm blood, and why it is ſo little ſo in animals that have the blood cold, is a much nicer enquiry. I propoſe to ſpeak on this ſubject in a work *On Factitious and Natural Airs*, (*Sur les Airs Factices et Naturels*) which I hope to publiſh very ſoon.

End of the Firſt Volume.

INDEX

TO THE

FIRST VOLUME.

A

ACIDS the Mineral, do not effervesce with the venom of the viper - - - - - 254
——— they unite with it - - 255
——— they do not dissolve it when it is dried *ibid*
Acids the Vegetable, do not dissolve the dried venom of the viper - - - - - 256
Acid the Vitriolick, is not mortal to animals when introduced into their wounds - - - - 247
——— does not dissolve the venom of the viper when it is dried - - - - 255 259
Acidity the, of the Venom of the Viper, has been falsely supposed by several authours - - - 241
——— an acidity is really found in the venom of bees 266
——— but is not the cause of the swelling and inflammation of the parts - - - - 267
Adder the, is not killed by the venom of the viper - 39
Air fixed, the $\frac{1}{22360}$ part of a grain of this air discovers itself by the changes it causes in the dye of the turnesol 246
Airs, that may be obtained by fire, and by the nitrous acid, from the venom of the viper and gum arabick - 260
Alkalies, do not effervesce with the venom of the viper 254
——— do not dissolve it when it is dry - - 256

Alkali Volatile, owes its reputation to a falſe theory - 9
——— recommended in a late publication - 122
——— experiments on its effects againſt the bite of the viper - - - - - - 132
——— is not a ſpecifick againſt the venom of this animal to pigeons - - - - 138 142
——— nor to ſparrows - - 136
Amputation, of the gills of fowls, after the bite had been made in the comb - - - - 207
Anatomy, of the viper's head - - - 234
Animals in general, the venom of the viper occaſions their death, by an internal derangement it produces - 229
Animals with warm Blood, are all ſenſible to the deadly effects of the venom of the viper - - 41 272
——— cauſe of their death - 297
Animals with cold Blood, live a long time without heart, and without viſcera - - - - 106
——— are not all ſuſceptible of the effects of the venom of the viper - - - - 275
Animals venomous, are probably not venomous to their own ſpecies - - - - - 32
Aſpick of the environs of Piſa, is nothing more than an innocent ſnake - - - - 38

Bag, or *Sheath*, *that covers the Teeth of the Viper*, its deſcription - - - 8 235
——— it is not the receptacle of venom - - 234
Baker aſſerts, that the venom of the viper cauſes a diſſolution of the humours - - - - - 79
Bees, experiments on their venom - - 263
—— it, in ſeveral reſpects, reſembles that of the viper 264
—— but differs from it in changing the juice of the ſkin of radiſhes red 266

Bite

Bite of the Viper, in the breaſt of fowls more dangerous than in the legs, contrary to what was obſerved in rabbits and guinea pigs - - - - - 191
——— not very dangerous on the ears of rabbits 197
——— more dangerous in the gills than in the comb of fowls - - - - 206
——— leſs dangerous on the noſe of quadrupeds than elſewhere - - - - - 214
——— its effects on limbs recently cut off 289
——— on limbs in which the circulation had been ſtopped - - - - - 371
——— on limbs ſheltered from the air - 391
——— on parts cut off after the circulation had been interrupted by a ligature - - 392
——— on frogs, the head of which had been previouſly cut off - - - 361
——— on frogs, the ſpinal marrow of which had been divided - - - 362
——— on rabbits, in the ſame predicament 369
——— on animals with warm blood, deprived of the head - - - - - 394
——— has no effect on the tendons - 231
Blood globules of, errours of writers in regard to their figure 80
Blood, whether it loſes any of its principles when in contact with the air - - - - 309
—— when drawn from the veſſels does not ſeem at all changed by the mixture of the venom of the viper 313
—— is found coagulated in the heart, &c. of animals killed by this venom - - - - 319
—— Mead thought at firſt, that the blood was the *medium* through which the action of poiſons rendered itſelf general - - - - - 89
—— the blood of cold animals is acted on in the ſame way as that of the animals in which it is warm - 384
—— that of the viper excepted - - *ibid*
Bones, not acted on by the venom of the viper - 198

Bonguer, his account of the American ſnake, that may be dried in the air, or in the ſmoke of a chimney, and afterwards reſtored to life by means of water - 112
Brain, inſenſible to the action of the venom of the viper 199
Brogiani, his Treatiſe on the Venoms of Animals - 76
Buffon Monſ. de, his idea of the action of the venom of the viper - - - - - 86
——— of the nature of the pus of wounds 87

C

Cats, the conſequences of having them bit on the noſe by the viper - - - - 216
——— experiments with the volatile alkali on theſe animals 153
Cavity, new one diſcovered in the canine teeth of the viper 10
Charas, his opinion of the venom of the viper - 158
Cock, its gills ſwell when it has been bit in the comb 203
Convulſions, by what cauſes they may be excited 80 81
Cornea tranſparent, the venom of the viper produces no change in it when ſimply applied to its ſurface - 201
——— but when introduced by a wound, produces a leucoma and ſwelling - - *ibid*

D

Death, definition of life and death - - - 115
Diſeaſe produced by the Bite of the Viper, the ſymptoms that characterize it - - - 305
——— conſidered by Mead as a nervous one - - - 330
——— the circumſtances that render it more violent - - 164
——— no appearance of diſeaſe in the legs cut off immediately after having been bit by the viper - - - - 390

Dogs and *Cats*, effects of the volatile alkali on these animals after they have been venomed by the viper 153 155 172 173
Drones, experiments on the wounds made with the sting of these insects - - - - 265
Dura mater, not acted on by the venom of the viper - 199

E

Ears, effects of the bite of the viper on these parts - 194
Earth-worm, very speedily killed by the polypus - 100
Eau de luce, the ingredients that enter into its composition 121
Eels in smutty Wheat, dried till they were dead, and restored to life by humudity - - - 112

F

Foam of the Viper, is not venomous - - 25
Fowls, the uncommon disease they are attacked by when bit in the comb by a viper - - 202
——— they die when bit in the breast - - 191
——— although the disease is not so dangerous, they likewise die when bit in the leg - - *ibid*
——— in the last case they die much later than pigeons 144
——— experiments to determine the effect of the volatile alkali, as an antidote, on these animals - 143 166
——— the absolute inutility of it - - 145
Frog, the venom of the viper destroys in it the principle of irritability - - - - - 97
——— experiments that prove the inefficacy of the volatile alkali, in the bites received by this animal - 157
——— it is killed by the bite of the viper - 273
——— effects of the venom of the viper on the brain and nerves of this animal - - - 332
——— it is very soon killed by mephitick vapours - 91

G

Gills of Fowls, the diſeaſe produced in them by the bite of the viper - - - - - 205
——— the cutting them off does not prevent the effects of the bites of this animal, made in the comb 208
Gland, what appears to be ſuch, and to be deſtined to the ſecretion of the venom of the viper - - 235
Gordius, a worm that recovers its life, if after having been dried it is wet with water - - 112
Guineapigs, experiments with the volatile alkali on theſe animals, which determine its efficacy againſt the bite of the viper - - - - 147 148
——— the venom of the viper ſimply introduced into the muſcles of theſe animals does not act on them ſo as to occaſion death - - - 186
——— the bite of the viper on the breaſt of theſe animals very dangerous - - - 190
——— conſequences that reſult from their being bit on the neck - - - - 208 209
——— ——— ——— from their being bit on the noſe - - - - 213
Gum Arabick, its general characteriſticks compared with thoſe of the venom of the viper - - 258
——— is perfectly innocent when applied to wounds 261
——— effects of this gum on blood drawn from the veſſels of an animal - - - 386
Gums, their characteriſtick properties - - 256
——— the venom of the viper is a ſpecies of gum - 257

H

Head of the Viper, its deſcription - - 234
Heart, that of the wheel-polypus in a voluntary muſcle 110

I.

James, Dr., believed in the acidity of the venom of the viper 240
——— the errours of this writer as to the ſeat of the reſervoir of venom - - - 22
Jaundice that ſupervenes in thoſe that have been bit by the Viper, the cauſe of it explained - - 83
Injection of the Venom of the Viper into the Blood-veſſels, and its Effects - - - - 315
Inſects that wound with a Sting, introduce into the wounded part, a liquor of a gummy nature - - 265
Inteſtines, effects of the venom of the viper on theſe viſcera 192
Irritability, deſtroyed by mephitick vapours - 93
——— ——— by the venom of the viper - 98 102
——— it is rather one of the circumſtances that attends it, than the cauſe of the death of animals bitten - 401
Juſſieu, recommended the uſe of the volatile alkali againſt the bite of the viper, in conſequence of deeming the venom of this animal acid - - - 244

L

Leeches, employed againſt the bite of the viper - 36
——— deſcription of their mouth - - 72
Leg, the part of animals on which the greateſt number of experiments were made with the venom of the viper—what ought to be underſtood by this expreſſion - 132
Lewenhoeck, his errour in relation to the ſuppoſed wheels, or arms of the wheel-polypus - - 108
——— motion of the heart of this animal, which appears to be a voluntary muſcle - - 110
Ligaments, the venom of the viper has no effect on them 223
Limbs, bit by the viper, and cut off immediately after, do not appear to be changed - - - 389
Liver, effects of the bite of the viper on this viſcus - 193
Lungs, of animals bit by the viper, are covered with livid ſpots - - - - 322
Lymph, its characteriſticks - - - 250

M

Marrow of the Bones, not acted on by the venom of the viper - - - - 209
Mead, this authour is the firſt who has written on the nature of the venom of the viper - - 239
—— he believes this venom to be hot and acrid to the taſte 252
—— ſuppoſes the exiſtence of certain ſalts - 47 247
—— he ingenuouſly agrees, at length, that it is neither acid nor alkaline - - - - - 254
—— his hypotheſis, as to the action of this venom 78 244
—— remarks, that it produces no change in blood drawn from the veſſels - - - 312
—— believes that it acts on the animal ſpirits - 89
—— has a dog bit in the noſe by a viper—the concluſions he draws from thence - - - - 211
—— the drawings he gives of the teeth, &c. of the viper, are defective - - - - 234
—— his opinion of the venom of bees - 263
Membrane adipoſe, experiments made with the venom of the viper on this part - - - 183
Mephitick vapours, different hypotheſes on their mode of action on animals - - - - 90
Morgagni, his opinion of the pain excited by the ſting of the ox-fly - - - - 69
Muſcle the conſtrictor, of the veſicle of Venom of the Viper, its deſcription - - - - 21
Muſcles, action of the venom of the viper on theſe parts 185 187

N

Napel, effects of this juice on the fleſh of animals - 101
Narcoticks, have no effect on dogs - - 35
Neck, bite of the viper on this part - - - 208

Nerves, effects of the venom of the viper on theſe organs 330
——— on the ſciatick nerve of rabbits - - - 337
——— on the ſame nerve divided above the bite - - 343
——— divided below - - - 345
——— with a ligature, without being cut - 349
——— compurative experiments, by making ſimple mechanical wounds on the ſciatick nerve - - 352
Nintipolenga zeilanica, mode of action of the venom of this ſnake on the animals it bites - - 105
Noſe, effects of the bite of the viper on this part - 210
Nicholls, his drawings of the head of the viper imperfect 234
——— his idea of the flowing of the venom from the point of the canine teeth of the viper - - 6

O

Oils eſſential, are not diſſolved by the venom of the viper 256
Opium, why it cauſes convulſions - - 81
——— its effects are very analogous to thoſe of the venom of the viper - - - - - 114
Orvai, a ſnake that reſiſts the bite of the viper - 39
Ox-fly, (the *Taon* of the French) ſeems to be the ſame as the Oeſtros of the Greeks, the Aſilus of the Latins, and the Tabanus of Varro and Pliny - - 70

P

Palſy, cauſed by the bite of the viper - - 99
Particles organical, falſely ſuppoſed in the venom of the viper, and likewiſe in the pus of wounds - 87 83
Pericranium, not acted on by the venom of the viper 197
Perioſteum, not acted on by the venom of the viper - 198
Pigeons, the quantity of venom required to kill a ſparrow, has in general very little effect on theſe animals - 284
——— it requires about four times as much to kill them 285

Pigeons, a ſingle viper can ſucceſſively kill ten, or even twelve of theſe animals - - - 161
——— they generally die within from eight to twelve minutes after they have been bit - - 141
——— the volatile alkali is of no uſe to them 138, 142
——— effects of the bite of the viper on their breaſt 189
——— on their pericranium - - 197
——— bones and perioſteum - - 198
——— duræ mater and brain - - 199
——— marrow of their bones, &c. - - 200
——— on their muſcles - - - 185
——— muſcular fibres - - - 187
——— the amputation of the leg does not occaſion the death of theſe animals - - 298
Poiſons of the animal kingdom; their mode of action 106
Polypus of freſh water; its venom is analogous to that of the viper - - - 100
——— it is the moſt active of all venoms 106
Principle; an unknown one that ſeems to exiſt in the blood of a living animal - - - 327
Pus of wounds, improperly thought to contain live inſects 88
Putrefaction is the real characteriſtick of death - 405

R

Rabbits; effects of the venom of the viper on their muſcles 185
——— its effects on the tranſparent cornea of theſe animals 200
——— the bite of the viper on the breaſt of theſe animals very dangerous to them - - - 190
——— the bite made on their belly occaſions a tumour 191
——— effects of the bite on their inteſtines - - 192
——— on their liver - - - 193
——— on their ears - - - - 194
——— neck - - - 209
——— noſe - - - - 211
——— on their tendons, ligaments, &c. - 221

—— on their nerves - - 337
—— effects that enſue from the injection of the venom of the viper into the blood veſſels of rabbits 318, 325
—— inefficacy of the volatile alkali, in checking the effects of the venom in theſe animals - 149, 171
—— it rather tends to encreaſe than diminiſh the diſeaſe of the venom - - - 152
Radiſhes; their ſkin affords a blue ſtain, very ſenſible to the action of the ſlighteſt degree of acidity - 242
Rage or fury of the viper does not render its venom more active - - - 25, 26
Reaumur; his opinion of the pain produced by the ſting of the ox-fly - - - 69
Receptacle of the venom of the viper; its deſcription 21, 235, 236
Redi; the firſt who had any determinate ideas as to the venom of the viper - - 238
—— his errour in regard to the receptacle of the venom in the viper - - - 6, 235, 239
—— his opinion of the paſſage of the venom along the outer part of the tooth - - - 5
Rotifer, or wheel-polypus; its wheels are not real, but apparent - - - 108

S

Saline net-work, which Mead thought he ſaw in the venom of the viper; what it is - 50, 247, 249
Saliva of the viper is not venomous - - 85
Salts; Mead thought he could perceive ſalts in the venom of bees - - - 263
—— there are no ſalts in the venom of the viper 54, 102
Scorpion; miſtake as to its deſtroying itſelf with its own venom - - - 33
—— authors do not agree as to the number of orifices in its ſting 64

Skin; experiments to determine the action of the venom of the viper on this part - - - 177
Slugs are not destroyed by the venom of the viper 38
Snails are insensible to the venom of the viper - 38
Sparrows; the time in which they die after they have been bit by the viper - - - 141
——— the quantity of venom required to kill them 285
——— effect of the volatile alkali on these animals bit by the viper - - - 133
Spirit of wine does not dissolve the venom of the viper 257
——— precipitates it after it has been dissolved in water - - - 258

T

Tartar stibiated; has no action on the eyes - 34
Taste, hot and acrid, falsely supposed by Mead in the venom of the viper - - - - 56
Teeth canine of the viper; their description - 9, 236
——— their number - - - 7, 237
——— their double channelling - - 10
Tendons; experiments made on them with the venom of the viper - - - 220
——— in these experiments the death of the animal is occasioned by the denudation of the tendon, and not by the venom - - - 231
——— a reproduced vascular substance covers the tendon after it had been laid bare, and restores it to its former state - - - - 232
Tooth venomed; what it is - - 185
Toxicodendron; microscopical examination of the juice of this plant - - - - 102
Tumour that succeeds to the bite of the viper in the belly, in rabbits and guinea-pigs - - 191
——— it likewise appears when they are bit in the ear 194
——— and in the nose - - 211

——— fowls that have been bit in the comb have a tumour in the gills - - - 202
Tunica conjunctiva; not acted on by the venom of the viper 57
Turtles; are not ſuſceptible of the venom of the viper 40

V

Valiſnieri; his opinion of the paſſage of the venom 6
Venom of the Viper flows out at the elliptical hole at the point of the canine teeth - - 16, 23
——— manner in which it is to be employed in experiments, to obtain uniform conſequences 141
——— it has no taſte - 55, 252
——— when liquid, unites with the mineral acids without efferveſcence - - 255
——— the nature of it - - 238
——— it was thought acid by Mead 240
——— is not ſo - - 45
——— neither is it alkaline - 254
——— it diſſolves in water, but not in ſpirit of wine - - - 257
——— does not melt in the fire - 258
——— microſcopical examination of a drop of this venom - - - - 48, 259
——— it is a poiſon to every ſpecies of animals with warm blood - - 272
——— has no action on leeches - 36
——— nor on ſnails - - 38
——— a determinate quantity of it is required to kill an animal - - - 277, 287
——— a thouſandth part of a grain in weight is ſufficient to kill a ſparrow - - 285
——— it requires a certain time to produce its effect - - - 288, 294, 299, 302
——— it is not mortal if it does not penetrate beneath the adipoſe membrane - - 185

——— its action on parts bitten - 271
——— it has in general no action on the muſcles of animals - - - 186
——— nor on the bone, perioſteum, and pericranium - - - 197, 198
——— nor on the dura mater and brain - 199
——— nor on the marrow of the bones 200
——— nor on the tranſparent cornea 57, 200
——— has no effect on the tongue of rabbits 201
——— nor on amputated limbs - 289
——— when injected into the blood-veſſels, it kills an animal inſtantly - - - 318
——— cauſes no change in the blood drawn from the veſſels - - 313
——— prevents the coagulation of it *ibid.*
——— has no effect on the nerves - 356
——— according to Mead, it acts on the animal ſpirits - - - 330
——— does not loſe its deadly qualities, after having been employed in killing animals - 188
——— nor after having been kept a long time 65
Venom of the polypus analogous to that of the viper 100
Venomed; what ſhould be underſtood by this word 98
Vipers are not always provided with venom - 140
—— are not deſtroyed by their own venom - 29
—— it probably requires nearly the venom of two vipers to kill a man - - - 287

W

Waſps; experiments on their venom - - 263
Water diſſolves the venom of the viper perfectly 257
Wheel-polypus; its wheels are not real, but apparent 108

Digitized with financial assistance from the
Government of Maharashtra
on 18 July, 2018